

**A NARRATIVE REVIEW OF *Libidibia ferrea*:
BOTANICAL ASPECTS,
ETHNOPHARMACOLOGICAL PROPERTIES,
PHYTOCHEMICAL CHARACTERISTICS,
TOXICITY, AND EXPERIMENTAL TESTS**

ABSTRACT

Introduction: Jucá or pau-ferro (*Libidibia ferrea*) is an arboreal plant from the Fabaceae family. It is commonly used in traditional medicine in the treatment of various diseases, including inflammatory process.

Aims: The objective of this narrative review is to present botanical aspects, ethnopharmacological properties, phytochemical characteristics, toxicity highlighting, and experimental models with *L. ferrea*.

Results: **Botanical Aspects:** Jucá has several uses such as in landscaping (stem and canopy), in arborization of urban areas. **Ethnopharmacological Properties:** It is used in the treatment of various diseases such as diabetes, flu, asthma and, inflammatory processes of which different parts are used (root, stem bark, leaves, fruits, seeds).


Phytochemical Characteristics: Phenolic compounds, fatty acids, and terpenoids are among the compounds monthly used. **Toxicity:** *In vivo* models have been used to verify toxicity and in most studies the plant presented no toxicity in its use.

Experimental studies: Animals, such as mice, dogs, rats, etc. and different models of studies to analyze the action of the plant were used.

Conclusions: Such low toxicity, associated with its widespread use in folk medicine and its various effects demonstrated in the studies included in this Review have corroborated for the continuity of the research with *L. ferrea*. New studies, however, ought to follow methodological guidelines, such as the Animal Research: reporting *in vivo* Experiments (ARRIVE) so that, a methodological design secures more homogeneous studies capable of quantifying the actual size of the effect in the plant may have in clinical studies.

Keywords: Jucá, Medicinal plant, Animal experimentation.

GRAPHICAL ABSTRACT



1. INTRODUCTION

Brazil features a wide biodiversity of flora (20 to 22% of the world's total namely, 45 000 plants species) with pharmacological potential, but many of these plants have not yet been well studied to become targets of clinical studies [1].

According to the "Formulário de Fitoterápicos Farmacopeia Brasileira" [2], a medicinal plant is defined as a "plant species, cultivated or not, used for therapeutic purposes". Due to their great biological and chemical diversity, medicinal plants have been widely used for the treatment of various diseases, besides having a wide range of biological active compounds [1,3].

More than 35 000 species of medicinal plants can be found in the Amazon, of which approximately 5.000 have a great economic potential, not only for use in humans, but also in animals and environment. Popular knowledge observation about medicinal plants provides the opportunity to obtaining their active substances indicating the way to go with respect to biological activities [4]. In many developed countries the large part of the population, in primary care, depends on traditional medicine [5].

Approximately 25% of circulating drugs directly or indirectly derive from medicinal plants [6]. New substances of plant origin have been sought for the development of new medicines [1,7]. Among the plants associated with medicinal use, *Libidibia ferrea* (Mart. ex Tul.) L.P. Queiroz is one of them.

Libidibia ferrea (Mart. ex Tul.) L.P. Queiroz was included by the Ministry of Health (MS) in the National List of Medicinal Plants (SUS-RENISUS) in February 2009 [8,9]. Given the above, this study aims to fully review the literature on the species *Libidibia ferrea*.

2. BOTANICAL ASPECTS

Libidibia ferrea was designated by Carl (Karl) Friedrich Philipp von Martius (Mart.) in 1828 as *Caesalpinia ferrea* (basionym) [10]. The genus named after the Italian botanist Andrea

Caesalpinia, described by Carl Linnaeus [11] however, was described in 1844 by Louis René ('Edmon') Tulasne (Mart. ex Tul.) [10]. It suffered a taxonomic genus change from *Caesalpinia* to *Libidibia* by Lewis in 2005 and Luciano Paganucci de Queiroz [(Mart. ex Tul.) L.P. Queiroz], in 2009, allocated all variations of *Caesalpinia* to the genus *Libidibia*, named *Libidibia ferrea* (Mart. ex Tul.) L.P. Queiroz [12, 13] (Fig. 1).




Figure 1: *Libidibia ferrea* (Mart. Ex Tul.) L.P. Queiroz timeline.

Libidibia ferrea presents the following variations (var.): *ferrea* and *glabrescens*, with distribution in the Caatinga domain; *leiostachya* and *parvifolia* in the Atlantic Forest (Benth.) with differentiated distributions [12]. Regarding its taxonomic classification it belongs to the Plantae Kingdom, Magnoliophyta Phylum (Angiospermae), Magnoliopsida Class (Dicotyledonea), Fabales Order, Fabaceae Family, Caesalpinieae subfamily (Caesalpinoideae, Leguminosae), *Libidibia* genus, and *Libidibia ferrea* (Mart. ex Tul.) L.P. Queiroz species [10,14] (Fig. 2).




Figure 2: *Libidibia ferrea* (Mart. ex Tul.) L.P. Queiroz taxonomic classification.

Libidibia ferrea can be found in the phylogeographic domains of the Caatinga, Atlantic Forest, and Brazilian Cerrado [15]; in the Northern region [16] and in Northeastern region

[17]. The name *ferrea* is related to the hardness existing in its wood [15]. It is known by popular names such as jucá, pau-ferro or true pau-ferro and by its indigenous names: imirá-itá, ibirá-obi [11, 18], muiré-itá, muirá-obi [11], jucaína [19].

Jucá is an arboreal plant, heliophile, native to Brazil that can reach up to 15 meters (m) in height [11]. It features a lush and wide canopy [15]. As for its morphology, this species has a hard core, smooth trunk [11] and squamous [15, 19], with 40-60 centimeters (cm) in diameter [18].

Galdino et al. (2007) have collected leaves, fruits, and seeds in the city of Manaus and in parallel planted the seeds for monitoring their germination (average of 14 days) and the jucá seedling. Through biometric measurements the leaf of the adult plant was estimated in 7-30 cm and the seedling in 3-5 cm [20].

The fruit presented the following dimensions 8.3 x 1.8 x 0.8 cm (length x width x thickness) and 5.27 g (3.55-7.30 g) of weight. The fruits could be green and brown in color when immature and mature, respectively. It is considered an indehiscent fruit since no seed is released when the fruit is ripe. The fruits base can be rounded to curved, and it has a protruding ventral suture and the apex is rounded with mucro (oval) [20].

The flowering period occurs towards the end of November [19] exhibiting small flowers in the form of yellowish brunches [20] the smallest petal has red stains [17], several fruits coming from the inflorescence [20].

The seeds are found in individual cavities inside the fruit evident in a transverse and uniseriate arrangement of 6-12 seeds per fruit. Seeds have an average size of 0.9 x 0.5 x 0.5 and 0.15 g weight, light green to yellowish color, firm in its consistency and slightly wrinkled integument, discoid to ovoid shape, with rounded apex and flattened base [20], which are responsible for the species propagation [21].

L. ferrea has several applications, such as civil construction [11] for its quite heavy and hard wood, but it is also popular as fodder for cattle [22], used in urban landscaping and reforestation [15, 18], planks, fences, firewood, in animal feed [23,24], in the fight against gastrointestinal parasites, such as sheep parasites [14,25], ornamental species (avenues and streets) [11,18] and its phytotherapy used [23,24].

In addition to the above applications studies presenting its ethnopharmacological characteristics are also developed.

ETHNOPHARMACOLOGICAL PROPERTIES

Herbs, leaves and roots ingestion to cure diseases and relieve illnesses is possibly the first form of plant use [27]. And for *L. ferrea* the biological activities have been researched and many authors have used aqueous extract to verify the biological activities of the fruits [26,28-34], leaves [35], stem bark [26,36,37] and seeds [38,39].

Libidibia ferrea is one of the species that stand out in this use in folk medicine in the form of "garrafada" from stem bark, for the treatment of dysentery, diarrhea and anemia [40], in the use of leaf and fruit for the therapeutic treatment of hypoglycemia in the form of infusion and *in natura* [41], in the use of the left soaking for the treatment of fruit bronchitis and influenza [42].

In an ethnobotanical survey with residents around Serra da Capivara National Park in Piauí conducted by Reis et al. (2017) the use of *C. ferrea* leaves, pods, stem bark, root, and whole plant for the treatment of influenza, injuries, action on the liver, lungs, heart, throat, and as anti-inflammatory was reported [43].

Several parts of this plant are used in Amazon region. Leaves, in decoction, are used in the treatment of hemorrhoid (externa use), and in amebiasis and liver problems (internal use); treatment of tuberculosis with infusion of leaves with the fruits; in the form of syrup in the treatment of bronchitis and asthma [11]; use of the pods in the form of syrup, tea, and

infusion for the treatment of gastric problems [44]. The use of fruits immersed in alcohol is used in the Lower Amazon in the treatment of several dermal wounds [45].

Investigation of the action from *L. ferrea* is carried out in the in the treatment of analgesic and inflammatory conditions [27,29,46]; cancer chemopreventive [47,48]; larvicidal activity against *Aedes aegypti* and presenting cellulosic, anticoagulant and amylase activity with the use of the crude aqueous extract of the seed [38]; antimicrobial activity [33,36,49,50]; antiglycemic and the treatment of diabetes [37,51,52]; wound healing potential [45,53,54]; antioxidant and hepatoprotective, viral activity against the Herpes Simplex Virus e Poliovirus [55] and against the dengue virus (DENV-2) [56]; repellent action against species of the Calliphoridae family [57].

Other activities are performed by this plant: anthelmintic [25]; antileishmania action [58]; gastroprotective and antiulcerogenic [44]; cosmetic anti-whitening and anti-wrinkle potential effects as cosmetic [59]. In a study conducted with zebrafish (*Danio rerio*) that oral use of alcoholic extract can be used as an oral drug with an acceptable safety was observed [60]. Cutaneous treatment of wounds in goats was also observed in veterinary medicine with the use of stem bark as the basis for ointment production [61]. And there is a potential use for wound healing in dogs in a formulation containing 5% of jucá ethanol extract [45].

PHYTOCHEMICAL CHARACTERISTICS

Among various applications of medicinal plants extracts the anti-inflammatory action which has reports of some compounds, such as flavonoids, terpenes and, phenolic compounds [1] is highlight. In *L. ferrea* phenolic compounds the presence of gallic and ellagic acids, catechins and, epicatechins presence in aqueous extract from the stem bark [37] and bark [36].

Ueda et al. (2001) on analyzing the dried fruits of *C. ferrea* have observed the presence of ellagic acid and 2-(2,3,6-trihidroxy-4-carboxyphenyl) ellagic acid [62]. Identification of ellagic and gallic acids in fruits was also identified [63]. Studies have corroborated with the identification of steroids, phenolic compounds, saponins, coumarins, flavonoids and tannins in the hydroalcoholic extract from leaves and stem bark of *C. ferrea* [64]. Silva et al. (2013) have identified gallic acid and methylated gallate derivative from hydroalcoholic extract from the fruits [65]. Phenolic compounds, such as gallic acid and methyl gallate have been isolated from ethyl acetate extract from jucá fruits [48].

Frasson et al. (2003) upon analyzing the crude ethanolic extract of the stem of *C. ferrea* (Mart. ex Tul.) var. *leiostachya* Benth have observed the absence of saponins, low content of total tannin, presence of flavonoids in greater quantity in the ethyl acetate fraction, terpenes in petroleum ether and dichloromethane fractions [66].

Sawada et al. (2004) have observed the presence of different fatty acids such as palmitolenic, oleic, linoleic, linolenic, stearic, capric, palmitic in the lipidic portion seed [39]. Corroborating the data from Dias et al. (2013) who have identified by supercritical extract unsaturated fatty acid (52%), saturated fatty acids (26%) and terpenoids (13%) in pods of *L. ferrea* [53]. The presence of monosaccharide compounds (D-galactose e D-mannose) was observed at *C. ferrea* sulfated polysaccharide from seeds [55].

Comandolli-Wyrepkowski et al. (2017) have identified high levels of phenolic compounds and flavonoids from *L. ferrea* fruit and identified terpenoids in the plant [58]. Prazeres et al. (2019) have also identified phenolic compounds such as gallic acid and ellagic acid in the *Libidibia ferrea* dry extract from fruits [44]. Phenolic compounds, tannins and flavonoids have been identified in the lyophilized extract of the pods used as topical phytopharmaceutical [45]. Hassan et al. (2015) have identified carbohydrates and/or glycosides, tannins, and phenolic compounds in aqueous ethanolic extract of *C. ferrea* [51].

A study testing four different types of jucá extract have identified phenolic compounds, and carbohydrates (aqueous ethanol extract), lipids and predominance of organic acids (ethyl acetate extract), organic acids and predominance of lipids (chloroform extract), and alcohol and lipids (hexane extract) [67].

Table 1 presents the most compound observed in the articles included in this narrative review.

Table 1: *L. ferrea* compounds identified in the included studies.

Compounds		References	
Phenolic compounds	----	----	
	Hydrolysable tannins	gallic acid	[51], [58], [64], [67] [36], [37], [44], [48], [63], [65]
		ellagic acid	[37], [44], [62], [63]
		methyl gallate	[48], [65]
		2-(2,3,6-trihydroxy-4-carboxyphenyl) ellagic acid	[62]
	Condensed tannins	catechin	[36], [37]
		epicatechin	[37]
		tannins	[45], [51], [64]
	Polyphenols	Total tannins	[66]
	Flavonoids	----	[45], [54], [66]
Terpenes	Triterpenoid	----	
	lupenone	[53]	
Steroids and Lipids	----	[45] and [67]	
Fatty acids	Unsaturated	linoleic acid	[39], [53]
		oleic acid	[39]
		palmitolenic acid	
		elaidic acid	[53]
	gamma-sitosterol		
	Saturated	palmitic acid	[39], [53]
		stearic acid	
capric acid		[39]	

Carbohydrates	----	----	[51], [67]
Glycosides	----	----	[51]
	saponin	----	[58], [64]
Monosaccharide	D-Galactose, D-Mannose	----	[55]
Coumarin	----	----	[64]
Organic acids	----	----	[67]
Alcohols	----	----	

An observation about these compounds is that they vary according to the time of year, age of the plant, type of soil, climate, among others [66]. Among this diversity of existing compounds in medicinal plants it is known that they may or may not present toxicity. The evaluation of the toxicity degree is necessary since it may make the use of the plants impossible despite the plant showing a medicinal effect [68].

TOXICITY

The degree of plant toxicity might be related to the method of its preparation as well as the part used and the route of administration [41,69]. Few studies have been described regarding the toxicity of *L. ferrea* [70].

Souza et al. (2006) have investigated the effects of aqueous extract of the fruits on red bone marrow using micronucleus model and chromosomal aberration in Wistar rats. Results have shown that there was no cytotoxic or clastogenic effect [71]. Reboredo et al. (2007) have demonstrated that the use of this extract had shown alteration only in the weight of the seminal vesicle but had not alter the weight of the other organs of the male reproductive system of male Wistar rats in the subacute toxicity tests, used at the dose of 300 mg/kg once a day intragastrically for five days [34].

Study in female Wistar rats (3 months, 160-190 g) during the blastocyst implementation period (5th and 7th days of pregnancy) 300 mg/kg by gavage of *C. ferrea* aqueous extract of the fruit was applied the presence of toxicity in rats or into blastocyst implantation [32]. Cavalheiro et al. (2009) have also tested acute toxicity in six male Swiss mice and found that the crude aqueous extract of seeds applied intraperitoneally route at a dose of 0.3 mL/10 g showed no toxicity, weight loss, diarrhea, or behavioral changes [38].

Using 300 mg/kg of the aqueous extract applying 1 mL per gavage in a 52-day treatment caused no toxicity in Wistar rats [31]. In an acute oral toxicity test using aqueous extract and F80 fraction of *L. ferrea* var. *parvifolia* (Mart. ex Tul.) L.P. Queiroz at dose 2,500 mg/kg in female albino Swiss mice caused low toxicity [30].

No death of any animal has been recorded in the acute toxicity test of the ethanolic extract from pods. The test was performed in female Swiss mice with the application of a single dose of 2,000 mg/kg orally [46]. An *in vitro* toxicity test with macrophage RAW 264.7 cells using supercritical extract of raw fruits of jucá was carried out. And through the release of lactate dehydrogenase (LHD) it the induction of certain toxicity in these cells at the beginning of their release in the dressings was observed. However, there were no significant morphological changes in cells, led to the inference that the extract would not be toxic, but it

had inhibited the adhesion of fibroblasts [53]. In the study with *C. ferrea* (Mart. ex Tul.) var. *ferrea* using fruits/seeds low toxicity in HEP-2 ATCC CCL-23 cells was verified [55].

In 2015, a study using ethanolic aqueous extract from *C. ferrea* Martius leaves in Sprague-Dawley rats applying the dose of 1.500 mg/kg, no animal death or toxic reaction were observed, and much less mood change in animals until the end of the experiment [51]. Kobayashi et al. (2015) have tested oral toxicity at a dose of 5 g/kg using ethanol extract from *L. ferrea* (Mart. ex Tul.) var. *ferrea* fruits in 10 female Wistar rats and showed no acute toxicity [54].

Through the cell viability assay, at a concentration of 20 µg/mL of the dry extracts of the stem bark and pods, no cytotoxicity was significantly observed in normal human fibroblast cells [59]. Comandoli-Wyrepkowski et al. (2017) have observed low toxicity in J774 macrophages cells using methanol extract of fruits used to make gel in the topical treatment against infection with promastigotes and amastigote of *Leishmania* (*Leishmania*) *amazonensis* and *Leishmania* (*Viannia*) *guyanensis* in golden hamster (*Mesocricetus auratus*) [58].

Cunha et al. (2017) in a preparation of jucá seed gum containing polysaccharides (galactomannan) have observed, *in vitro*, through the LDH test in human leukocytes that there was no toxicity in the plasma membrane of neutrophils [72]. Guerra et al. (2017) using raw extract from the fruits of *L. ferrea* (Mar. ex Tul.) L.P. Queiroz var. *ferrea* tested cell viability in HT-29 and HEK-293 cells and have observed that there was no toxicity in HEK-293 in 40T, 60T and 80T extracts [73].

C. ferrea seed extract coated with silver nanoparticles (AgNP) presenting toxicity in L929 murine fibroblast cells exposed for 48h at the highest concentration (1,000 mg/mL), and this cytotoxicity was dose-dependent with the concentration of AgNP [74].

A cell viability test with raw extract and hydroalcoholic, aqueous and ethyl acetate fractions of the fruits of *L. ferrea* (Mart. ex Tul.) L.P. Queiroz var. *ferrea* observed that there was an increase in cell viability [29]. Prazeres et al. (2019) have performed acute toxicity test in female Wistar rats orally treated with dry extract of the fruit of *L. ferrea* at a dose of 2,000 mg/kg. Results have shown that this dose caused no death, change in behavior, change in food consumption, or weight gain during the 14 days of treatment, only an increase in water consumption [44]. In a study with zebrafish (*Danio rerio*) with hydroalcoholic extract of the aerial parts with the fruits at a dose of 2 g/kg presented toxicity in the heart of concentration-dependent embryos have been observed. In adults significant histopathological changes in the gills were caused [60].

In the preparation of the *L. ferrea* (Mart. ex Tul.) L.P. Queiroz var. *ferrea* bark and seed hydroalcoholic extract, Pickler et al. (2019) have observed that both bark and seed extract were not proven to be safe when used in the gestational phase in Wistar rats [75]. Calandrini de Azevedo et al. (2020) have observed that only with the aqueous ethanolic extract from the jucá pods in the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) test there was no change in cell survival in any concentration tested, nor did it present mutagenic or genotoxic effects [67]. Figure 3 presents the survey about the toxicity effect and 21 studies were included.




Figure 3: *In vitro* and *in vivo* toxicity observed in articles with *L. ferrea*. a. Three *in vitro* studies presented toxicity action [56, 59, 75] and six not presented toxicity action [29, 53, 59, 67, 72, 73]. b. Four *in vivo* studies presented toxicity action [32, 34, 60, 75] and six not presented toxicity action [38, 44, 46, 51, 54, 71].

In view of the above, to verify the effects and toxicological analysis, most experiments are carried out in animal models, especially in rodents.

EXPERIMENTAL TESTS

Many of the studies involving scientific research with *L. ferrea* are carried out in animal model, among which are rodents. Mice (*Mus musculus*) are easy to handle since they are small, have a rapid reproduction rate, ease of keeping them in laboratories, short life span [76, 77] and supply of many strains of blood-related [78] and, throughout history, has represented the *in vivo* model that predominates in biomedical research [76].

Studies, which have described experimental models with *L. ferrea*, identified in this research, will be described below in chronological (ascending) order.

Carvalho et al. (1996) used Swiss albino mice for analgesia testing and Wistar rats to verify anti-inflammatory action. Administration of the crude aqueous extract was oral, at concentrations of 10 and 20 mg/kg for analgesia and 100 mg/kg in the hot plate test observing analgesic property of the extract. And at a dose of 300 mg/kg of the extract administered orally inhibited paw edema [28].

Aqueous extract of the fruits of *Caesalpinia ferrea* showed a positive regulatory effect on myelopoiesis and may also act against opportunistic *Listeria monocytogenes* infection at concentrations of 500 and 1,000 mg/kg in oral BALB/c mice [79].

Female mice were used in two-stage cutaneous carcinogenesis experimentation and tested using gallic acid and methyl gallate isolated from the fruits of *Caesalpinia ferrea* Mart. demonstrating that there was a reduction in the number of papilloma [48].

Use of ellagic acid and ellagic acid 2-(2,3,6-trihydroxy-4-carboxyphenyl) (ASD) of *Caesalpinia ferrea* dried fruits *in vitro* in Diabetes-induced Wistar rats with streptozotocin (STZ) at doses of 50, 75, 100 mg/kg orally measuring sorbitol accumulation was tested by Ueda et al. (2004) suggesting that diabetic complications could be alleviated with the ingestion of fruits and vegetables containing ellagic acid [52].

There was no change in the blastocyst implementation process in three-month Wistar females in the use of 300 mg/kg via gavage of the crude aqueous extract of *C. ferrea* Mart fruits [32].

Stem bark aqueous extract used at doses of 450 mg/kg per day orally as a promising alternative in the treatment of diabetes was shown by Vasconcelos et al. (2011) in a model of streptozotocin-induced diabetes in Wistar rats [37].

Kobayashi et al. (2015) have topically tested ethanolic extract of jucá fruit and have observed that wound-healing at the dose of 12.5% in Wistar rats [54]. Topical use of hydrogel with methanol extract of fruits at a dose of 50 mg/day have demonstrated antileishmania effect in hamster (*Mesocricetus auratus*), where there was a reduction in both volume and inflamed region and presented the lower parasitic load of *Leishmania (Leishmania) amazonensis* [58].

Galactomannan of the jucá seed at dose of 10 mg/kg was used in diabetic Wistar rats induced with streptozotocin suggesting its use as a potential functional food in the treatment of type 2 diabetes [72].

Dry extract of jucá pods used at doses of 50, 100, 200 and 400 mg/kg administered orally influenced the treatment of gastric ulcer caused by indomethacin in Wistar rats. Data have also shown that the extract presents antioxidant activity, which could be used as a prevention of various diseases associated with oxidative stress [44].

In addition to studies in rats, mice, and hamsters, two studies were also found using animal models, goats, and dogs. Oliveira et al. (2010) have observed that the application of an ointment based on the stem bark of *C. ferrea* in the cutaneous treatment in experimental wounds induced in male goats aided in the wound-healing process, reducing inflammatory exudate, edema, and hyperemia [61]. Another study was that of Américo et al. (2020) in which dermal wounds were induced in dogs and a 5% ethanolic extract of *L. ferrea* fruits and flowers of the adult plant were used topically (ointment). Extract presented a potential for wound healing, suggesting its veterinary use [45].

This study has shown the following limitations: diversity of animal models, plant parts, extract types and differentiated doses found in the articles, which precludes in-depth analysis of the plant's effect on investigated disease model. Another important limitation is related to survey in databases and the method of selection of studies which may have missed some relevant publications.

CONCLUSIONS

This research aimed to bring a state-of-the-art survey of what has been published on *Libidibia ferrea* (Mart. ex Tul.) L.P. Queiroz, since jucá has been described as a medicinal plant by popular belief and by RENISUS.

Demonstration that there is low, or no toxicity related to its use, as well as the positive and/or regulatory effect in various diseases, in addition to the use in folk medicine corroborate the use of this plant in future clinical research. However, it is observed that experimental studies require a standardized design such as (Guideline Animal Research: Reporting of *in Vivo* Experiments - ARRIVE) for the quantitative confirmation of pharmacological effects.

ACKNOWLEDGEMENT

To the Foundation for “Fundação de Amparo à Pesquisa do Estado do Amazonas” (FAPEAM)/POSGRAD for the grant assistance (doctoral scholarship) to the first author of this review. Thanks to “Coordenação de Aperfeiçoamento de Pessoal de Nível Superior” (CAPES).

AUTHORS' CONTRIBUTIONS

NCOSA: Investigation; Writing – original draft; SCF: Supervision; Writing – review and editing; JFMB: Writing – review and editing.
All authors have read and approved the final manuscript.

CONSENT

Not applicable.

ETHICS APPROVAL

Not applicable.

REFERENCES

1. Ribeiro VP, Arruda C, Abd El-Salam M, Bastos JK. Brazilian medicinal plants with corroborated anti-inflammatory activities: a review. *Pharmaceutical Biology*. 2018; 56(1): 253-68. Available: <https://doi.org/10.1080/13880209.2018.1454480>.
2. ANVISA. Primeiro Suplemento - Formulário de Fitoterápicos Farmacopeia Brasileira. Agência Nacional de Vigilância Sanitária. 1ª ed. 2018:160. Portuguese.
3. Yang R, Yuan BC, Ma YS, Zhou S, Liu Y. The anti-inflammatory activity of licorice, a widely used Chinese herb. *Pharmaceutical Biology*. 2017; 55(1): 5-18. Available: <https://doi.org/10.1080/13880209.2016.1225775>.
4. Cardenas JDR. Diversidade botânica. In: dos Santos GM, França LR. GEEA: Grupo de Estudos Estratégicos Amazônicos, v. 10, (Cardeno de Debates). Editora INPA, Manaus. ISBN 978-85-211-0168-0; 2017. Portuguese.
5. Brasil. Ministério da Saúde. Política e Programa Nacional de Plantas Medicinais e Fitoterápicos. Secretaria de Ciência, Tecnologia e Insumos Estratégicos, Departamento de Assistência Farmacêutica, Brasília: Ministério da Saúde, 2016. ISBN 978-85-2399-2. Portuguese.
6. Brasil. Ministério da Saúde. Práticas integrativas e complementares: plantas medicinais e fitoterapia na Atenção Básica. Ministério da Saúde. Secretaria de Atenção à Saúde. Departamento de Atenção Básica. Série A. Normas e Manuais Técnicos. Brasília: Ministério da Saúde, 2012. ISBN 978-85-334-1912-4. Portuguese.
7. Ghasemian M, Owlia S, Owlia MB. Review of Anti-inflammatory herbal medicines. *Advances in Pharmacological Sciences*. 2016. Available: <https://doi.org/10.1155/2016/9130979>.
8. Brasil, Ministério da Saúde. A fitoterapia no SUS e o Programa de Pesquisa de Plantas Medicinais da Central de Medicamentos. Série B. T. Brasília: Ministério da Saúde, Secretaria de Ciência, Tecnologia e Insumos Estratégicos, Departamento de Assistência Farmacêutica. ISBN 85-334-1187-1; 2006. Portuguese.

9. Brasil. Ministério da Saúde. Relação Nacional de Plantas Medicinais de Interesse ao SUS - RENISUS. DAF/SCTIE/MS. Fev/2009. Portuguese.
10. Tropicos. Missouri Botanical Garden. Accessed 29 Jul 2020. Available: <http://www.tropicos.org>.
11. Di Stasi LC, Himura-Lima CA. Plantas medicinais na Amazônia e na Mata Atlântica. 2ª ed. rev. e ampli. São Paulo: Editora UNESP. ISBN 85-7139-411-3; 2002. Portuguese.
12. *Libidibia* in Lista de Espécies da Flora do Brasil. Jard Botânico Do Rio Janeiro. Accessed 01 August 2020. Available: <http://floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB109828>. Portuguese.
13. Queiroz LP. Leguminosas da Caatinga. Feira de Santana: Universidade Estadual de Feira de Santana; 2009. Portuguese.
14. Da Costa LM, Guilhon-Simplicio F, De Souza TP. *Libidibia ferrea* (Mart. Ex tul) L. P. Queiroz var. *ferrea*: Pharmacological, phytochemical and botanical aspects. International Journal of Pharmacy and Pharmaceutical Sciences. 2015; 7(4): 48-53.
15. Maia-Silva C, Silva CI da, Hrcir M, Queiroz RT de, Imperatriz-Fonseca VL. Jucazeiro. In: Guia de plantas visitadas por abelhas na caatinga, 1ª ed. Editora Fundação Brasil Cidadão, Fortaleza, CE, ISBN 978-85-98564-05-0. pp. 43; 2012. Portuguese.
16. Silva MF da, Carreira LMM, Tavares AS, Ribeiro IC, Jardim MAG, Lobo M da GA, et al. As leguminosas da Amazônia brasileira: lista prévia. Acta Botanica Brasílica. 1989; 2(1):193-237. Portuguese.
17. Stehmann JR, Faria FS, Bragioni T. 50 árvores do Museu. Belo Horizonte: Formato Editora; ISBN 978-85-62164-13-2. 2019. Portuguese.
18. Lorenzi H. Árvores Brasileiras: Manual de identificação e cultivo de plantas arbóreas nativas do Brasil. Nova Odessa, São Paulo: Plantarum; 1992. Portuguese.
19. Balbach A. Plantas que curam. 2ª ed. São Paulo: Missionária; 1972. Portuguese.
20. Galdino G, Mesquita MR, Ferraz IDK. Descrição morfológica da plântula e diásporos de *Caesalpinia ferrea* Mart – Nota científica. Revista Brasileira de Biociências, Porto Alegre. 2007; 5(2): 747-749. Portuguese.
21. Matos ACB, Ataíde G da M, Borges EE de L. Physiological, physical, and morpho-anatomical changes in *Libidibia ferrea* (Mart. ex Tul.) L.P. Queiroz) seeds after overcoming dormancy. Journal of Seed Science. 2015; 37(1): 26-32. Available: <https://doi.org/10.1590/2317-1545v37n1140433>.
22. Sousa CC, Gomes SO, Lopes ACA, Gomes RLF, Brito FB, Lima PSC, et al. Comparison of methods to isolate DNA from *Caesalpinia ferrea* - Short Communication. Genetics and Molecular Research - Online Journal. 2014; 13(2): 4486-4493. Available: <https://doi.org/10.4238/2014.June.16.7>.
23. Carvalho SMC, Torres SB, Benedito CP, Nogueira NW, Souza AAT, Souza Neta ML de. Viability of *Libidibia ferrea* (Mart. ex Tul.) L.P. Queiroz var. *ferrea* seeds by tetrazolium test. Journal of Seed Science. 2017; 39(1): 7-12. Available: <https://doi.org/10.1590/2317-1545v39n1163784>.
24. Santana JA da S, Ferreira L da S, Coelho RRP, Vieira F de A, Pacheco MV. Tecnológicas de baixo custo para superação de dormência em sementes de *Caesalpinia ferrea* var. *ferrea* Mart. ex. Tul. (pauffero). Revista Verde de Agroecologia e Desenvolvimento Sustentável. 2011; 6(1): 225-229. Portuguese.
25. Salles HO, Braga ACL, do Nascimento MT dos SC, Sousa AMP, Lima AR, Vieira L da S, et al. Lectin, hemolysin and protease inhibitors in seed fractions with ovicidal activity against *Haemonchus contortus*. Brazilian Journal of Veterinary Parasitology. 2014; 23(2): 136-143. Available: <https://doi.org/10.1590/S1984-29612014050>.
26. Barros AO, De Souza RS, Aranha ESP, Da Costa LM, De Souza TP, De Vasconcellos MC, et al. Antioxidant and hepatoprotective activities of *Libidibia ferrea* bark and fruit extracts. International Journal of Pharmacy and Pharmaceutical Sciences. 2014; 6(11): 71-76.

27. Santos, SF dos, Santos, AS, Corpes, RS, Leão, NV. Aspectos dos cultivo *in vitro* de *Libidibia ferrea* (Mart. ex Tul.) L.P. Queiroz (Leguminosae-Caesalpinioideae) como fonte alternativa para produção de metabólitos secundários. *Revista Espacios*. 2018, 39(37): 17-24. Portuguese.
28. Carvalho JCT, Teixeira JRM, Souza PJC, Bastos JK, Dos Santos Filho D, Sarti SJ. Preliminary studies of analgesic and anti-inflammatory properties of *Caesalpinia ferrea* crude extract. *Journal of Ethnopharmacology*. 1996; 53: 175-178. Available: [https://doi.org/10.1016/0378-8741\(96\)01441-9](https://doi.org/10.1016/0378-8741(96)01441-9).
29. Falcão TR, Araújo AA De, Soares LAL, Farias IB De, Silva WAV Da, Ferreira MRA, et al. *Libidibia ferrea* fruit crude extract and fractions show anti-inflammatory, antioxidant, and antinociceptive effect *in vivo* and increase cell viability *in vitro*. *Evidence-Based Complementary Alternative Medicine*. 2019; 2019, ID 6064805, 14 pages. Available: <https://doi.org/10.1155/2019/6064805>.
30. Freitas ACC, Ximenes NCA, Aguiar JS, Nascimento SC, Lins TUL, Magalhães LR, et al. Biological activities of *Libidibia (Caesalpinia) ferrea* var. *parvifolia* (Mart. ex Tul.) L. P. Queiroz pod preparations. *Evidence-Based Complementary Alternative Medicine*. 2012; 2012: ID 514137, 7 pages. Available: <https://doi.org/10.1155/2012/514134>.
31. Lucinda LMF, Rocha CB, Reboredo MM, Faria VC, Sá RCS. Assessment of sperm production and reproductive organs of Wistar rats to long-term exposure of *Caesalpinia ferrea*. *Anais da Academia Brasileira de Ciências*. 2010; 82(4) :907-914. Available: <https://doi.org/10.1590/s0001-37652010000400013>.
32. Peters VM, Souza SO, Carvalho JCT, Borges L V., Guerra MO. Evaluation of reproductive toxicity of aqueous extract of the fruits from *Caesalpinia ferrea* Mart. in rats. *Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas*. 2008; 7(5): 268-272.
33. Sampaio FC, Pereira M do S V., Dias CS, Costa VCO, Conde NCO, Buzalaf MAR. *In vitro* antimicrobial activity of *Caesalpinia ferrea* Martius fruits against oral pathogens. *Journal of Ethnopharmacology*. 2009; 124(2019): 289-94. Available: <https://doi.org/10.1016/j.jep.2009.04.034>.
34. Reboredo M de M, Lucinda LMF, Rocha CB, Queiroz GT de, Faria VC de, Vieira V de A, et al. Avaliação da toxicidade do extrato aquoso de *Caesalpinia ferrea* em órgãos vitais, no sistema reprodutor e na produção de espermatozoides de ratos Wistar submetidos a tratamento subagudo. *Boletim do Centro de Biologia da Reprodução*. 2006; 25: 17-29. Portuguese.
35. Falcão TR, Rodrigues CAO, De Araújo AA, De Medeiros CACX, Soares LAL, Ferreira MRA, et al. Crude extract from *Libidibia ferrea* (Mart. ex. Tul.) L.P. Queiroz leaves decreased intra articular inflammation induced by zymosan in rats. *BMC Complementary and Alternative Medicine*. 2019; 19(47): 1-10. Available: <https://doi.org/10.1186/s12906-019-2454-3>.
36. De Araújo AA, Soares LAL, Assunção Ferreira MR, De Souza Neto MA, Da Silva GR, De Araújo RF, et al. Quantification of polyphenols and evaluation of antimicrobial, analgesic and anti-inflammatory activities of aqueous and acetone-water extracts of *Libidibia ferrea*, *Parapiptadenia rigida* and *Psidium guajava*. *Journal of Ethnopharmacology*. 2014; 156(2014): 88-96. Available: <https://doi.org/10.1016/j.jep.2014.07.031>.
37. Vasconcelos CFB, Maranhão HML, Batista TM, Carneiro EM, Ferreira F, Costa J, et al. Hypoglycaemic activity and molecular mechanisms of *Caesalpinia ferrea* Martius bark extract on streptozotocin-induced diabetes in Wistar rats. *Journal of Ethnopharmacology*. 2011; 137(2011): 1533-1541. Available: <https://doi.org/10.1016/j.jep.2011.08.059>.
38. Cavalheiro MG, Farias DF, Fernandes GS, Nunes EP, Cavalcanti FS, Vasconcelos IM, et al. Atividades biológicas e enzimáticas do extrato aquoso de sementes de

- Caesalpinia ferrea* Mart., Leguminosae. *Revista Brasileira de Farmacognosia*. 2009; 19(2B): 586-591. Available: <https://doi.org/10.1590/S0102-695X2009000400014>. Portuguese.
39. Sawada LA, Monteiro VSDC, Rabelo GR, Dias GB, Da Cunha M, Do Nascimento JLM, et al. *Libidibia ferrea* mature seeds promote antinociceptive effect by peripheral and central pathway: Possible involvement of opioid and cholinergic receptors. *BioMed Research International*. 2014; 2014, ID 508725, 10 pages. Available: <https://doi.org/10.1155/2014/508725>.
 40. Agra M de F, Freitas PF de, Barbosa-Filho JM. Synopsis of the plants known as medicinal and poisonous in Northeast of Brazil. *Revista Brasileira de Farmacognosia*. 2007; 17(1): 114-140. Available: <https://doi.org/10.1590/S0102-695X2007000100021>.
 41. Santos KA dos, Vilanova CM. Estudo etnobotânico de plantas medicinais utilizadas como hipoglicemiantes por usuários do Programa de Fitoterapia da Universidade Federal do Maranhão, Brasil. *Scientia Plena*. 2017; 13(03): 1-12. Available: <https://doi.org/10.14808/sci.plena.2017.034501>. Portuguese.
 42. Gomes TMF, Lopes JB, Barros RFM de, Alencar NL. Plantas de uso terapêutico na comunidade rural Bezerro Morto, São João da Canabrava, Piauí, Brasil. *Gaia Scientia*. 2017; 11(1): 253-268. Available: <https://doi.org/10.22478/ufpb.1981-1268.2017v11n1.33683>. Portuguese.
 43. Reis CRM, Prereira AF de N, Cansação IF. Levantamento etnobotânico de plantas medicinais utilizadas por moradores do entorno do Parque Nacional Serra da Capivara - PI. *Biofarm*. 2017; 13(04): 7-21. Portuguese.
 44. Prazeres LDKT, Aragão TP, Brito SA, Almeida CLF, Silva AD, De Paula MMF, et al. Antioxidant and antiulcerogenic activity of the dry extract of pods of *Libidibia ferrea* Mart. ex Tul. (Fabaceae). *Oxidative Medicine and Cellular Longevity*. 2019; 2019. Available: <https://doi.org/10.1155/2019/1983137>.
 45. Américo AVL dos S, Nunes KM, Assis FFV de, Dias SR, Passos CTS, Morini AC, et al. Efficacy of phytopharmaceuticals from the Amazonian plant *Libidibia ferrea* for wound healing in dogs. *Frontiers in Veterinary Science*. 2020; 7: 1-11. Available: <https://doi.org/10.3389/fvets.2020.00244>.
 46. Lima SMA, Araújo LCC, Sitônio MM, Freitas ACC, Moura SL, Correia MTS, et al. Anti-inflammatory and analgesic potential of *Caesalpinia ferrea*. *Revista Brasileira de Farmacognosia*. 2012; 22(1): 169-175. Available: <https://doi.org/10.1590/S0102-695X2011005000197>.
 47. Nakamura ES, Kurosaki F, Arisawa M, Mukainaka T, Okuda M, Tokuda H, et al. Cancer chemopreventive effects of constituents of *Caesalpinia ferrea* and related compounds. *Cancer Letters*. 2002; 177(2002): 119-124. Available: [https://doi.org/10.1016/S0304-3835\(01\)00708-X](https://doi.org/10.1016/S0304-3835(01)00708-X).
 48. Nakamura ES, Kurosaki F, Arisawa M, Mukainaka T, Takayasu J, Okuda M, et al. Cancer chemopreventive effects of a Brazilian folk medicine, Juca, on *in vivo* two-stage skin carcinogenesis. *Journal of Ethnopharmacology*. 2002; 81(2002): 135-137. Available: [https://doi.org/10.1016/S0378-8741\(02\)00047-8](https://doi.org/10.1016/S0378-8741(02)00047-8).
 49. Oliveira GP, Souza TP, Caetano SK, Farias KS, Venancio GN, Bandeira MFCL, et al. Atividade antimicrobiana *in vitro* de extratos da casca do caule e da vagem de *Libidibia ferrea* L. frente a microrganismos da cavidade bucal. *Revista Fitos*. 2013; 8(2): 95-102. Available: <https://doi.org/10.5935/1808-9569.20130004>. Portuguese.
 50. Silva LCN da, Sandes JM, Paiva MM de, Araújo JM de, Figueiredo RCBQ de, Silva MV da, et al. Anti-*Staphylococcus aureus* action of three Caatinga fruits evaluated by electron microscopy. *Natural Product Research: Formerly Natural Products Letters*. 2012; 1-5. Available: <https://doi.org/10.1080/14786419.2012.722090>.
 51. Hassan SK, El-Sammad NM, Mousa AM, Mohammed MH, Farrag A el RH, Hashim ANE, et al. Hypoglycemic and antioxidant activities of *Caesalpinia ferrea* Martius leaf

- extract in streptozotocin-induced diabetic rats. *Asian Pacific Journal of Tropical Biomedicine*. 2015; 5(6): 462-471. Available: <https://doi.org/10.1016/j.apjtb.2015.03.004>.
52. Ueda HU, Awanishi KK, Oriyasu MM. Effects of Ellagic Acid and 2-(2, 3, 6-trihydroxy-4-carboxyphenyl) ellagic Acid on Sorbitol Accumulation *in vitro* and *in vivo*. *Biological and Pharmaceutical Bulletin*. 2004; 27(10): 1584-1587.
53. Dias AMA, Rey-Rico A, Oliveira RA, Marceneiro S, Alvarez-Lorenzo C, Concheiro A, et al. Wound dressings loaded with an anti-inflammatory jucá (*Libidibia ferrea*) extract using supercritical carbon dioxide technology. *The Journal of Supercritical Fluids*. 2013; 74(2014): 34-45. Available: <https://doi.org/10.1016/j.supflu.2012.12.007>.
54. Kobayashi YT da S, Almeida VT de, Bandeira T, Alcântara BN de, Silva ASB da, Barbosa WLR, et al. Avaliação fitoquímica e potencial cicatrizante do extrato etanólico dos frutos de Jucá (*Libidibia ferrea*) em ratos Wistar. *Brazilian Journal of Veterinary Research and Animal Science*. 2015; 52(1): 34-40. Available: <https://doi.org/10.11606/issn.1678-4456.v52i1p34-40>. Portuguese.
55. Lopes N, Faccin-Galhardi LC, Espada SF, Pacheco AC, Ricardo NMPS, Linhares REC, et al. Sulfated polysaccharide of *Caesalpinia ferrea* inhibits herpes simplex virus and poliovirus. *International Journal of Biological Macromolecules*. 2013; 60(2013): 93-99. Available: <https://doi.org/10.1016/j.ijbiomac.2013.05.015>.
56. Marques MMM, Morais SM De, Silva ARA Da, Barroso ND, Pontes Filho TR, Araujo FMDC, et al. Antiviral and Antioxidant activities of sulfated Galactomannans from plants of Caatinga Biome. *Evidence-Based Complementary Alternative and Medicine*. 2015; 2015. Available: <https://doi.org/10.1155/2015/591214>.
57. Fernandes CPM, Machado C, Lopes TV, Cunha Filho N, Bretanha PR, Schons S, et al. Repellent Action of *Carapa guianensis* and *Caesalpinia ferrea* for flies species of Calliphoridae family. *Ciência Rural*. 2016; 46(5): 867-870. Available: <https://doi.org/10.1590/0103-8478cr20150727>.
58. Comandoli-Wyrepkowskil CD, Jensen BB, Grafova I, Santos PA dos, Barros AMC, Soares FV, et al. Antileishmanial activity of extracts from *Libidibia ferrea*: development of *in vitro* and *in vivo* tests. *Acta Amazonica*. 2017; 47(4): 331-340. Available: <https://doi.org/10.1590/1809-4392201700871>.
59. Pedrosa T do N, Barros AO, Nogueira JR, Fruet AC, Rodrigues IC, Calcagno DQ, et al. Anti-wrinkle and anti-whitening effects of jucá (*Libidibia ferrea* Mart.) extracts. *Archives of Dermatological Research*. 2016. Available: <https://doi.org/10.1007/s00403-016-1685-0>.
60. Ferreira DQ, Ferraz TO, Araújo RS, Cruz RAS, Fernandes CP, Souza GC, et al. *Libidibia ferrea* (jucá), a traditional anti-inflammatory: A study of acute toxicity in adult and embryos zebrafish (*Danio rerio*). *Pharmaceuticals*. 2019; 12(175): 1-15. Available: <https://doi.org/10.3390/ph12040175>.
61. Oliveira AF, Batista JS, Paiva ES, Silva AE, Farias YJMD, Damasceno CAR, et al. Avaliação da atividade cicatrizante do jucá (*Caesalpinia ferrea* Mart. ex Tul. var. *ferrea*) em lesões cutâneas de caprinos. *Revista Brasileira de Plantas Mediciniais*. 2010; 12(3): 302-310. Available: <https://doi.org/10.1590/s1516-05722010000300007>. Portuguese.
62. Ueda H, Tachibana Y, Moriyasu M, Kawanishi K, Alves SM. Aldose reductase inhibitors from the fruits of *Caesalpinia ferrea* Mart. *Phytomedicine*. 2001; 8(5): 377-381. Available: <https://doi.org/10.1078/0944-7113-00043>.
63. Ferreira MA, Fernandes MM, da Silva W V, Bezerra IF, de Souza T, Pimentel M, et al. Chromatographic and spectrophotometric analysis of phenolic compounds from fruits of *Libidibia ferrea* Martius. *Pharmacognosy Magazine*. 2016; 0:0. Available: <https://doi.org/10.4103/0973-1296.179665>.
64. Gonzalez FG. Estudo farmacognóstico e farmacológico de *Caesalpinia ferrea* Martius. Universidade de São Paulo, Doctoral's Thesis. 2005. Portuguese.

65. Silva LCN, Miranda R de CM de, Gomes E de B, Macedo AJ, Araújo JM de, Figueiredo RCBQ de, et al. Evaluation of combinatory effects of *Anadenanthera colubrina*, *Libidibia ferrea* and *Pityrocarpa moniliformis* fruits extracts and erythromycin against *Staphylococcus aureus*. *Journal of Medicinal Plants Research*. 2013; 7(32): 2358-2364. Available: <https://doi.org/10.5897/JMPR2013.2597>.
66. Frasson APZ, Bittencourt CF, Heinzmann BM. Caracterização físico-química e biológica do caule de *Caesalpinia ferrea* Mart. *Revista Brasileira de Farmacognosia*. 2003; 13(1): 35-39. Available: <https://doi.org/10.1590/s0102-695x2003000100004>. Portuguese.
67. Calandrini de Azevedo LF, Alves Ferreira TA, Melo KM, Porfírio Dias CL, Bastos CEMC, Santos SF, et al. Aqueous ethanol extract of *Libidibia ferrea* (Mart. ex Tul) L.P. Queiroz (juca) exhibits antioxidant and migration-inhibiting activity in human gastric adenocarcinoma (ACP02) cells. *PLoS One*. 2020; 15(1):e0226979. Available: <https://doi.org/https://doi.org/10.1371/journal.pone.0226979>.
68. Pereira VS, Saraiva CRN, Rocha JE, Lopes J da C, Silva MK do N, Bandeira SMF, et al. Estudo químico, toxicidade e atividade antimicrobiana do óleo essencial de *Ocimum gratissimum*. *Revista Interfaces: Saúde, Humanas e Tecnologia*. 2014; 2(Número especial): 2014. Portuguese.
69. Negri G. Diabetes melito: Plantas e princípios ativos naturais hipoglicemiantes. *Revista Brasileira de Ciências Farmacêuticas*. 2005; 41(2): 121-42. Available: <https://doi.org/10.1590/S1516-93322005000200002>. Portuguese.
70. Ferreira MRA, Soares LAL. *Libidibia ferrea* (Mart. ex Tul.) L. P. Queiroz: A review of the biological activities and phytochemical composition. *Journal of Medicinal Plants Research*. 2015; 9(2): 140-150. Available: <https://doi.org/10.5897/jmpr2014.5706>.
71. Souza AB de, Mara L, Souza S, Carlos J, Carvalho T, Maistro EL. No clastogenic activity of *Caesalpinia ferrea* Mart. (Leguminosae) extract on bone marrow cells of Wistar rats. *Genetics and Molecular Biology*. 2006; 29(2): 380-383.
72. Cunha AP, Ribeiro ACB, Ricardo NMPS, Oliveira AC, Dávila LSP, Cardoso JHL, et al. Polysaccharides from *Caesalpinia ferrea* seeds - Chemical characterization and anti-diabetic effects in Wistar rats. *Food Hydrocolloids*. 2017; 65: 68-76. Available: <https://doi.org/10.1016/j.foodhyd.2016.10.039>.
73. Guerra ACV de A, Soares LAL, Ferreira MRA, Araújo AA de, Rocha HA de O, Medeiros JS de, et al. *Libidibia ferrea* presents antiproliferative, apoptotic and antioxidant effects in a colorectal cancer cell line. *Biomedicine and Pharmacotherapy*. 2017; 92: 696-706. Available: <https://doi.org/10.1016/j.biopha.2017.05.123>.
74. Soares MRPS, Corrêa RO, Stroppa PHF, Marques FC, Andrade GFS, Corrêa CC, et al. Biosynthesis of silver nanoparticles using *Caesalpinia ferrea* (Tul.) Martius extract: Physicochemical characterization, antifungal activity and cytotoxicity. *PeerJ*. 2018; 2018: 1-16. Available: <https://doi.org/10.7717/peerj.4361>.
75. Pickler TB, Lopes KP, Magalhães SA, Krueger CMA, Martins MM, Filho VC, et al. Effect of *Libidibia ferrea* bark and seed in maternal reproductive and biochemical outcomes and fetal anomaly in rats. *Birth Defects Research*. 2019; 111: 863-871. Available: <https://doi.org/10.1002/bdr2.1520>.
76. Franco NH. Animal experiments in biomedical research: A historical perspective. *Animals*. 2013; 3: 238-273. Available: <https://doi.org/10.3390/ani3010238>.
77. Baumans V. The Welfare of Laboratory Rats: In: *The Welfare of Laboratory Animals*. vol. 2. Springer. 2007. Available: <https://doi.org/10.1007/978-1-4020-2271-5>.
78. Perlman RL. Mouse Models of Human Disease: An Evolutionary Perspective. *Evolution, Medicine & Public Health*. 2016: eow014, 170-176. Available: <https://doi.org/10.1093/emph/eow014>.
79. Queiroz MLS, Justo GZ, Valadares MC, Pereira-da-Silva FRR. Evaluation of *Caesalpinia ferrea* extract on bone marrow hematopoiesis in the murine models of listeriosis and

Ehrlich ascites tumor. *Immunopharmacology and Immunotoxicology*. 2001; 23(3): 367-382. Available: <https://doi.org/10.1081/IPH-100107337>.

UNDR PEER REVIEW