

## Original Research Article

# Impact of Acute Kidney Injury pRIFLE Criteria Implementation in Pediatric Critical care

### ABSTRACT

**Aims:** The study aims to assess the impact of using pRIFLE criteria to define the incidence, the main causes, outcome, and associated morbidity of AKI at SQUH pediatric ICU. Also, we aim to implement a guideline for the early detection and management of AKI in critically sick Omani children.

**Study design:** A retrospective study included Omani children, less than 14 years, admitted to Pediatric intensive care, with Acute kidney injury.

**Place and Duration of Study:** The study was conducted between 2016-2017 in Sultan Qaboos University Hospital, PICU.

**Methodology:** AKI diagnosis and staging were based on pRIFLE criteria. Serum creatinine level and Urine output (UOP) were enrolled for all patients. Estimated Creatinine Clearance (eCrCL) was calculated using the Schwartz formula. Data was compiled using SPSS software using the chi-square test and fisher's exact test.

**Results:** Out of 499 cases, 469 cases were included while 30 cases were excluded. The incidence of AKI was 9.17% (43). Stage 'Risk (R)', 'Injury (I)' and 'Failure (F)' constituted 32.6% (14), 27.9% (12) and 39.5% (17) respectively. Children below five years were more developing AKI with a frequency of 79.07% ( $p= 0.80$ ). UOP and serum creatinine classification were matching in 42.11%. In regards the primary cause of AKI, Ventilators had the highest frequency (88.4%), followed by Hypotension (65.1%), Sepsis (58.1%), and nephrotoxic drug (14.50%). Prerenal causes constituted 88.47% and renal 11.63%.

**Conclusion:** pRIFLE criteria are an easy tool for early identification and stratification of AKI. A higher incidence of AKI patients was in the failure stage on admission with preventable and treatable Pre-renal causes. Ventilation is still the main concern of AKI in critical children. Nevertheless, Hypotension, sepsis, and drugs are the main risk factor contributing to AKI.

*Keywords: Acute Kidney Injury; PICU; SQUH; pRIFLE; Oman*

### 1. INTRODUCTION

Acute Kidney Injury (AKI) is a frequent and sometimes devastating syndrome, with high costs to patients and healthcare systems.<sup>1</sup> AKI is a sudden

reduction of a kidney function in which there is a significant decrease in glomerular filtration rate (GFR).<sup>2</sup> The most consequence of AKI are increasing serum creatinine (or cystatin c), retention of nitrogenous waste, and inability to regulate electrolyte and fluid hemostasis.<sup>2,3</sup> Serum creatinine elevation is the most characteristic marker that is used for the diagnosis of AKI. However, Serum creatinine level does not reflect the kidney function accurately because of the delay in creatinine rise in those individuals with volatile kidney function. Moreover, it is influenced by many non-renal factors, such as the nutritional and hydration status of the child.<sup>2,4</sup> So, serum creatinine cannot be used alone to describe an abrupt decline of renal function. Ideal AKI staging criteria should be simple, sensitive, specific, low cost, and good correlation between stage and outcome.<sup>1</sup> Therefore, standardized definitions for pediatric AKI appear to be multidimensional, which depends on staging and accurate assessment using pRIFLE criteria "Pediatric Risk, Injury, Failure, Loss, End-Stage Renal Disease", which depends on serum creatinine and urine output.<sup>1,5</sup>

Unfortunately, the incidence of AKI increases globally with a simultaneous increase in mortality and morbidity. A world meta-analysis showed that 13.3 million people develop AKI, out of which 1.7 million die every year.<sup>6</sup> In addition, the incidence of AKI is 33.7% in children, when compared to 21% in adults.<sup>6</sup> AKI is considered a major cause that increases mortality in pediatric patients.<sup>7</sup> The incidence of AKI is varying from one region to another at the same time. Many studies approve that AKI has various burdens between the developing countries and developed countries. Moreover, the younger age patients in developing countries are more susceptible to developing AKI, and infection is considered the main cause.<sup>8</sup> Among the infection causes, infection was found the most prevalent cause of AKI followed by acute gastroenteritis.<sup>9</sup>

A careful history, physical examination, and investigations are essential to determine the cause of AKI.<sup>9,10</sup> The etiological classification of AKI is divided into three main categories; Pre-renal injury, Intrinsic renal disease, and Obstructive uropathy. The main causes of Pre-renal injury are reducing intravascular volume (true or effective intravascular volume). While there are several etiologies for intrinsic renal injuries such as acute tubular necrosis, Uric acid nephropathy, Toxin mediated, Drug-induced, and vascular glomerulonephritis. However, it can be Idiopathic in many cases. Whereas, bilateral ureteral obstruction, ureteral Obstruction in a solitary kidney, and Urethral obstruction are the main causes of Obstructive uropathy.<sup>10</sup>

AKI is usually treatable and preventable with few long-term health outcomes. A high incidence of AKI in some countries means a lack of early fundamental care. So, we need to raise awareness about AKI which includes early detection and appropriate management to prevent any complications. In addition, no studies regrettably were conducted in Oman related to AKI in pediatric patients. So, the observed data are an important base for further studies of AKI in Oman. Our study aims to assess the impact of using pRIFLE criteria to define the incidence, the main causes, outcome, and associated morbidity of AKI in critically ill Omani children.

## 2. METHODOLOGY

### 2.1 STUDY DESIGN AND SUBJECTS

This retrospective observational study was conducted in the Department of Child Health, Sultan Qaboos University Hospital (SQUH), in Muscat, Oman. Medical records of children admitted to Pediatric Intensive care Unit (PICU), in the period between January 2016 and December 2017, were obtained from SQUH electronic database. All children aged 1 month to 14 years old were enrolled for the study. Children with chronic kidney diseases were excluded.

### 2.2 DATA COLLECTION

All demographic characteristics which include age, gender, weight, height, geographical origin, and disease duration were obtained. In addition, the cause of admission and clinical and laboratory investigation were recorded. However, AKI diagnosis and staging were based on pRIFLE (pediatric RIFLE) criteria (table1).<sup>8</sup> Therefore, Urine output (ml/kg/hour) and Serum creatinine level also were recorded for each child diagnosed with AKI. Recording of serum creatinine was at diagnosis then after 6 hours, 12 hours, and 24 hours of (6h,12h,24h). Baseline creatinine was determined by taking into consideration previous baseline readings, age-related creatinine range, and lowest creatinine measurement during hospitalization.<sup>1</sup> In addition, the estimated glomerular filtration rate (eGFR) was calculated by using the Schwartz formula. Moreover, detailed clinical history was used to determine the risk factors, the main cause, and outcome of AKI in diagnosed patients.

**Table 1: pRIFLE criteria for classification of Acute Kidney Injury in children**

Class	Serum creatinine/ GFR criteria	Urine output criteria
Risk	Serum creatinine *1.5 or GFR loss > 25%	0.5 < ml/kg/hour * > 6 hrs
Injury	Serum creatinine *2 or GFR loss > 50%	0.5 < ml/kg/hour * > 12 hrs
Failure	Serum creatinine *3 or GFR loss > 75%	0.3 < ml/kg/hour * > 24 hrs <b>Or</b> anuria > 12 hrs
Loss	Persistent failure of kidney function > 4 weeks	
End stage	Persistent failure of kidney function > 3 months	

Abbreviations: GFR = Glomerular Filtration Rate, hrs = hours.

### 2.3 STATISTICAL ANALYSIS


Data was analyzed utilizing SPSS (Statistical Package for Social Sciences) program 23, IBM, Chicago, Illinois, USA. Analysis included demographic features of the examined patients, Descriptive statistics were utilized including frequencies and percentages for categorical variables. In addition to mean ( $\pm$  standard deviation) for numerical variables. A Comparison was made between patients using Chi-squared

( $\chi^2$ ) test or fisher's exact test as appropriate. An a priori two-tailed level of significance was set at 0.05.

### 3. RESULTS

During the study period, a total of 499 cases were admitted to PICU, with 30 cases meeting the exclusion criteria. 43 out of 469 patients developed AKI, with an incidence of 9.17% in the age range 0.1-11.79 years, mean 3.03 years old, with male predominance. According to PRIFLE criteria, stage 'Risk' comprised maximum cases with 32.6% (14 cases), followed by 'Injury' comprised 27.9% (12 cases) and 'Failure' comprised 39.5% (17 cases). Out of 43 AKI patients, 30 (69.77%) patients had AKI at admission while 25.58% developed within 72 h.

**FIGURE 1:** Flowchart and demographics of children with Acute Kidney Injury (AKI)



Abbreviations: AKI = Acute Kidney Injury, PICU- Pediatric Intensive care unit, SQUH = Sultan Qaboos University Hospital, n = number, IQR= inter-quartile ratio

Serum creatinine criteria were screened amongst all cases to classify AKI patients according to pRIFLE criteria to risk, injury, and failure stages. For serum creatinine criteria, 43 cases had AKI. Incidence of Risk stage (R) was 14 (32.56%), Injury stage (I) was 12 (27.9%) and Failure stage (F) was 17 (39.53%). Urine output criteria were only applied to 19 cases out of 43 cases of AKI. Among these 19 cases, Stage 'R', stage "I" and stage "F" cases were 6 (31.58%), 5(26.32%), and 8(42.12%) (Table2). There was matching between serum creatinine criteria and urine output criteria in 8 cases (42.11%). The matching rate in stage "R" was 16.67%, while there was 40.00 % matching in stage "I" and 62.50% in stage "F" (Table 3).

**TABLE 1:** Relation of creatinine, urine output criteria and incidence of Acute Kidney Injury

Stage	Serum Creatinine Criteria			Urine Output Criteria		
	No AKI	AKI	Total	No AKI	AKI	Total
No AKI	426(100%)	0(0%)	<b>426(90.83%)</b>	442(100%)	0(0%)	<b>450(95.95%)</b>
Risk	0(0%)	14(32.56%)	<b>14(2.99%)</b>	0(0%)	6(31.58%)	<b>6(1.28%)</b>
Injury	0(0%)	12(27.90%)	<b>12(2.56%)</b>	0(0%)	5(26.32%)	<b>5(1.07%)</b>
Failure	0(0%)	17(39.53%)	<b>17(3.62%)</b>	0(0%)	8(42.12%)	<b>8(1.70%)</b>
<b>Total</b>	<b>426(100%)</b>	<b>43(100%)</b>	<b>469(100%)</b>	<b>442(100%)</b>	<b>19(100%)</b>	<b>469(100%)</b>

*P= 0.30 (kappa test)*

**TABLE 3:** Pairing of creatinine criteria staging and urine output staging

Final RIFLE Stage		Urine Output Stages (U)			Total
		R(U)	I(U)	F(U)	
Serum Creatinine Stages	R	<b>1(16.67%)</b>	2(40.00%)	1(12.50%)	<b>4(21.05%)</b>
	I	3(50.00%)	<b>2(40.00%)</b>	2(25.00%)	<b>7(36.84%)</b>
	F	<b>2(33.33%)</b>	1(20.00%)	<b>5(62.50%)</b>	<b>8(42.11%)</b>
<b>Total</b>		<b>6(100%)</b>	<b>5(100%)</b>	<b>8(100%)</b>	<b>19(100%)</b>

Abbreviations: R = Risk, I = Injury, F = Failure, U = Urine.

In our study, the frequency of children developing AKI below five years was abundant 79.07% (34 cases) but there was no association ( $p= 0.8$ ). However, the percentage of patients of age (5-9 years) and (9-14 years) are 11.63% (5 cases), 9.3% (4 cases) respectively. The predominant causes of AKI in our studied group were the pre-renal causes which represented 38 cases and a percentage of 88.47% of AKI cases, while renal causes appeared in 11.63% only and no post-renal causes. Sepsis had the highest frequency as a primary cause in the studied group and it was present in 44.19% of AKI cases. Cardiopulmonary arrest and dehydration were the second most frequent cause of AKI and they were present in 18.6% of AKI cases (Table 4). The frequency of risk factors among AKI patients varied. Patients with ventilators had the highest frequency which was 88.37%, then followed by hypotension then drug mediated AKI (Table 5). Sepsis, pneumonia, drug-mediated, and ventilator were the risk factors that get a significant association with developing AKI ( $p<0.05$ ). However, the nephrotoxic drug did not have a reliable association with AKI ( $p=0.57$ ) in our group study. (Table 6). Among AKI patients, full recovery is the common outcome which occupied 55.81%, in contrast, the rate of mortality was 44.19%.

**Table 4:** Primary causes among AKI patients

	AKI cases	
	Frequency	Percentage
Dehydration	3	18.60%
Hypotension	1	4.65%
Sepsis	22	44.19%
Hemorrhage	1	2.33%
Metabolic Disease	3	2.32%
Hematologic Disease	1	2.33%
Respiratory Disease	2	2.32%
Drug mediated	1	2.33%
Epilepticus status	1	2.33%
Cardiopulmonary arrest	8	18.60%
<i>Total</i>	<b>43</b>	<b>100%</b>

**Table 5:** Correlation of clinical variable in relation to incidence of AKI

Variables		Final Rifle Stage			
		AKI (43 Cases)	NON-AKI (426 cases)	Total (469 cases)	P-Value
<b>Gender</b>					
	<i>Male</i>	19 (44.2%)	188(44.1%)	262(55.9%)	<i>P=.</i> 559
	<i>Female</i>	24 (55.8%)	238(55.9%)	207(44.1)	
<b>Hypotension</b>					
	<i>Yes</i>	28 (65.1%)	25 (5.9%)	53(11.3%)	<i>P&lt;.0001</i>
	<i>No</i>	15(34.9%)	399(94.1%)	414(88.7%)	
<b>Sepsis</b>					
	<i>Yes</i>	25(58.1%)	52(12.2%)	77(16.5%)	<i>P&lt;.0001</i>
	<i>No</i>	18(41.9%)	373(87.8%)	391(83.5%)	
<b>Nephrotoxic Drug</b>					
	<i>Yes</i>	20 (46.5%)	88 (20.7%)	108 (23.1%)	<i>P&lt;.0001</i>
	<i>No</i>	23 (53.5%)	338 (79.34%)	360 (76.9%)	
<b>Ventilator</b>					
	<i>Yes</i>	38 (88.4%)	82 (19.2%)	120 (25.6%)	<i>P&lt;.0001</i>
	<i>No</i>	5 (11.6%)	344 (80.7%)	349 (74.4%)	

**Table 6:** Assessment of the risk factors for development of AKI

Variable	Logit coefficient	S.E	Wald	P Value	Adj. OR	95% C.I	
						Lower	Upper
Hypotension	3.335	0.555	36.115	<0.0001	28.085	9.464	83.345
Sepsis	1.280	0.498	6.604	<0.01	3.597	1.355	9.550
Ventilation	3.520	0.606	33.685	<0.0001	33.783	10.291	110.902
Nephrotoxic Drug	0.283	0.503	0.317	0.573	1.328	0.495	3.558

#### 4. DISCUSSION

AKI was defined in our study by using pRIFLE criteria. This was comparable to lots of studies that were done in this field. The cause of widely used pRIFLE criteria is simplicity and high sensitivity in the pediatric age group.<sup>11</sup> The Rustagi et al study showed a higher incidence of AKI (14%) which may reflect the inclusion of patients up to 18 years old.<sup>12</sup> On the other hand, the incidence in our study was lower than in other studies that were conducted in developing countries as the incidence of AKI is 26.1% in Bangalore, India where Dengue spread.<sup>8</sup> Meanwhile, Gupta et al reported an incidence of 42.9% in Bikaner, India.<sup>10</sup> We must put into consideration that these two last studies are prospective studies, which can avoid the limitations of the retrospective studies. In addition, this variation can be explained by regional differences, heterogeneity of patients, and sample size. Our incidence can markedly increase if we added the missed UOP criteria. Koeze et al concluded urine output criteria detect AKI earlier and double incidences in critically ill patients.<sup>13</sup>

In our study, 69.77% of patients developed AKI from admission while 25.58% developed AKI early in the course of PICU, in comparison Gupta et al identify that 50% had AKI on admission and 40% develops AKI within 72h of PICU stay, emphasizing the importance of early diagnosis of AKI.<sup>10</sup> Also, grading of AKI according to pRIFLE criteria displayed that 32.6%, 27.9%, and 39.5% that patients were in the "R" stage, "I" stage, and "F" stage, respectively. This pattern of distribution is different from other studies for example, in Gupta et al AKI stratum R, I, and F in diagnosis is 49.1%, 29.5%, 21.3% respectively<sup>9</sup> and these results are comparable to Srinivasa et al.<sup>6</sup> Regarding "F" stage in our study, 16 patients out of 17 having AKI in failure stage, had AKI from admission. This was an indication of the importance of raising doctors' awareness about AKI Characteristics at primary and secondary health facilities before reaching our tertiary care facilities.

Serum creatinine and urine output are used in pediatric RIFLE criteria for AKI severity staging. In our study, the matching between urine output classification and serum creatinine classification was 42.11%. In comparison with Srinivasa et al study where the matching between the two criteria is 77%.<sup>7</sup> They applied urine output only to those patients who were catheterized to increase the accuracy of urine output classification and the matching will be more accurate.<sup>7</sup> In our study, it was observed the matching between urine output and serum creatinine increased with AKI progression which was explained by the insertion of a urinary catheter with progressive stages of AKI. Also, missing documentation of accurate urine output in the early stage of AKI.

The study showed that patients below five years occupied 77.83% of patients who developed AKI. However, there was no significant association between age and developing AKI. However, this association is approved in Metha et al study which shows that younger children develop AKI more than others.<sup>6</sup> In addition, Srinivasa et al also show patients who are below one year more likely to develop AKI.<sup>7</sup>

The etiological spectrum of AKI was different in developing countries and developed countries.<sup>6</sup> Sepsis and glomerulonephritis are the common causes of AKI in developing countries. In contrast, nephrotoxic drugs, major surgery, and pulmonary failure are the predominant causes of AKI in developed countries.<sup>6</sup> In our study, the most common primary cause was sepsis (44.19%) followed by dehydration and cardiopulmonary arrest (18.60%). Sepsis had a significant association with developing AKI and that was comparable to Sinha et al and other studies, where they also find that sepsis is the most common etiology associated with AKI.<sup>14,15</sup> Furthermore, hypotension and ventilator were associated with a high incidence of AKI. The postulated mechanism was attributed to hemodynamic instability and, therefore, it predisposes impairment of circulation and consequent reduction of renal blood flow, with risk of evolution to hypoxic-ischemic AKI.<sup>15</sup>

It is observed that there were no post-renal causes detected in the AKI patients. According to Sinivasa et al where no post-renal causes were also detected and they explained that the chronic course and progressive deterioration of renal function in post-renal causes so the detection is usually in the pediatric ward than in PICU.<sup>7</sup> In our study, prerenal causes were more predominant (88.47%) and they are usually preventable. We have potential limitations in our study. It is a retrospective observational study so we depend on medical records to obtain the information where there were some missing values and information. In addition, a small sample size restricts the researcher in a certain range of findings.

## 5. CONCLUSION

Our study aimed to form a basis and focus on the burden of AKI in critically ill children. Higher incidence of AKI patients was in the failure stage on admission which indicates the importance of early identification of disease, by increasing awareness, adopting a unified definition and staging of AKI to guide quick intervention. Pre-renal causes, which were usually preventable and treatable, were prevalent in AKI Patients. Elaborating preventable and non-preventable risk factors will improve the prognosis of AKI. The ventilator is still the main concern of AKI in a critically ill child in SQUH. Nevertheless, hypotension, sepsis, and nephrotoxic medications are the main significant risk factors contributing to AKI. Such findings are worrisome, Pediatric intensivists should be alert, initiating measures to secure adequate renal perfusion. The Mortality rate was high in AKI patients, which requires raising awareness of AKI in the health care community.

## ETHICAL APPROVAL

Ethical approval was obtained from the Research and Ethics Committee, College of Medicine and Health Sciences, Sultan Qaboos University, Oman.

## Consent

As per international standard, parental written consent has been collected and preserved by the author(s).

## COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## REFERENCES

1. Thomas ME, Blaine C, Dawnay A, Devonald MA, Ftouh S, Laing C, Latchem S, Lewington A, Milford DV, Ostermann M. The definition of acute kidney injury and its use in practice. *Kidney International* 2015; 87, 62–73. doi:10.1038/ki.2014.328.
2. Mohd Ashraf, Naveed Shahzad, Mohd Irshad, Sheikh Quyoom Hussain, Parvez Ahmed. Pediatric acute kidney injury: A syndrome under paradigm shift. *Indian J Crit Care Med* 2014; 18(8): 518–526. doi: 10.4103/0972-5229.138156.
3. Khwaja, A. KDIGO Clinical Practice Guidelines for Acute Kidney Injury. *Nephron Clin Pract* 2012; 120(4): c179-84. DOI: 10.1159/000339789.
4. Shemesh, O, Golbetz, H, Kriss, J, Myers BD. Limitations of creatinine as a filtration marker in glomerulopathic patients. *Kidney Int* 1985; 28(5): 830-8. doi: 10.1038/ki.1985.205.
5. Bellomo R, Ronco C, Kellum JA, Mehta RL, Palevsky P, Acute Dialysis Quality Initiative workgroup. Acute renal failure - definition, outcome measures, animal models, fluid therapy and information technology needs: The Second International Consensus Conference of the Acute Dialysis Quality Initiative (ADQI) Group. *Crit Care* 2004; 8(4): R204–12. doi: 10.1186/cc2872.
6. Mehta, R, Cerdá, J, Burdmann E, Tonelli M, García-García G, Jha, V et al. International Society of Nephrology's 0by25 initiative for acute kidney injury (zero preventable deaths by 2025): a human rights case for nephrology. *Lancet* 2015; 385(9987): 2616-43. doi.org/10.1016/S0140-6736(15)60126-X.

7. Gupta S, Sengar G, Meti P, Lahoti A, Beniwal M, Kumawat M. Acute kidney injury in pediatric intensive care unit: Incidence, risk factors, and outcome. *Indian J Crit Care Med* 2016; 20(9): 526. doi: 10.4103/0972-5229.190368.
8. Srinivasa, S, Reshmavathi V. Incidence and etiology of acute kidney injury in children admitted to PICU using pRIFLE criteria. *Curr Pediatric Res* 2016; 20(1-2): 1–6.
9. Maqbool, J, Mir A, Bhat N, Moona W. Incidence and etiology of acute kidney injury in children admitted to PICU using pRIFLE criteria. *Int J of Contemp Pediatr* 2018; 5(3): 917. doi: 10.18203/2349-3291.ijcp20181513
10. Akcay A, Turkmen K, Lee D, Edelstein CL. Update on the diagnosis and management of acute kidney injury. *Int J Nephrol Renovasc Dis* 2010; 3: 129. doi: 10.2147/IJNRD.S8641.
11. Srinivasa S, Reshmavathi V., Srividya S. A comparison of pRIFLE and AKIN criteria for acute kidney injury in pediatric intensive care unit patients. *Int J Contemp Pediatrics* 2016; 3(2): 398–402. doi: 10.18203/2349-3291.ijcp20161022.
12. Rustagi RS, Arora K, Das RR, Pooni PA, Singh D. Incidence, risk factors and outcome of acute kidney injury in critically ill children - a developing country perspective. *Paediatr Int Child Health* 2017; 37(1): 35-41. doi: 10.1080/20469047.2015.1120409.
13. J Koeze, F Keus, W Dieperink, I C C van der Horst, J G Zijlstra, M van Meurs. Incidence, timing and outcome of AKI in critically ill patients varies with the definition used and the addition of urine output criteria. *BMC Nephrol* 2017 Feb 20;18(1):70. doi: 10.1186/s12882-017-0487-8.
14. Mehta P, Sinha A, Sami A, Hari P, Kalaivami M, Gulati A et al. Incidence of acute kidney injury in hospitalized children. *Indian pediatr* 2012; 49(7): 537-42. doi: 10.1007/s13312-012-0121-6.
15. Louzada CF, Ferreira AR. Evaluation of the prevalence and factors associated with acute kidney injury in a pediatric intensive care unit. *J Pediatr (Rio J)* 2021;97(4):426-432. doi: 10.1016/j.jpmed.2020.08.003.