
Short Communication

Pre - sowing seed treatment of chemicals, botanicals, plant growth regulators on growth, yield and yield attributing traits of linseed (*Linum usitatissimum* L.) under late sowing conditions var. Uma

*X. XXXXX, ¹ X. XXXX, ² X. X. XXXX AND ³ X. XXX

Division of XXXXXX, XXXX,

ABSTRACT

The experiment was conducted at Field Experimentation and Farm Research Centre, Department of Genetics and Plant Breeding, (Rabi season, 2020-21) Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (U.P.). The experimental material for present experiment comprised of thirteen priming treatments on linseed seed. The experiment was conducted in Randomized Block Design (RBD) with three replications. Linseed seeds of variety Uma was subjected to various pre sowing treatments like $MnSO_4$, $FeSO_4$ Pongamia leaf powder, Arappu leaf powder, Notchi leaf powder, Kinetin at various concentrations along with distilled water control. The study revealed that linseed seeds treated with Pongamia leaf powder T_6 (5%) showed maximum field emergence (97.48%), plant height 60 days (77.46cm) and 90 days (92.69cm) no of primary branches per plant (5.97), no of seeds per capsule (8.95), no of capsules per plant (25.91), test weight (9.43), seed yield per plant (4.78g), seed¹ yield per plot (128.4g), biological yield (8.26), harvest index (79.53) and reduced days to 50% flowering (65.89), days to maturity (115.89), followed by kinetin T_{12} 50(ppm), in comparison to control T_0 .

Article History

Received: 11.01.2020

Revised : 08.12.2020

Accepted: 17.12.2020

Keywords

Arappu leaf powder, $FeSO_4$, kinetin, $MnSO_4$, notchi leaf powder, pongamia leaf powder

Linseed (*Linum usitatissimum* L.) is an annual self-pollinated crop with origin in the Middle East or the Indian regions (Vavilov, 1951). The generic name *Linum* derives from Celtic word "lin" means "thread" and the species name *usitatissimum* (Latin word) means "very useful". *Linum usitatissimum*, only cultivated species of the genus *Linum* of family Linaceae is cultivated for oil from the beginning of agriculture eight thousand years ago and somewhat later for fibre (Zohary and Hopf, 1999). India ranks second in area after Canada in the world, but is at fourth place in term of production after Canada, China and U.S.A. In term of productivity, India (449kg/ha) is far below to Canada (1492 kg/ha), U.S.A (1484 kg/ha), Egypt (1465 kg/ha). (Alukedi *et al.*, (2021). Linseed is a *Rabi* crop in India which is a member of family Linaceae. Linseed (*Linum usitatissimum* L.) is an erect annual herbaceous plant 30-120 cm, in height with slender glabrous, grayish green stem. Flowers are showy, variously shaped regular, hermaphrodite, pentamerous, hypogynous and borne in loose terminal raceme or open cyme with blue, white or pink colour. *Linum usitatissimum* L. is the only species with non-dehiscent or semi-dehiscent capsules for modern cultivation of the family Linaceae. It contains up to 10 smooth, glossy an apple pip shaped, light brown color seeds which is 4-7 mm long. Linseed is predominantly self-pollinating. Natural cross pollination can occur at level of 6.75% by insects. (Alemayehu *et al.*, 2019). In recent years, the use of local botanicals has gained much importance, mainly among researchers, because of its high benefits in plant growth, yield and seed quality attributes. Some botanicals like Neem extract, Arappu leaf powder, Notchi leaf powder, Pongamia leaf powder are used for seed treatment and various effect of them can be seen on growth and yield of a plant. Botanical seed treatment is extracted from naturally occurring sources based on botanical ingredients. It has synergistic effect on early and uniform seed germination and enhances tolerance to pest and disease during early crop stage. (Mavi *et al.*, 2015). Hormonal priming can also be done to attain good plant growth and productivity. Hormonal priming refers to treating seeds in hormones such as GA₃, salicylic acid, ascorbate, kinetin, jasmonic acid etc. which helps in promoting seedling growth. The present study was conducted to evaluate the effect of pre-sowing seed treatment of chemicals, botanicals and plant growth regulators on plant growth and yield of linseed and to find out suitable pre-sowing seed treatment favorable for linseed crop.

MATERIALS AND METHODS

The experimental material for present investigation comprised of thirteen priming treatments on linseed variety (Uma). The experiment was laid out in Randomized Block Design (RBD) with three replications. Seeds of

(Uma) variety was obtained from CSAUAT Kanpur developed for high yield purpose and was subjected to various pre-sowing seed treatments like Mnso₄ T₁ (0.3%) and T₂ (0.5%), Feso₄ n T₃ (0.3%) T₄ (0.5%), Pongamia leaf powder T₅ (3%) T₆ (5%), Arappu leaf powder T₇ (3%) T₈ (5%) Notchi leaf powder T₉ (3%) T₁₀ (5%) Kinetin T₁₁ (25ppm) T₁₂ 50 (ppm) at various concentrations along with distilled water control T₀.

After pre-sowing seed treatments seeds were dried and then sowing was done in the field and following data was recorded field emergence percentage, plant height (cm), number of primary branches per plant, days to 50% flowering, days to maturity, number of capsules per plant, number of seeds per capsule, seed yield per plot (g), biological yield, harvest index, test weight in (g), seed yield per plant (g) data were collected from the field. The data was subjected to statistical analysis.

RESULTS AND DISCUSSION

All the treatments significantly affected growth and yield attributes. Here we will see about performance of various treatments depend on data which is recorded.

Table 1 Mean performance of field parameters of linseed


Treatments	Chemicals	FM	DFF	PH		NPBPP	DM	NCP	NSPC	TW	SYP P (g)	SYLOT (g)	BY	HI
				60 DAS	90DAS									
T₀	Control	88.63	77.89	70.89	75.12	4.29	128.67	18.41	6.94	5.98	2.12	48.6	4.56	38.90
T₁	MnSO ₄ (0.3%)	90.12	73.96	62.11	77.34	5.01	123.96	22.14	7.08	6.88	2.88	71.4	5.23	55.07
T₂	MnSO ₄ (0.5%)	92.34	76.55	65.35	80.58	5.41	124.16	23.45	7.51	7.06	2.96	73.8	6.69	44.25
T₃	Feso ₄ (0.3%)	89.14	70.12	72.94	88.17	4.88	120.12	18.46	8.04	8.14	2.69	65.7	5.34	50.37
T₄	Feso ₄ (0.5%)	93.54	69.35	73.21	89.44	4.98	119.35	19.87	8.28	8.41	2.94	73.2	7.48	39.30
T₅	Pongamia leaf powder (3%)	94.22	76.61	73.83	89.06	5.77	116.56	24.77	8.64	8.77	3.55	81.2	5.78	42.95
T₆	Pongamia leaf powder (5%)	97.48	65.67	77.46	92.69	5.97	115.89	25.91	8.95	9.43	4.78	128.4	8.26	79.53
T₇	Arappu leaf powder (3%)	93.64	72.96	60.91	76.14	4.66	122.96	25.33	8.26	8.16	3.35	85.5	7.78	43.06
T₈	Arappu leaf powder (5%)	94.11	74.65	63.42	78.65	4.84	124.65	25.86	8.46	8.89	3.65	91.5	7.26	42.98
T₉	Notchi leaf powder (3%)	90.89	69.24	58.66	73.89	5.11	119.24	21.54	7.99	7.56	2.98	74.4	7.66	38.90
T₁₀	Notchi leaf powder (5%)	91.49	70.84	61.71	76.94	5.36	120.84	23.55	8.14	8.61	3.19	80.7	7.14	44.68
T₁₁	Kinetin (25ppm)	93.47	74.33	65.43	80.66	5.44	124.33	19.38	8.31	8.36	3.06	76.8	7.31	41.86
T₁₂	Kinetin (50ppm)	95.55	66.56	74.86	90.09	5.59	127.61	24.88	8.74	9.06	3.85	121.5	8.01	78.72
	MEAN	92.82	72.29	66.83	82.06	5.18	122.18	22.27	8.10	8.10	3.27	83.05	6.71	49.86
	MIN	88.63	65.67	70.89	75.12	4.29	115.89	18.41	6.94	5.98	2.12	48.60	4.56	38.90
	MAX	97.48	77.89	77.46	92.69	5.97	128.67	25.91	8.95	9.43	4.78	128.40	8.26	79.53
	C.D @ (5%)	3.26	0.51	4.73	4.17	0.58	0.68	1.85	2.58	4.30	3.13	2.99	3.64	2.04
	SE(d)	1.58	0.25	2.29	2.02	0.28	0.33	0.90	1.25	2.08	1.52	1.45	1.76	0.99
	C.V.	2.60	3.81	7.64	7.64	0.45	3.76	2.64	0.58	0.94	0.69	10.69	1.15	11.25

Field emergence percentage

The maximum field emergence percent (97.48%) was recorded with pongamia leaf powder T₆ (5%), followed by (95.55%) kinetin T₁₂ 50 (50ppm). Whereas minimum field emergence

percent recorded with T₀ control (88.63%).

This experiment provided information that field emergence percent increases when linseed seeds are treated with pongamia leaf powder (5%) when compared with other treatments




Field emergence as influence by priming treatments on linseed

Days to 50% flowering

The minimum days to 50% flowering (65.89) recorded with pongamia leaf powder T₆ (5%), followed by T₁₂ kinetin (50PPM). The maximum days to 50% flowering recorded with (77.89)

T₀ control. This experiment provided information that days to 50% flowering reduced when linseed seeds treated with pongamia leaf powder (5%) in comparison with other treatments.




Days to 50% flowering as influence by priming treatments on linseed

Plant height at 60 DAS

The maximum plant height at 60 DAS (77.46 cm) is recorded with pongamia leaf powder T₆ (5%) followed by (74.86 cm) kinetin T₁₂ 50 (ppm). And minimum plant height at 60 days was recorded with

T₀ control (70.89cm).

This experiment provided information about plant height at 60 DAS increased when linseed seeds are treated with pongamia leaf powder (5%) in comparison with other treatments.




Plant height at 60 DAS as influence by priming treatments on linseed

Plant height at 90 DAS

The maximum plant height at 90 DAS (92.69cm) recorded with pongamia leaf powder T₆ (5%), followed by (90.09cm) kinetin T₁₂ (50ppm). Whereas minimum plant height (75.12cm) was recorded with T₀ control. The data collected was

statistically analysed and found significant difference among the triats. This experiment provided information that plant height at 90 DAS increased when linseed seeds treated with pongamia leaf powder (5%) in comparison to other treatments.




Plant height at 90 DAS as influence by priming treatments on linseed

Number of primary branches per plant

The maximum number of primary branches per plant (5.97) recorded with pongamia leaf powder T₆ (5%) followed by kinetin T₁₂ (50 ppm). And minimum number of primary branches per plant (4.29) was observed in T₀ control. Number of primary branches per plant data was collected

and statistically analysed found significant difference among the traits. This experiment provided information that number of primary branches per plant increased when linseed seeds treated with pongamia leaf powder (5%) in comparison with other treatments.




Number of primary branches per plant as influence by priming treatments on linseed

Days to maturity

The maximum number of days taken to maturity (128.67) recorded with T₀ control followed by (125.61) kinetin T₁₂ (50 PPM) and reduced days to maturity to minimum recorded with pongamia leaf

powder T₆ (5%). This experiment provided information that Days to maturity decreased when linseed seeds treated with pongamia leaf powder (5%) in comparison with other treatments.




Days to maturity as influence by priming treatments on linseed

Number of capsules per plant

The maximum number of capsules per plant (25.91) recorded with pongamia leaf powder T_6 (5%), followed by (24.84) kinetin T_{12} (50ppm). Whereas minimum number of capsules per plant (18.41) recorded with T_0 control.

This experiment provided information that no of capsules per plant increased when linseed seeds treated with pongamia leaf powder (5%) when compared with other treatments.




Number of capsules per plant as influence by priming treatments on linseed

Number of seeds per capsule

A range of 6.94 to 8.95 of number of seeds per capsule with 8.10 mean value were obtained. The maximum number of seeds per capsule (8.95) recorded with pongamia leaf powder T_6 (5%), followed by (8.74) kinetin T_{12} (50 PPM).

Whereas minimum number of seeds per capsule (6.94) recorded with T_0 control. This experiment provided information that no of seeds per capsule increased when linseed seeds treated with pongamia leaf powder (5%) when compared with other treatments.




Number of capsules per plant as influence by priming treatments on linseed

Test weight

The maximum test weight (9.43g) recorded with pongamia leaf powder T₆ (5%) followed by (9.06g) kinetin T₁₂(50ppm). Whereas minimum test weight was recorded with (5.98) T₀ control. This

experiment provided information that test weight increased when linseed seeds are treated with pongamia leaf powder (5%) in comparison with other treatments.




Test weight as influence by priming treatments on linseed

Seed yield per plant (g)

The maximum seed yield per plant (4.78g) recorded with pongamia leaf powder T₆ (5%), followed by (3.85g) kinetin T₁₂ (50 ppm). Whereas minimum seed yield per plant recorded with (2.12

g) T₀ control. This experiment provided information about seed yield per plant that it increased when linseed seeds treated with pongamia leaf powder (5%) in comparison to other treatments.




Seed yield per plant (g) as influence by priming treatments on linseed

Seed yield per plot (g)

The maximum seed yield per plot (128.4g) recorded with pongamia leaf powder T₆ (5%), followed by (121.5g) kinetin T₁₂ (50ppm). Whereas minimum seed yield per plot recorded with (48.6g)

T₀ control This experiment provided information that seed yield per plot increased when linseed seeds are treated with pongamia leaf powder (5%) in comparison with other treatments.




Seed yield per plot (g) as influence by priming treatments on linseed

Biological yield

The maximum biological yield recorded with (8.26) pongamia leaf powder T₆, followed by Kinetin T₁₂ 50 ppm (8.01). Whereas minimum biological yield recorded with (4.56) T₀ control This experiment

provided information that biological yield increased when linseed seeds treated with pongamia leaf powder (5%) in comparison with other treatments.




Biological yield as influence by priming treatments on linseed

Harvest index

The maximum harvest index (79.53) recorded with pongamia leaf powder T_6 (5%), followed by (78.72) kinetin T_{12} (50ppm) and minimum harvest index (38.90) recorded with T_0 control. This

experiment provided information about harvest index that it increased when linseed seeds are treated with pongamia leaf powder (5%) when compared with other treatments.



Harvest index as influence by priming treatments on linseed

CONCLUSION

It is concluded from the present study that pongamia leaf powder 5% treated with linseed seeds variety Uma can significantly affect growth, yield and yield attributing traits. It resulted in increase in field emergence (97.48%), plant height 60 days (77.46cm) and 90 days (92.69cm) no of primary branches per plant (5.97), no of seeds per capsule (8.95), no of capsules per plant (25.91), test weight (9.43), seed yield per plant (4.78g), seed yield per plot (128.4g), biological yield (8.26), harvest index (79.53). And lower down days to 50% flowering (65.67) and days to maturity (115.89). Followed by kinetin T_{12} (50ppm), in comparison with other treatments T_0 control.

ACKNOWLEDGEMENT

I express my sincere gratitude to all members of Seed science and technology, Department of Genetics and Plant Breeding, Sam Higginbottom University of Agriculture, Technology and Sciences, Naini, Prayagraj, Uttar Pradesh for providing all the necessary requirements and support for research.

REFERENCES

Alemayehu, Nigussie, and Gemechu Keneni. "Genetic Consequences of Failure to Control Outcrossing in Breeding Often Cross-Pollinated Pulse and Oil Crops"

Alukedi, Ahmed O, Ahmed A Almarie, Ma Alaloussi, Ss Farhan, And Alif Almehemdi. (2021). "Effect Of Laser Exposure As Pre Sowing Seed Priming In Three Flax Cultivars (*Linum Usitatissimum* L.)." *Plant Cell Biotechnology and Molecular Biology*: 43-52

Chormule, SR, NM Changade, and JB Patel. (2018). "Effect of seed treatments on storability of

vegetable seeds: a review." *Plant Archives*: 18(1): 28-32.

Mavi, Reza Kiani. (2015). "Green supplier selection: a fuzzy AHP and fuzzy ARAS approach." *International Journal of Services and Operations Management*: 22(2):165-188.

Mondal, Sananda, and Bandana Bose. (2019) "Impact of micronutrient seed priming on germination, growth, development, nutritional status and yield aspects of plants." *Journal of plant nutrition*: 42(19): 2577-2599.

Prabha, Deepti, Swati Negi, Poonam Kumari, Yogesh Kumar Negi, and JS Chauhan. (2016). "Effect of seed priming with some plant leaf extract on seedling growth characteristics and root rot disease in tomato." *International Journal of Agriculture System*: 4(1):46-51.

Prakash, M, G Sathiya Narayanan, B Sunil Kumar, and A Kamaraj. (2013). "Effect of seed hardening and pelleting on seed quality and physiology of rice in aerobic condition." *Agricultural Science Digest* 33(3):172-177.

Rasool, Tassadduq, Riaz Ahmad, and Muhammad Farooq. (2019). "Seed priming with micronutrients for improving the quality and yield of hybrid maize." *Gesunde Pflanzen* 71(1):37-44.

Rehman, H, Muhammad Kamran, Shahzad Maqsood Ahmed Basra, Irfan Afzal, and Muhammad Farooq. (2015). "Influence of seed priming on performance and water productivity of direct seeded rice in alternating wetting and drying." *Rice Science* 22(4):189-196.

Sajjan, Ashok S, MS Dhanelappagol, and RB Jolli. (2017). "Seed quality enhancement through seed priming in pigeonpea [*Cajanus cajan* (L.)

- Millsp.]. " *Legum Res* 40(1):173-177.
- Singh, Harmeet, Rupinder Kaur Jassal, JS Kang, SS Sandhu, Harrajdeep Kang, and Kamaljit Grewal. (2015). "Seed priming techniques in field crops- A review." *Agricultural Reviews* 36(4):
- Umesha, BC, K Bhanuprakash, J Lakshmi, and Soumya S Desai. (2017). "Influence of Seed Fortification on Crop Growth and Seed Yield in Cluster Bean [*Cyamopsis tetragonoloba* (L.) Taub.] var Pusanavbahar." *Environment & Ecology* 35(3C):2232-2236
- Vakeswaran, V, A Vijayakumar, and R Jerlin. (2003). "Effect of Seed Priming in Pea (*Pisum sativum* L.) Seeds." *Seed Technology*: 203-203.
- Vavilov, Nikolai Ivanovich. (1951). *The origin, variation, immunity and breeding of cultivated plants*. Vol. 72: LWW.
- Zohary, Daniel. (1999). "Monophyletic vs. polyphyletic origin of the crops on which agriculture was founded in the Near East." *Genetic Resources and Crop Evolution* 46(2):133-142.