

Seed yield and economic returns in coriander in relation to different nutrient combinations.

ABSTRACT

Organic manures along with the chemical fertilizers tend to reduce the total cost of cultivation and supplement the crop with the essential nutrients for growth and development of the plant. In view of this an experiment was conducted in coriander during winter season of 2012-13 at college of agriculture, Jabalpur, Madhya Pradesh to assess the effect of organic and inorganic sources of nutrient with different levels on various parameters of coriander. Results with regard to various levels of organic manures shows that poultry manure 5 t ha^{-1} produced the maximum number of umbels per plant (33.67), the maximum number of umbellets per umbel (7.06), maximum number of seeds per umbel (33.69), maximum weight of seeds per umbel (0.44g), maximum number of seeds per plant (603.75) and significant maximum seed yield per plant (6.09 g), seed yield per plot (1.85 kg) and seed yield per hectare (17.30 q). Results in relation to fertilizer levels shows that the maximum number of umbels per plant (33.46), maximum number of umbellets per umbel(7.04) , maximum number of seeds per umbel (33.29), maximum weight of seeds per umbel (0.45 g), maximum seed yield per plant (6.01g), yield per plot (1.87kg) and seed yield per hectare (17.53 q) were recorded with 100 % RDF. Results in relation to interaction effect of different treatments shows that, the significant maximum number of umbels per plant (36.27), maximum number of umbellets per umbel(7.71), maximum number of seeds per umbel (36.74), maximum weight of seeds per umbel (0.49 g), maximum number of seeds per plant (701.97) and the significant maximum seeds yield per plant (7.14g), seed yield per plot (2.04 kg) and seed yield hectare (19.16 q) were observed in poultry manure $5 \text{ t ha}^{-1} + 100 \%$ RDF. The highest net return and cost benefit ratio 1: 2.98 with treatment combination T₅ (Poultry manure $5 \text{ t ha}^{-1} + 100 \%$ RDF).

Key words: Coriander , seed yield, economic returns, organic manures and fertilizers.

INTRODUCTION

Coriander (*Coriandrum sativum*) is an annual herb which possesses nutritional and medicinal properties in the family Apiaceae (Sharma and Sharma, 2012). Coriander is originated from the Eastern Mediterranean region. Coriander is one of the most commonly used spices (Leena *et al.*, 2012). It is one of the important seed spices occupying a prime position throughout the globe to add taste, flavor and pungency in various food items. It is a multipurpose herb grown mainly for its foliage and seeds (Burdock and Carabin, 2009). Since the coriander seeds have strong and typical scent, they are appreciated worldwide as basic ingredients of many traditional foods, particularly curry powder (Mahendra and Bisht, 2011; Sahib *et al.*, 2013). Moreover, it is a frequent ingredient in the preparation of ayurvedic medicines and is a traditional home therapy for different ailments viz., rheumatism, joint pain, gastrointestinal, complaints, flatulence, indigestion, insomnia, convulsions, anxiety, loss of appetite (Emamghoreishi *et al.*, 2005), etc. Health benefits of coriander include treatment of swellings, diarrhoea, mouth ulcers, anaemia, digestion, menstrual disorders, small pox, eye care, conjunctivitis, skin disorders, blood sugar disorders, protects and soothes liver etc (Diwan *et al.*, 2018). Green leaf is having high essential oil with good aroma and flavour (Lal *et al.*, 2020). The essential oil content in seeds of coriander is 0.47–0.53 per cent whereas major chemical constituents of essential oil are linalool, linalyl acetate along with other important constituents such as thymol, geraniol, carophyllene and pinene. (Lal *et al.*, 2020).

In India, coriander is mainly cultivated in Rajasthan, Madhya Pradesh, Uttar Pradesh and Southern States like Andhra Pradesh, Karnataka and Tamil Nadu. It

takes approximately 100 days for maturity whereas the smaller-seeded coriander takes a longer growing period. The spice is an important item of international trade. Producers thinking about growing coriander should investigate the current pricing of the crop as prices can vary. Due to global competition, the Canadian coriander is uniform in size with good seed quality and reported more competent for export market. In India the area under coriander is increasing as per our domestic demand. The increase in overall production of coriander seeds in India is mainly due to the advancement of production technology through the National Agriculture Research System and educating the farmers through various training programmes. Introduction of high yielding varieties, new production techniques, integrated nutrient management and sowing of crops as per soil/ land suitability are the major production factor for the higher productivity. Application of fertilizer has been documented to enhance plant growth and development. Particularly nitrogen (N) is one of the greatest production inputs. Nitrogen is an essential nutrient in creating the plant dry matter, as well as many energy-rich compounds that regulate photosynthesis and plant production. Comparisons between inorganic and organic fertilizer-N sources are hard to perform since there is usually a dramatic difference in N availability from these two sources of N. Comparisons on the basis of similar amounts of total N applied are therefore of limited relevance to agricultural practice, whereas comparisons on the basis of similar N availability are hindered by the lack of reliable nitrogen release estimates for organic fertilizer sources derived from animal manures (Van Kessel and Reeves, 2002). Organic fertilizers in comparison with the chemical one have lower nutrient content and are slow to release but they are as effective as chemical fertilizers over longer periods of use (Sharafzadeh and Ordoorkhani, 2011). Organic manure must be added to conventional NPK fertilizer to improve soil structure, make the soil easier to cultivate, encourage root development, provide with nutrients and enable their

increased uptake by plants. The application of nutrients through various sources is newer approach and being advocated for sustainable production. The integrated nutrient management has a crucial role in improving the plant physiology characters that builds levels of resistance and reduces the incidence of disease and pest attacks (Mirchandani and Mirchandani, 2005). Therefore, present study was taken to develop a suitable nutrient management practice for coriander crop adopting organic manures.

MATERIALS AND METHODS

The field experiment was conducted at Horticulture complex Department of Horticulture, JNKVV, Jabalpur, Madhya Pradesh during the winter seasons (Rabi) of 2012-2013. The Design of experiment was Randomized complete block design (RCBD) with 3 replications. The experiment consisted of 12 treatments T1-FYM @ 20 t/ha + 100 % RDF , T2-FYM @ 20 t/ha + 50 % RDF, T3-FYM @ 10 t/ha + 100 % RDF, T4-FYM @ 10 t/ha + 50 % RDF, T5-Poultry manure @ 5 t/ha + 100 % RDF , T6-Poultry manure @ 5 t/ha + 50 % RDF, T7- Poultry manure @ 2.5 t/h + 100 % RDF, T8- Poultry manure @ 2.5 t/ha + 50 % RDF, T9-Vermicompost @ 5 t/ha + 100 % RDF, T10-Vermicompost @ 5 t/ha + 50 % RDF, T11-Vermicompost @ 2.5 t/ha +100% RDF, T12-Vermicompost @ 2.5 t/ha +50 % RDF. Proper leveling was done in order to facilitate the irrigation. Layout of the experiment was done as per plan of the investigation and treatments as given. The well decomposed FYM, Poultry manure and Vermicompost was applied in required plots before sowing of seeds. It was mixed well in each plot by light ploughing. Half amount of N with full amount of P and K were given per plot as basal dose and rest amount of N was given as top dressing after 40 days of sowing. Prior to sowing coriander variety JD-1 seeds were split into two halves by rubbing seeds were treated with thiram @ 2 g/kg of seeds was done thoroughly

against seed borne diseases. The sowing of seeds was done as per treatments in prepared plots. Seeds were sown at 30 cm row spacing apart. The first light irrigation was provided just after sowing and subsequent irrigations were given at 15-20 days intervals to maintain the soil moisture till crop maturity. All the cultural operations were done as and when required for good crop stand. Other operations were done as per operation schedule. Precautionary two sprays of Dithane M-45 (0.03%) were done against diseases in the main field. Similarly two sprays of Rogor (0.03%) were done against insects in the main field as precautionary measure. Due to large population of plants in the plots, it was rather difficult to record the observation in each plant in the experiment field. Since, all the plants have equal opportunity for their growth and development. Therefore, a technique of random sampling was adopted and a sample of five plants from each plot was drawn at random to record the yield and yield attribute. The following growth, yield and quality components (Number of umbels per plant, Number of umbellets per umbel, Number of seeds per umbel, Weight of seeds per umbel (g), Number of seeds per plant, Seed yield per plant (g), Seed yield per plot (kg), Seed yield per hectare) were taken into observation and they were subjected to statistical analysis in order to draw the valid conclusion on the basis of C.D. value. To find out the economic feasibility of various treatment combinations, economics of the various treatments were worked out by calculating parameters like cost of cultivation, gross returns, net returns and benefit cost ratio using the prevailing price of inputs and outputs in local market.

RESULT AND DISCUSSION.

EFFECT OF VARIOUS LEVELS OF ORGANIC MANURES AND FERTILIZERS ON YIELD AND YIELD PARAMETERS.

The application of organic manures significantly improves the various yield parameters and consequently yield viz., number of umbels per plant, number of umbelets per umbel, number of seeds per umbel, weight of seeds per umbel, seed yield per plant (g), seed yield per plot (kg) and seed yield per hectare ($q\ ha^{-1}$) of coriander.

NUMBER OF UMBELS PER PLANT .

Number of umbels per plant of coriander were significantly affected by various organic manures. Among the organic manures, poultry manure @ $5\ t\ ha^{-1}$ produced the maximum number (33.67) of umbels per plant followed by vermicompost @ $5\ t\ ha^{-1}$ (32.33). The least number (30.57) of umbels per plant was recorded in FYM @ $10\ t\ ha^{-1}$. Poultry manure @ $5\ t\ ha^{-1}$ was significantly superior over vermicompost and FYM. The present findings are in accordance with Ibrahim *et al.*, (2006) and El- Mekawey *et al.*, (2010). This may be due to poultry manure is due to enriched nitrogen content as compared to farmyard manure and compost. Nitrogen, phosphorus and potash are the three major nutrient required for good production or yield of coriander. Increasing seed yield of coriander ultimately depends on good yield contributing characters, which results from proper dose of fertilizers. Number of umbels per plant were significantly influenced with regard to application of fertilizer levels. The maximum number (33.43) of umbels per plant were observed in the application of 100 % RDF and the minimum (30.31) in the application of 50 % RDF, 100 % RDF was found to be significant over 50 % RDF. The present findings are in accordance with Singh and Jat (2002) ; Okut and Ydrm (2005); Kumar *et al.* (2007); Nagar *et al.*, (2009); Nayak *et al.* (2009); Jan *et al.*, (2011) and Khalid (2012). Fikadu *et al.*, (2019) observed maximum number of umbels plant⁻¹ (17.93) with treatment N ($60\ kg\ ha^{-1}$) and P ($50\ kg\ ha^{-1}$). In case of interaction effects, number of umbels per plant

was significantly influenced by the treatment combinations. The maximum number (36.27) of umbels per plant were recorded with the application of poultry manure @ 5 t ha⁻¹ + 100 % RDF followed by vermicompost @ 5 t ha⁻¹ + 100 % RDF (34.13). The least number (29.03) of umbels per plant were recorded in FYM @ 10 t ha⁻¹ + 50% RDF. Poultry manure @ 5 t ha⁻¹ + 100 % RDF was significant over all other interactions. The present findings corroborated the results of Choudary *et al.* (2011) for number of pods per plant in fenugreek. The increase in number umbels per plant might be due to increased supply of major plant nutrients that are required in larger quantities for growth and development of plants. Nitrogen accelerates the growth, development reproductive phases and protein synthesis in plants, thereby promoting higher number of umbels per plant. It's that balanced fertilizer application improved the availability of phosphorous, nitrogen and other nutrients, helped in increasing the yield attributing characters (Mounika *et al.*, 2018).

NUMBER OF UMBELLETS PER UMBEL .

Number of umbellets per umbel were not found to be significantly influenced by the treatments with various organic manures. However, Poultry manure @ 5 t ha⁻¹ produced the maximum number (7.06) of umbellets per umbel followed by vermicompost @ 5 t ha⁻¹ (6.82). Whereas the least number (6.37) of umbellets per umbel were recorded in FYM @ 10 t ha⁻¹. Number of umbellets per umbel were significantly influenced with regard to fertilizer levels. The maximum number of umbellets per umbel (7.04) were found with 100 % RDF and the minimum (6.27) with 50 % RDF. 100 % RDF is found to be significant over 50 % RDF. The present findings are in propinquity with Kumar *et al.*, (2007). Fikadu *et al.*, (2019) observed maximum number of umbellate umbel⁻¹ (6.69), with treatment N (60kg ha⁻¹) and P (50 kg ha⁻¹) instead of highest dose of both nutrients. But,

umbellate per umbel interaction effect was found to be non-significant by Pooja *et al.*, (2017). The interaction effect on number of umbellets per umbel was observed to be non significant. However, the maximum number (7.71) of umbellets per umbel were recorded with poultry manure @ 5 t ha⁻¹ + 100 % RDF followed by vermicompost @ 5 t ha⁻¹ + 100 % RDF (7.11). The least number (5.97) of umbellets per umbel was recorded in FYM @ 10 t ha⁻¹ + 50 % RDF.

NUMBER OF SEEDS PER UMBEL .

Number of seeds per umbel was significantly improved by various levels of organic sources. Among the organic sources, poultry manure @ 5 t ha⁻¹ recorded the maximum number (33.69) of seeds per umbel followed by vermicompost @ 5 t ha⁻¹ (32.52). However, the least number (29.85) of seeds per umbel were recorded in vermicompost @ 2.5 t ha⁻¹. Effect of poultry manure @ 5 t ha⁻¹ was significantly superior over vermicompost and FYM. Number of seeds per umbel was significantly influenced with regard to fertilizer levels. The maximum number (33.29) of seeds per umbel were recorded in 100 % RDF and the minimum (29.33) in 50 % RDF. 100 % RDF was found to be significant over 50 % RDF. The present findings are in accordance with Singh and Jat (2002) and Nayak *et al.*, (2009). Fikadu *et al.*, (2019) observed significantly maximum number of seeds umbel⁻¹ (18.34), with treatment N (60kg ha⁻¹) and P (50 kg ha⁻¹). But, seeds per umbel interaction effect was found to be non-significant by Pooja *et al.*, (2017). The interaction effects on number of seeds per umbel were found to be significant. Significantly maximum number (36.74) of seeds per umbel were recorded in poultry manure @ 5 t ha⁻¹ + 100 % RDF followed by vermicompost @ 5 t ha⁻¹ + 100 % RDF (34.14). The least number (27.73) of seed per umbel were recorded in vermicompost @ 2.5 t ha⁻¹ + 50% RDF. These findings are quite similar findings were reported by Choudary *et al.*, (2011) for seeds per pod in

FYM @ 20 t/ ha	32.67	30.73	31.70	6.99	6.29	6.64	33.07	29.07	31.07	33.07	29.07	31.07	0.47	0.34	0.41
FYM @ 10 t/ ha	32.10	29.03	30.57	6.78	5.97	6.37	31.43	28.33	29.88	31.43	28.33	29.88	0.42	0.33	0.37
PM @ 5 t/ ha	36.27	31.07	33.67	7.71	6.40	7.06	36.74	30.63	33.69	36.74	30.63	33.69	0.49	0.39	0.44
PM @ 2.5 t/ ha	33.33	30.73	32.03	6.87	6.13	6.50	32.37	29.33	30.85	32.37	29.33	30.85	0.45	0.36	0.40
VC @ 5 t/ ha	34.13	30.53	32.33	7.11	6.54	6.82	34.13	30.90	32.52	34.13	30.90	32.52	0.48	0.38	0.43
VC @ 2.5 t/ ha	32.07	29.77	30.92	6.81	6.20	6.51	31.97	27.73	29.85	31.97	27.73	29.85	0.42	0.32	0.38
MEAN	33.43	30.31	31.87	7.04	6.26	6.65	33.29	29.33	31.31	33.29	29.33	31.31	0.45	0.35	0.40
	(OM)	(RDF)	(OM) x (RDF)	(OM)	(RDF)	(OM) x (RDF)	(OM)	(RDF)	(OM) x (RDF)	(OM)	(RDF)	(OM) x (RDF)	(OM)	(RDF)	(OM) x (RDF)
SEm±	0.259	0.150	0.367	NS	0.162	NS	0.355	0.205	0.502	0.355	0.205	0.502	0.041	0.024	0.058
CD @ 5%	0.760	0.439	1.075	-	0.475	-	1.042	0.601	1.473	1.042	0.601	1.473	-	0.070	-

FYM- Farmyard manure, PM- Poultry manur, VC- Vermicompost , OM- organic manures, RDF- recommended dose of fertilizers, DAS- Days after sowing.




Fig 1. Graph representing variation in number of umbels per plant and number of seed per umbel due to various treatment interactions.

NUMBER OF SEEDS PER PLANT .

Number of seeds per plant was found to be significantly influenced by various organic manures. Among the organic manures, poultry manure @ 5 t ha⁻¹ produced the highest number of seeds per plant (603.75) followed by vermicompost @ 5 t ha⁻¹(550.44). The least number of seeds per plant (488.54) was noted in FYM @ 10 t ha⁻¹ . Poultry manure @ 5 t ha⁻¹ were significant over vermicompost and FYM. Number of seeds per plant were significantly influenced with regard to fertilizer levels. The maximum number of seeds per plant (590.48) were recorded with 100 % RDF and the minimum (475) in 50 % RDF. 100 % RDF was observed to be significant over 50 % RDF . The present findings are in propinquity with Manure *et al.*, (2000) and Channabasavanna *et al.*, (2002). The interaction effect on number of seeds per plant was found to be significant. The maximum number of seeds per plant (701.97) were found with poultry manure 5 t ha⁻¹ + 100 % RDF followed by vermicompost @ 5 t ha⁻¹ + 100 % RDF (598.34). While it was observed to be minimum (440.51) in FYM @ 10 t ha⁻¹ + 50% RDF. Application of poultry manure @ 5 t ha⁻¹ +100 % RDF were significant over all the other interactions .

SEED YIELD PER PLANT .

Seed yield per plant (g) was influenced significantly by the use of various organic manures . Among the organic manures, poultry manure @ 5 t ha⁻¹ produced the maximum seed yield per plant (6.09g) followed by vermicompost @ 5 t ha⁻¹ (5.53 g). Whereas, the minimum seed yield per plant (4.25 g) was recorded in FYM @ 10 t ha⁻¹ . Effect of poultry manure @ 5 t ha⁻¹ was significant over

vermicompost and FYM . The present findings are in accordance with El-Mekawey *et al.*, (2010). Seeds yield per plant was significantly influenced by fertilizer levels. The highest seed yield per plant (6.01 g) were recorded in 100 % RDF and minimum (4.25g) in 50 % RDF. 100 % RDF was observed to be significant over 50 % RDF . The present findings are in accordance with Manure *et al.*, (2000); Kumar *et al.*, (2007); Nayak *et al.*, (2009) and Khalid (2012). Fikadu *et al.* (2019) recorded maximum, seed yield plant⁻¹ (4.01 g) with treatment N (60kg ha⁻¹) and P (50 kg ha⁻¹) instead of highest dose of both nutrients .The interaction effect on seed yield per plant was found significant. The maximum seed yield per plant (7.14 g) was recorded in poultry manure @ 5 t ha⁻¹ + 100 % RDF followed by vermicompost @ 5 t ha⁻¹ + 100 % RDF (6.72 g) and the minimum seed yield per plant (3.62 g) was recorded in FYM @ 10 t ha⁻¹ + 50 % RDF. However, critical difference between poultry manure @ 5 t ha⁻¹+ 100 % RDF and vermicompost @ 5 t ha⁻¹ + 100 % RDF was found at par .

SEED YIELD PER PLOT AND SEED YIELD PER HECTARE .

Seed yield per plot and seed yield per hectare was influenced significantly by the use of various organic manures. It was observed that among the organic manures, poultry manure @ 5 t ha⁻¹ produced the maximum seed yield per plot (1.85 kg) and seed yield (17.07 q ha⁻¹) followed by vermicompost @5 t ha⁻¹(1.82 kg and 17.07 q ha⁻¹ respectively). The least seed yield per plot (1.49kg) and seed yield (14.02 q ha⁻¹) was noted in FYM @ 10 t ha⁻¹. However, difference in between poultry manure and vermicompost was non significant and which were at par . To standardize the rate of three major nutrients nitrogen, phosphorus and potash , studies have shown that, significant maximum seed yield per plot (1.87 kg) and seed yield per hectare (17.53 q) was found in 100 % RDF and the minimum (1.53 kg and 14.30 q respectively) in RDF 50 % . 100 % RDF was

found to be significant over 50 % RDF . Similar results are reported by Manure *et al.*, (2000); Naghera *et al.*, (2000); Singh *et al.*, (2000); Singh and Jat (2002); Channabasavanna (2002); Kumar *et al.*, (2002); Garg *et al.*, (2004); Gujar *et al.*, (2005); Tripathi (2006); Akbarinia *et al.*, (2006); Oliveira *et al.*, (2006); Kumar *et al.*, (2008) and Nagar *et al.*, (2009). Fikadu *et al.*, (2019) recorded significantly maximum seed yield plot⁻¹ (341g) and seed yield ha⁻¹ (12.36 q) with treatment N (60kg ha⁻¹) and P (50 kg ha⁻¹) instead of highest dose of both nutrients. Based on the interaction effects on the maximum seed yield per plot (2.04 kg) and seed yield per hectare (19.16 q) were recorded with poultry manure @ 5 t ha⁻¹ + 100 % RDF followed by vermicompost @ 5 t ha⁻¹ + 100 % RDF (1.98kg and 18.53q respectively). While the minimum seed yield per plot (1.27 kg) and seed yield per hectare (11.90 q) was recorded in FYM @ 10 t ha⁻¹ + 50% RDF. However, difference in between poultry manure 5 t ha⁻¹ and vermicompost @ 5 t ha⁻¹ was not found to be significant and values were at par. Similar results have also been reported by Mohamed and Abdu (2004); Sadanandan and Hamza (2006) in black pepper. Aishwath *et al.*, (2010); Choudary *et al.*, (2011) in fenugreek and Jan *et al.*, (2011). The probable reason for enhanced seed yield might be due to cumulative effects of nutrient (macro and micro) on vegetative growth which ultimately led to more photosynthetic activities while, application of organic, inorganic and bio-fertilizers enhance carbohydrate and nitrogen metabolism of pectic substances, as well as improve the water metabolism and water relation in the plants.

Table.2. Number of seeds per plant, seed yield per plant (g), seed yield per plot (kg) and seed yield per hectare (q) with regard to various treatments of organic manures and fertilizer levels.

ORGANIC MANURES	NUMBER OF SEEDS PER PLANT	SEED YIELD PER PLANT (G)	SEED YIELD PER PLOT (KG)	SEED YIELD PER HECTARE (Q)
-----------------	---------------------------	--------------------------	--------------------------	----------------------------

	100% RDF	50% RDF	Mean	100% RDF	50% RDF	Mean	100% RDF	50% RDF	Mean	100% RDF	50% RDF	Mean
FYM @ 20 t/ha	591.37	475.65	533.51	5.78	4.50	5.14	1.90	1.59	1.74	17.81	14.89	16.35
FYM @ 10 t/ha	536.49	440.59	488.54	4.89	3.62	4.25	1.72	1.27	1.49	16.14	11.90	14.02
PM @ 5 t/ha	701.97	505.53	603.75	7.14	5.04	6.09	2.04	1.65	1.85	19.16	15.43	17.30
PM @ 2.5 t/ha	571.76	480.02	525.89	6.27	4.13	5.20	1.82	1.52	1.67	17.08	14.20	15.64
VC @ 5 t/ha	598.34	502.54	550.44	6.72	4.34	5.53	1.98	1.67	1.82	18.53	15.61	17.07
VC @ 2.5 t/ha	542.92	440.51	491.72	5.23	3.88	4.56	1.75	1.47	1.61	16.43	13.77	15.10
MEAN	590.48	474.14	532.31	6.01	4.25	5.13	1.87	1.53	1.70	17.53	14.30	15.91
	(OM)	(RDF)	(OM) x (RDF)	(OM)	(RDF)	(OM) x (RDF)	(OM)	(RDF)	(OM) x (RDF)	(OM)	(RDF)	(OM) x (RDF)
SEm±	5.101	2.945	7.213	0.206	0.119	0.291	0.019	0.011	0.028	0.177	0.102	0.300
CD @ 5%	14.960	8.637	21.157	0.604	0.349	0.855	0.057	0.033	0.081	0.520	0.251	0.735

FYM- Farmyard manure, PM- Poultry manur, VC- Vermicompost , OM- organic manures, RDF- recommended dose of fertilizers.




Fig 2. Graph representing variation in seed yield due to various treatment interactions.

ECONOMICS OF VARIOUS TREATMENTS.

Coriander is one of the most important economic crops (Khan *et al.*, 2019). To obtain an excellent economic return, optimum fertilizer application has great importance. Accordingly, the study found that the use of RDF (100%) through fertilizers and combinations of different organic sources produces the maximum grain yield, net return, and benefit cost ratio in coriander production.

Cost of cultivation (Rs./ha):

The cost of cultivation was directly associated with various inputs *viz.* cost of chemical fertilizers, FYM, poultry manure and vermicompost. The highest cost of cultivation (Rs 57650.6 /ha.) was calculated under T9 (Vermicompost @ 5 t ha⁻¹+ 100 % RDF) followed by T10 (Vermicompost @ 5 t ha⁻¹ + 50 % RDF) (Rs-56825.3 /ha.). While the minimum cost of cultivation (Rs 43025.30 /ha.) was recorded by the use of T8 (Poultry manure @ 2.5 t ha⁻¹+ 50 % RDF).

Gross income (Rs./ha):

The maximum gross income (Rs 134120.00 /ha.) was calculated in the treatment T5 (Poultry manure @ 5 t ha⁻¹ + 100 % RDF) followed by T9 (Vermicompost @ 5 t ha⁻¹ + 100 % RDF) with (Rs 129710.00 /ha). While the minimum (Rs 89810.00 /ha.) was found in the treatment T4- (FYM @ 10 t ha⁻¹ + 50 % RDF)

Net profit (Rs. /ha):

The maximum net profit (89069.4 Rs/ha.) was obtained in T5 (Poultry manure @ 5t ha⁻¹+ 100 % RDF) which is the best as compared to other treatments followed by T1(FYM @ 20 t ha⁻¹+ 100 % RDF) (Rs 79419.4 Rs ha⁻¹), however the minimum net income (Rs 46684.7/ha) was obtained in treatment T4 ((FYM @ 10 t ha⁻¹+ 50 % RDF).

Cost: Benefit ratio:

Application of different fertilizers alone leads to the lowest net profit and benefit cost ratio. However, when fertilizers are combined during application, the profit is increased. The maximum cost: benefit ratio (1: 2.98) was noted under the treatment T5 (Poultry manure @ 5 t ha⁻¹+ 100 % RDF) followed by T1 (FYM @ 20 t ha⁻¹ + 100 % RDF) (1:2.76). However, the minimum cost: benefit ratio (1: 1.95) was recorded from T12 (Vermicompost @ 2.5 t ha⁻¹ +50 % RDF). Which is closely followed by 50 % RDF through fertilizers and 50 % RDF through vermicompost (Godara *et al.*, 2014). Fikadu *et al.*, (2019) found that the use of RDF (100%) through fertilizers and combinations of different organic and inorganic sources produces the maximum grain yield (1024 kg ha⁻¹), net return (Rs. 59556 ha⁻¹) and benefit cost ratio (3.66) in coriander production. Jhankar *et al.*, (2017) observed, the maximum net profit (Rs. 107689 ha⁻¹) and benefit: cost (2.09) in the treatment of inorganic fertilizer (100% of RDF), bio-fertilizers;

azotobacter, azospirillum and PSB (2.5 kg ha^{-1} and farmyard manure (5 t ha^{-1})). Similarly, it's also reported that 60 kg N ha^{-1} (half dose of nitrogen as basal and remaining half dose at 30 days after sowing) and $60 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ (full doses of phosphorus as basal) is optimum for higher production & net returns from coriander (Javiya *et al.*, 2017). Therefore, integrated nutrient application is very important in improving the physicochemical and biological conditions of the soils and finally helped in increasing the net profit by maximizing benefit cost ratio (Tripathi *et al.*, 2013; Jhankar *et al.*, 2017; Pooja *et al.*, 2017).

Table .3. Economics of crop in relation to different treatments.

CONCLUSION.

The study shows that both the nutrient sources i.e. organic manures and inorganic

Symbol	Treatments	Yield (q/ha.)	Cost of cultivation (Rs. /ha.)	Gross income (Rs. /ha.)	Net income (Rs./ha.)	Cost : Benefit Ratio
T ₁	FYM 20 t/ ha + 100 % RDF	17.81	45250.6	124670	79419.4	1:2.76
T ₂	FYM 20 t/ ha + 50 % RDF	14.92	44425.3	104440	60014.7	1:2.35
T ₃	FYM 10 t/ ha + 100 % RDF	16.14	43950.6	112980	69029.4	1:2.57
T ₄	FYM 10 t/ ha + 50 % RDF	12.83	43125.3	89810	46684.7	1:2.08
T ₅	Poultry manure 5 t/ ha + 100 % RDF	19.16	45050.6	134120	89069.4	1:2.98
T ₆	Poultry manure 5 t/ ha + 50 % RDF	15.43	44225.3	108010	63784.7	1:2.44
T ₇	Poultry manure 2.5 t/h +100 % RDF	17.08	43850.6	119560	75709.4	1:2.73
T ₈	Poultry manure 2.5 t/ ha + 50 % RDF	14.2	43025.3	99400	56374.7	1:2.31
T ₉	Vermicompost 5 t/ ha + 100 % RDF	18.53	57650.6	129710	72059.4	1:2.25
T ₁₀	Vermicompost 5 t/ ha + 50 % RDF	15.61	56825.3	109270	52444.7	1:1.92
T ₁₁	Vermicompost 2.5 t/ ha +100 % RDF	16.43	50150.6	115010	64859.4	1:2.29
T ₁₂	Vermicompost 2.5 t/ ha +50 % RDF	13.77	49325.3	96390	47064.7	1:1.95

fertilizers responded well in terms of yield and yield parameters. The interaction of both the nutrient sources showed significant effect on yield and economics. It is concluded that the application of poultry manure @ 5 t ha⁻¹ + 100 % RDF (50:30:60 kg NPK/ ha) recorded the maximum seed yield of coriander variety JD-1 along with highest net profit and cost benefit ratio 1:2.98 followed by FYM @ 20 t ha⁻¹ + 100% RDF . Hence present findings have practical utility in successful cultivation of coriander for Kymore pleatue and Satpura hills of Madhya Pradesh.

REFERENCES.

Aishwath, O. P., Mehta, R. S. and Anwer, M. M. (2010). Integrated nutrient management in seed spice crops. *Indian Journal of Fertilisers*.6 (11): 132-139.

Akbarinia, A., Daneshian, J. and Mohmmadbiegi, F. (2006). Effect of nitrogen fertilizer and plant density on seed yield, essential oil and oil content of *Coriandrum sativum* L. [Persian] *Iranian Journal of Medicinal and Aromatic Plants*. 22 (4): Pe410-Pe419.

Burdock, G. A. and Carabin, I. G. (2009). Safety Assessment of Coriander (*Coriandrum sativum* L.). Essential Oil as a Food Ingredient. *Food and Chemical Toxicology*, 47, 22-34.

Channabasavanna, A. S. (2002). Standardization and economic analysis of fertilizer levels for coriander (*Coriandrum sativum* L.). *Journal of Maharashtra Agricultural Universities*. 27 (2): 160-162.

Channabasavanna, A. S., Yalamali S. G. and Biradar, D. P. (2002). Nutrient requirement of coriander in Tungabhadra project area of Karnataka . *Journal of Maharashtra Agricultural Universities*. 27(1): 38-39.

Choudhary, B.R., Gupta A.K., Parihar C.M., Jat S.L.and Singh D.K. (2011). Effect of integrated nutrient management on fenugreek (*Trigonella foenumgraecum*) and

its residual effect on fodder pearl millet (*Pennisetum glaucum*) Online published on 3 December.

Choudhary, G. R., Jain, N. K. and Jat, N. L. (2008). Response of coriander (*Coriandrum sativum*) to inorganic nitrogen, farmyard manure and biofertilizer. *Indian Journal of Agricultural Sciences*. 78 (9): 761-763 .

Diwan, G., Bisen, B. P. and Maida, P. (2018). Effect of nitrogen doses and row spacing on growth and seed yield of coriander (*Coriandrum sativum* L.) *International Journal of Chemical Studies* . 6(4): 2768-2772.

Emamghoreishi, M., Khasaki, M., Aazam, M. F. (2005). *C. sativum*. Evaluation of its anxiolytic effect in the elevated plus-maze. *J Ethnopharmacol*. 96:365–70.

El- Mekawey, M.A.M., Ali, M.A.M., Awad, A.E. and Hassan, H.M.S. (2010) . Effect of fertilization and growth regulators on *coriandrum sativum* L. plants productivity under north sinai conditions. *J. Agric. Res.* 36 :314 -339.

Fikadu, L. W., Diriba, S. G. and Muluaalem, A. M. (2019). The need of integrated nutrient management for coriander (*Coriandrum sativum* L.) production. *International Journal of Food & Nutrition* .4 (1): 1 – 13.

Grangeiro, L. C., Oliveira, F. C. L., Negreiros, M. Z., Marrocos, S. de T. P., Lucena, R. R. M. de and Oliveira, R. A. de . (2011). Nutrients growth and accumulation in coriander and rocket. [Portuguese]. *Revista Brasileira de Ciencias Agrarias*. 6 (1): 11-16.

Gujar, S. M., Warade, A. D., Mohariya, A. and Paithankar, D. H. (2005). Effect of dates of sowing and nitrogen levels on growth, seed yield and quality of coriander. *Crop Research* (Hisar). 29 (2): 288-291.

Godara, A. S., Gupta, U. S., Lal, G. and Singh, R.. (2014). Influence of organic and inorganic source of fertilizers on growth, yield and economics of coriander (*Coriandrum sativum* L.) *International J. Seed Spices* .4(2): 77-80.

Ibrahim, S. M., El-Labban, H. M., Mohamed. F. I. and Naga, N. M. (2006). Effect of Organic manures and chemical Fertilizers on *Foeniculum vulgare*, Mill and *Carum Carvi*, L. Bull. Pharm. Sci., Assiut University. 29 (1):187-201.

Jan, I., Sajid, M., Shah, A. H., A. Rab, Khan, N. H., Wahid, I. F., Rahman, A., Alam. R and H. Alam. (2011). Response of seed yield of coriander to phosphorus and row spacing. Sarhad. J. Agric. 27(4): 549-552.

Javiya, P.P., Solanki, J.N., Kaneria, S.C. and Rupareliya, V.V. (2017). Response of Coriander (*Coriandrum sativum* L.) to Nitrogen and Phosphorus in South Saurashtra Condition. Int. J. Pure App. Biosci. 5(4): 860-866.

Jhankar, P., Panda, C. and Sethi, D. (2017). Effect of INM practices on yield, yield attributes and economics of coriander (*Coriandrum sativum* L.). Int. J. Curr. Microbiol. App. Sci. 6(5): 1306- 1312.

Khalid, K. A. (2012). Effect of NP and foliar spray on growth and chemical compositions of some medicinal Apiaceae plants grow in arid regions in Egypt. Journal of Soil Science and Plant Nutrition. 12 (3): 617-632.

Khan, M., Habib, A., Abdullah, A., Tahir, A., Zahid, K., Asghar, J., Latif, S. and Mushtaq, M. (2019). Seed associated mycoflora of coriander (*Coriandrum sativum* L.) its effect on seed germination and management through seed treatment chemical. The Int. J. Biol. Res. 2: 101-117.

Kumar, A., Singh, R. and Chhillar, R. K. (2008). Influence of omitting irrigation and nitrogen levels on growth, yield and water use efficiency of coriander (*Coriandrum sativum* L.). Acta Agronomica Hungarica. 56 (1): 69-74.

Kumar, K., Singh, G. P., Singh, N., Bhatia, A. K. and Nehra, B. K. (2007). Performance of seed crop of coriander under different levels of row spacing, nitrogen and cycocel. Haryana Journal of Horticultural Sciences. 36 (1/2): 127-128.

Kumar, S., Choudhary, G. R. and Chaudhari, A. C. (2002). Effects of nitrogen and biofertilizers on the yield and quality of coriander (*Coriandrum sativum* L.). Annals of Agricultural Research. 23 (4): 634-637.

Lal, G., Lal, S., Choudhary, M.K., Meena R.S., and Shekhawat, N., (2020). Growth, yield and essential oil of coriander (*Coriandrum sativum* L.) variety ACr-2 as influenced by various nutrient levels and crop geometry. *International Journal of Chemical Studies* ; 8(4): 2749-2752.

Leena K, Sharma A, Lodi S (2012). Potential health benefits of coriander (*Coriandrum sativum*) . *Int. J. Pharm. Res. Dev.* 4(2):10-20.

Mahendra., P., Bisht, S. B., (2011). Anti-anxiety activity of *Coriandrum sativum* assessed using different experimental anxiety models. *Indian Journal of Pharmacology.* 43(5): 574-7.

Manure, G. R., Shivaraj, B., Farooqui, A. A. and Surendra, H. S. (2000). Yield attributes, seed, essential oil yield and oil content of coriander (*Coriandrum sativum* L.) as influenced by the graded levels of nitrogen, sulphur and zinc nutrition in red sandy loam soils. challenges and opportunities in the new century. Contributory papers. *Centennial conference on spices and aromatic plants.* 139-144

Mirchandani T.B, Mirchandani R. (2005) For balanced specialty nutrition in N. Ram (ed.) *The Hindu Survey of the Indian agriculture, Chennai,* 202- 205.

Mohamed M. A-H. and Abdu.M. (2004) .Growth and Oil Production of Fennel (*Foeniculum vulgare* Mill): Effect of Irrigation and Organic Fertilization. *Biological Agriculture and Horticulture.* 22:31-39.

Moslemi, M., Aboutalebi, A., Hasanzade, H. and Farahi, M. H. (2012). Evaluation the effects of different levels of vermicompost on yield and yield components of Coriander (*Coriandrum sativum* L.). *Annals of Biological Research.* 3 (10) : 4852-4853.

Mounika, Y., Sivaram, G., Reddy, P. and Ramaiah, M. (2018). Influence of biofertilizers and micronutrients on seed yield, essential oil and oleoresins of coriander (*Coriandrum sativum* L.) cv. Sadhana. *Bull. Env. Pharmacol. Life Sci.,* (7):36-39.

Nagar, R. K. Meena, B. S. and Dadheech, R. C. (2009). Effect of weed and nutrient management on growth, yield and quality of coriander (*Coriandrum sativum* L.). *Indian Journal of Weed Science*. 41 (3/4): 183-188.

Nagar, R. K., Meena, B. S., and Dadheech, R. C. (2009). Effect of integrated weed and nutrient management on weed density, productivity and economics of coriander (*Coriandrum sativum*). *Indian Journal of Weed Science*. 41 (1/2): 71-75.

Naghera, R. P., Sukhadia, N. M. and Ramani, B. B. (2000). Effect of sowing dates and varying levels of nitrogen and phosphorus on Coriander (*Coriandrum sativum* L.). *Gujarat Agricultural University Research Journal*. 26 (1): 52-54.

Nayak, B. R., Saini, S. S. and Sahu, G. S. (2009). Effect of farm yard manure, nitrogen and plant spacings on yield and attributing character of coriander (*Coriandrum sativum* L.). *Environment and Ecology*. 27 (3A): 1374-1377.

Okut, N. and Ydrn, B. (2005). Effects of different row spacing and nitrogen doses on certain agronomic characteristics of coriander (*Coriandrum sativum* L.). *Pakistan Journal of Biological Sciences*. 8 (6): 901-904.

Oliveira, A. P., de Alves, E. U. , Bruno, R. de L. A., Sader, R. and Alves, A. U. (2006). Yield and quality of coriander seeds in function of nitrogen levels. [Portuguese]. *Revista Brasileira de Sementes*. 28 (1): 193-198.

Pooja., Nagre, P. and Yadav, H. (2017). Influence of different levels of nitrogen and phosphorus on seed yield and economics of coriander (*Coriandrum sativum* L.). *Journal of Pharmacognosy and Phytochemistry*. 6(5): 157-160.

Sadanandan, A. K. and Hamza, S. (2006). Effect of Organic Farming on Soil Quality, Nutrient Uptake, Yield and Quality of Indian Spice. Scientific registration n^o:2445. Symposium n^o :40.

Sahib, N.G., Anwar, F., Gilani, A. H., Hamid, A.A., Saari, N., Khalid M. and Alkharfy, K.M. (2013). Coriander (*Coriandrum sativum* L.) A Potential Source of High-Value Components for Functional Foods and Nutraceuticals- Phytotherapy research. (wileyonlinelibrary.com) DOI: 10.1002/ptr.4897.

Sharafzadeh, S. and Ordookhani, K. (2011). Organic and Bio Fertilizers as a Good Substitute for Inorganic Fertilizers in Medicinal Plants Farming. *Australian Journal of Basic and Applied Sciences*, 5(12): 1330-1333.

Sharma, M. M. and Sharma, R. K. (2012). Handbook of Herbs and Spices (Second Edition), Volume 1.

Singh M .(2011).Effect of vermicompost and chemical fertilizers on growth, yield and quality of coriander (*Coriandrum sativum L.*) in a semi-arid tropical climate. *Journal of Spices and Aromatic Crops*. 20 (1) : 30-33.

Singh, J., Malik, Y. S., Thakral, K. K. and Mehla, C. P. (2000). Effect of sowing time, nitrogen levels and leaf cuttings on green and seed yield of Coriander. *Haryana Journal of Horticultural Sciences*. 29 (3/4): 225-228.

Singh, S. and Jat, N. L. (2002). Effect of phosphorus and zinc fertilization on growth and yield of coriander (*Coriandrum sativum L.*). *Annals of Agricultural Research*. 23 (4): 734-736

Tripathi, M. L. (2006). Response of coriander varieties to fertility levels. *Research on Crops*. 7 (3): 885-886.

Tripathi, M. L. (2006). Effect of potassium and sulphur levels on yield and uptake of N, P and K by coriander. *Crop Research (Hisar)*. 32 (3): 370-371.

Tripathi, M., Singh, H. and Chouhan, S. (2013). Response of coriander (*Coriandrum Sativum L.*) to integrated nutrient management. *A Journal of Multidisciplinary Advance Research*. 2(2): 43- 46.

Van Kessel, J.S., and Reeves J.B. (2002). Nitrogen mineralization potential of dairy manures and its relationship to composition. *Biol. Fertil. Soils*, 36 pp. 118-123.