

## Determining the Effect of Integrated Nutrient Management and Panchagavya spray on growth and yield of Rice in India

### Abstract

A field study was conducted during *Kharif* 2021 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P), India. The soil texture of the experimental plot was sandy loam, with a pH of 7.8, low organic carbon (0.35 percent), available N (243 kg/ha), available P (20.10 kg/ha), and available K (105 kg/ha). The Study was laid out in Randomized Block Design, with nine treatments consists of viz., 1 : (RDF 50% + FYM 5t/ha) +P<sub>1</sub> -3%, 2 : (RDF 50% + FYM 5t/ha) +P<sub>2</sub> - 6%, 3 : (RDF 50% + FYM 5t/ha) +P<sub>3</sub> - 9%, 4: (RDF 75% +FYM 5t/ha) + P<sub>1</sub> - 3%, 5 : (RDF 75% + FYM 5t/ha) + P<sub>2</sub> -6%, 6 : (RDF75% + FYM 5t/ha) + P<sub>3</sub>- 9%, 7 : (RDF 100% +FYM 5t/ha) + P<sub>1</sub> -3%, 8 : (RDF 100% + FYM 5t/ha) +P<sub>2</sub>- 6%, 9 : (RDF 100% + FYM 5t/ha) + P<sub>3</sub> - 9% and were replicated thrice. Results obtained that there was significant increase in yield parameters viz., Number of panicles /hill (12.53), Number of grains/hill (118.13), Panicle length (32.13 cm), Test weight (24.27 g), Grain yield (6.28 t/ha), Straw yield (7.91t/ha) and Harvest index (44.25%) were recorded with the application of (RDF 100% + FYM 5t/ha) + P<sub>2</sub>-6%. The maximum Gross return (169650.00 INR/ha), Net return (115561.00 INR/ha) and B:C ratio (2.14) is recorded in treatment with (RDF 100% + FYM 5t/ha) +P<sub>2</sub>-6%. Therefore, it is concluded that the application of (RDF 100% + FYM 5t/ha) +P<sub>2</sub>-6% was more productive and economically feasible.

**Keywords:** Rice, integrated nutrient management, FYM, panchagavya, Economics

## INTRODUCTION

Rice, (*Oryza sativa*), edible starchy cereal grain and the grass plant (family Poaceae) by which it is produced. Roughly one-half of the world population, including virtually all of East and Southeast Asia, is wholly dependent upon rice as a staple food; 95 percent of the world's rice crop is eaten by humans.

As a cereal grain, domesticated rice is the most widely consumed staple food for over half of the world's human population especially in Asia and Africa. It is the agricultural commodity with the third-highest worldwide production, after sugarcane and maize. Since sizable portions of sugarcane and maize crops are used for purposes other than human consumption, rice is the most important food crop with regard to human nutrition and caloric intake, providing more than one-fifth of the calories consumed worldwide by humans. There are many varieties of rice and culinary preferences tend to vary regionally.

Total production of Rice during 2019-2020 is estimated at record 117.94 million tonnes. It is higher by 8.17 million tonnes than the five years average production of 109.77 million tonnes. In the 2020/2021 crop year, China produced over 148 million metric tons of milled rice, a higher volume than any other country, India came in second place with 122 metric tonnes of milled rice in that year. The highest productivity is 6710 kg/ha in china followed by Vietnam. In India West Bengal is the largest producer of rice. The area of Rice in India in the year 2022/2021 is about 45,400 thousand hectare and in the year 2021/2022 the area increased to about 47,000 thousand hectares. In India West Bengal is the largest producer of rice. Two crops of rice are raised in a year in this state. Rice is regarded as the master crop of coastal India and in few regions of eastern India.

“Integrated Nutrient Management aims to use nutrients in a more rational way (yield-targeted, site- and soil specific); understanding the interrelation of different nutrients; use combinations of mineral and organic fertilizers; provide nutrients on a cropping-system/rotation basis; and use on-farm and off-farm waste through recycling. Nutrient cycling is an important component of Conservation Agriculture, in which minimum soil disturbance, intercropping, crop rotations and a permanent soil cover minimize the need for chemical fertilizers. Healthy crops are also less susceptible to pests, thus contributing to crop protection (IPM). A better application of nutrients will reduce runoff, and by this benefits the overall ecosystem, including marine areas. INM aims to improve soil health and sustain high level of productivity and production” (**Prasad et al. 1995**)

“It’s need in modern agriculture has arisen due to: (i) high price of chemical fertilizers; (ii) imbalance in the ratio of NPK consumption; (iii) imbalance between consumption and domestic production; (iv) deterioration of soil health; (v) consumption of non-renewable energy sources by inorganic fertilizers; (vi) pollution hazards of chemical fertilizers; (vii) loss of chemical productivity; (viii) deterioration in soil physical properties; (ix) deterioration in biological activity; (x) additive effect of organic and mineral fertilizers; (xi) organic materials as a source of secondary micronutrients; (xii) interaction benefit crops; and reduction in crop productivity. INM aims to improve soil health and sustain high level of productivity and production. Organic supply of nutrients at the peak period of absorption also provide micro nutrients and modify soil-physical behavior as well as increase the efficiency of applied nutrients” (**Pandey et al. 2007**). ). **“Application of FYM (50% or 100%) in kharif rice induces higher residual contribution of nutrients like N, P, K and Application of FYM (50% or 100%) in kharif rice induces higher residual contribution of nutrients like N, P, K and S to the succeeding rabi lentil crop” (Pankaj et al.2013)**

“Production efficiencies are gained through nutrient management practices that promote combined use of mineral, organic and biological resources in a reasoned way to balance efficient use of limited/finite resources and ensure ecosystem sustainability against nutrient mining and degradation of soil and water resources. **Farmyard manure (FYM) is being used as prominent source of organic manure in crops as it supplies all essential plant nutrients and increases activities of microbes in soil” (Sutaliya and Singh, 2005).** Limited availability of FYM is however a major disadvantage in its uses as source of nutrients.

Panchagavya is an organic product having the potential to play the role of promoting growth and providing immunity in plant system. Panchagavya consists of nine products viz. cow dung, cow urine, milk, curd, jaggery, ghee, banana, Tender coconut and water. When suitably mixed and used, these have miraculous effects. It has micro nutrients like nitrogen, phosphorus and potassium which are the major nutrients required for the growth of plants. Panchagavya also contains many vitamins, amino acids. It also contain Gibberellins and auxins which regulate the growth of plants. Panchagavya also contains microorganisms like Psuedomonas, Azotobacter, phospho bacteria which are considered to be beneficial for rice crop. Panchagavya has the capacity to restore the yield efficiency level. It can be easily prepared at the field level without any special techniques. Panchagavya, an organic source of nutrition, is an indigenous material which is used widely for agricultural and horticultural crops. Panchagavya had positive influence on beneficial microorganisms present in the soil and influence the crop growth and yield. Panchagavya is now gaining attention as an efficient organic growth promoter.

## MATERIALS AND METHODS

This experiment was carried out during *Kharif* 2021 at Crop Research Farm, Department of Agronomy, NAI, SHUATS, Prayagraj, (U.P.) which is located at 25.28°N latitude, 81.54°E longitude and 98 m altitude above the mean sea level. This area is situated on the right side of the river Yamuna by the side of Prayagraj, Rewa road, about 5 km away from Prayagraj city. The soil of experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.8), low in organic carbon (0.35%), available N (243 kg/ha), available P (20.10 kg/ha) and available K (105 kg/ha). The experiment was laid out in Randomized Block Design, with nine treatments consists of viz., 1 : (RDF 50% + FYM 5t/ha) +P<sub>1</sub> -3%, 2 : (RDF 50% + FYM 5t/ha) +P<sub>2</sub> - 6%, 3 : (RDF 50% + FYM 5t/ha) +P<sub>3</sub> - 9%, 4 : (RDF 75% +FYM 5t/ha) + P<sub>1</sub> - 3%, 5 : (RDF 75% + FYM 5t/ha) + P<sub>2</sub> - 6%, 6 : (RDF75% + FYM 5t/ha) + P<sub>3</sub>- 9%, 7 : (RDF 100% +FYM 5t/ha) + P<sub>1</sub> -3%, 8 : (RDF 100% + FYM 5t/ha) +P<sub>2</sub>-6%, 9 : (RDF 100% + FYM 5t/ha) + P<sub>3</sub> - 9% and were replicated thrice. Rice variety Shiats dhan 3 was taken as test crop. The crop matures in about 130-138 days. The yield potential of the crop is 6.5 t/ha. The main field was prepared by ploughing with the help of cultivator. The main field was puddled by tractor drawn puddler and later it was leveled by planker. The 21 days old seedlings were transplanted to the main field. The transplanting was done at a spacing of 20 × 10 cm All the nutrients such as (N, P<sub>2</sub>O<sub>5</sub>, K<sub>2</sub>O) were applied to the soil in form of urea, DAP and MOP respectively. The 100 % recommended dose of fertilizer is 120:60:60. FYM is incorporated into the soil after puddling and before transplanting of seedling at a recommended rate of 5 t/ha. For preparing panchagavya, a wide mouthed plastic jar was used, as metal containers are not recommended to be used. First the fresh cow dung and cow's ghee were put into container and mixed thoroughly twice daily for 3 days. On the fourth day rest of the ingredients were added and stirred twice daily for 15 days. The panchagavya stock solution was ready after the 20th day. It was kept in the shade and covered with a plastic mosquito net to prevent houseflies from laying eggs and the formation of maggots (worms) in the solution. Whenever required suitable quantity of water was added to keep the slurry in a liquid state. Three, six and nine per cent solutions were used for spray at 15, 30 and 45 DAT as per

the treatment. The 15, 30 and 45 litres of panchagavya in every 500 litres of water were applied to rice at 3,6 and 9 % solutions respectively for 1 ha of rice. After dilution, the panchagavya solution was filtered and sprayed through knapsack sprayer. The growth parameters were recorded at periodical intervals of 20,40,60,80 DAT and at harvest stage from the randomly selected five plants in each treatment. Statistically analysis was done for all the parameters by using one way anova and mean compared at 5% probability level of significant results.

## RESULTS AND DISCUSSION

### Effect of Integrated Nutrient Management and Panchagavya spray on yield parameters of Rice

Higher grain yield and availability of nutrients to the crop efficiently is achieved with the incorporation of FYM to the soil. Effect of Integrated Nutrient Management and Panchagavya spray on yield parameters of Rice are presented in Table 1. The maximum number of panicles /hill (12.53), Number of grains/hill (118.13), Panicle length (32.13 cm), Test weight (24.27 g), Grain yield (6.28 t/ha), Straw yield (7.91 t/ha) and Harvest index (44.25 %) were recorded with application of (RDF 100% + FYM 5t/ha) +P<sub>2</sub>- 6% closely followed by the number of panicles /hill (11.80), number of grains/hill (113.07), Panicle length (29.97 cm), Test weight (21.23 gm), Grain yield (5.80 t/ha), Straw yield (7.40 t/ha) and Harvest index (43.92 %) which was recorded in the treatment of (RDF 100% + FYM 5t/ha) +P<sub>2</sub>-3%.The higher grain and straw yield might be attributed because of rapid decomposition of Farm yard manure and release of nutrients throughout the crop growth period combined with proper utilization of nutrients. The lowest yield was obtained in the treatment combination of (RDF 50% + FYM 5t/ha) + P<sub>1</sub>- 3% when compared to the rest of the treatments because of the less application of recommended dose of fertilizer. The higher yield might be due to the better and optimum nutrient availability to the crop from the 100% RDF and Panchagavya spray 6 % compared to other levels. The improvement in nutrient uptake in Rice crop may be due to adequate availability of nutrients in the soil and positive interaction effects of organic and inorganic sources of nutrients. Similar results with the combination of RDF 100% +FYM 5 t/ha were also reported by **S.Maiti et al., (2006)**. **S.K.Prasad et al., (2018)** stated that application of Panchagavya spray @ 6% improved the grain yield and straw yield.

### Effect of Integrated Nutrient Management and Panchagavya spray on Economics of Rice

Effect of Integrated nutrient management and Panchagavya spray on yield and economics of Rice are presented in Table 2. Maximum Gross returns (169650.00 INR/ha), Net returns (115561.00 INR/ha) and Benefit cost ratio (2.14) was obtained with the treatment combination of (RDF 100% + FYM 5t/ha) +P<sub>2</sub>-6%. However the treatment combination of (RDF 100% +FYM 5t/ha) + P<sub>1</sub>-3% were statistically at par with (RDF 100% + FYM 5t/ha) +P<sub>2</sub>-6% with the Grain yield of (5.28 t/ha). Maximum gross returns is obtained because of

the high grain yield which in turn because of the incorporation of the FYM and application of 100% recommended dose of the fertilizer. Maximum net returns is obtained because of the high gross returns. Grain yield of Rice was significantly influenced by integrated nutrient management as well as application of inorganic fertilizer. The cost of cultivation increased with the increased application of inorganic fertilizers. The advantages of organics such as FYM and Panchagavya applied to the crop and to the soil have improved the soil health, which are not calculated in terms of money. Application of Panchagavya provides resistance to the crop against pests and diseases instead of use of chemical pesticides and reduces the cost of cultivation which in turn increases the gross returns. The higher yield with sufficient quantity of organics is due to its nature of providing balanced supply of all the essential nutrients, which balances with crop needs, uptake and thus result in significantly higher grain yield over other higher doses of the same organic source. Higher level of RDF having essential nutrients in adequate amount and balanced proportion might have increased photosynthesis and photosynthates translocation from leaf to grain leading to increased production which in turn increases the gross returns, net returns and B:C ratio (Sudhakar *et al.* 2006). Similar results with the combination of INM and panchagavya were also reported by (P K UPADHYAY *et al.* 2018)

**Table 1 Effect of Integrated Nutrient Management and Panchagavya spray on yield parameters of Rice**

<b>Treatment combinations</b>	<b>No.of panicles /hill</b>	<b>No. of grains /hill</b>	<b>Panicle length (cm)</b>	<b>Test weight (g)</b>	<b>Grain yield (t/ha)</b>	<b>Straw yield (t/ha)</b>	<b>Harvest index (%)</b>
1 : (RDF 50% + FYM 5t/ha) + P <sub>1</sub> - 3%	7.53	92.33	25.90	18.23	4.35	6.34	40.70
2 : (RDF 50% + FYM 5t/ha) + P <sub>2</sub> - 6%	9.07	94.13	26.53	18.37	5.13	7.20	41.59
3 : (RDF 50% + FYM 5t/ha) + P <sub>2</sub> - 9%	10.0	102.07	27.77	21.00	4.97	7.10	41.15
4 : (RDF 75% +FYM 5t/ha) + P <sub>1</sub> - 3%	9.47	99.40	27.30	19.03	4.90	6.97	41.27
5 : (RDF 75% + FYM 5t/ha) + P <sub>2</sub> - 6%	11.07	109.33	28.63	20.37	5.60	7.27	43.50
6 : (RDF75% + FYM 5t/ha) + P <sub>3</sub> - 9%	9.63	97.57	26.40	18.77	4.90	6.97	41.29
7 : (RDF 100% +FYM 5t/ha) + P <sub>1</sub> - 3%	11.8	113.07	29.97	21.23	5.80	7.40	43.92
8 : (RDF 100% + FYM 5t/ha) +P <sub>2</sub> -6%	12.53	118.13	32.13	24.27	6.28	7.91	44.25
9 : (RDF 100% + FYM 5t/ha) + P <sub>3</sub> - 9%	9.50	106.73	26.91	19.40	5.20	7.24	41.80
F test	S	S	S	S	S	S	S
SEm (±)	0.42	1.74	0.4	0.48	0.17	0.11	0.85
CD (5%)	1.25	5.21	1.2	1.45	0.51	0.33	2.55

**Table 2 Effect of Integrated Nutrient Management and Panchagavya spray on economics of Rice**

Treatment combinations	Cost of cultivation (INR/ha)	Gross returns (INR/ha)	Net returns (INR/ha)	B:C Ratio
1 : (RDF 50% + FYM 5t/ha) +P <sub>1</sub> -3%	50794	117450	66656	1.31
2 : (RDF 50% + FYM 5t/ha) +P <sub>2</sub> - 6%	50994	138600	87606	1.72
3 : (RDF 50% + FYM 5t/ha) +P <sub>3</sub> - 9%	51219	134100	82881	1.62
4 : (RDF 75% +FYM 5t/ha) + P <sub>1</sub> -3%	52341	133650	81309	1.55
5 : (RDF 75% + FYM 5t/ha) + P <sub>2</sub> -6%	52541	151200	98659	1.88
6 : (RDF75% + FYM 5t/ha) + P <sub>3</sub> - 9%	52766	132300	79534	1.51
7 : (RDF 100% +FYM 5t/ha) + P <sub>1</sub> -3%	53889	156600	102711	1.91
8 : (RDF 100% + FYM 5t/ha) +P <sub>2</sub> - 6%	54089	<b>169650</b>	<b>115561</b>	<b>2.14</b>
9 : (RDF 100% + FYM 5t/ha) + P <sub>3</sub> -9%	54314	140400	86086	1.58



## CONCLUSION

From the above experiment it is concluded that transplanting of Rice with the application of (RDF 100% + FYM 5t/ha) +P<sub>2</sub> - 6% has found to be more productive and remunerative. Therefore, it is recommended for farmers for receiving higher yield and economic benefits of Rice.

## REFERENCES

- Avjit sen, PK Upadhyay, Yashwanth singh, SK Prasad, JP Srivastava, RK Singh (2018) . Effect of Panchagavya and recommended dose of fertilizers on growth, nutrient content and productivity of transplanted rice (*Oryza sativa*) under middle gangetic plain of India. *Indian journal of Agricultural sciences* 88(6): 931-6
- Pandey, N., Verma, A.K., Anurag and Tripathi, R.S. 2007. Integrated nutrient management in transplanted hybrid rice (*Oryza sativa*). *Indian Journal of Agronomy* 52(1) 40-42.
- Pankaj P K, Mandal N, Singh R P and Nirmal D. 2013. Nutrient management in direct seeded rice and its residual effect on uptake, translocation and recovery of nutrients in rainfed lentil. *An Asian Journal of Soil Science* 8(2): 226-30
- P K UPADHYAY, AVIJIT SEN *et al.* 2013. Effect of panchagavya and recommended dose of fertilizers on growth, nutrient content and productivity of transplanted rice (*Oryza sativa*) under middle Gangetic plain of India. *Indian Journal of Agricultural Sciences* 88 (6): 931-6
- Prasad, B., Prasad, J. and Prasad , R. 1995. Nutrient management for sustained rice and wheat production in calcareous soil amended with green manures, organic manure and zinc (ENG). *Fertilizer News* 40(3):39-41.
- S.Maiti *et al.*, (2006). Integrated nutrient management under hybrid rice (*Oryza sativa*) hybrid rice cropping sequence. *Indian journal of Agronomy* 51 (3): 157-159
- Sutaliya, R. and Singh, R.N. 2005. Effect of planting time, fertility level and phosphatesolubilizing bacteria on growth, yield and yield attributes of winter maize (*Zea mays*) under rice (*Oryza sativa*)- maize cropping system. *Indian Journal of Agronomy* 50(3):173-75.
- Sudhakar S K. 2008. 'Characterization and evaluation of jeevamruta formulations on aerobic rice.' M Sc (Agri.) thesis University of Agricultural Sciences, Bengaluru, Karnataka, India.