

# **The Role of Nanozeolite and Biocapsule in improving growth, yield and fruit quality of low chilling Apple (*Malus × domestica* Borkh.) Cv. HRMN-99 in Prayagraj Agro-climatic condition**

## **ABSTRACT**

The present investigation was conducted at the Research farm of Department of Horticulture, SHUATS, Prayagraj, on 30-month-old plant during January 2021 to July 2021 with the objective to find out suitable treatment of Biocapsule and Nanozeolite with combination of NPK in relation to plant growth, yield and quality of apple. The statistical design adopted for the experiment was randomized block design (RBD) with three replications and nine treatment combinations. viz, T<sub>1</sub> (NPK (RDF)), T<sub>2</sub> (Biocapsule 500 ppm), T<sub>3</sub> (Nanozeolite 250 ppm), T<sub>4</sub> (Nanozeolite 250 ppm + Biocapsule 500 ppm), T<sub>5</sub> (Nanozeolite 250 ppm + Biocapsule 250 ppm), T<sub>6</sub> (NPK (RDF) + Biocapsule 500 ppm), T<sub>7</sub> (NPK (RDF) + Nanozeolite 250 ppm), T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm), T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm). The treatments were applied by soil drenching at initiation of experiment, pre-flowering stage and the last one at the fruit setting stage to assess the effect on growth, yield and quality of apple. The results of present investigation revealed that, the treatment T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) outstand in all the aspects like Maximum plant height (2.8 m), Maximum number of Primary and Secondary branches (3), (28) respectively, maximum number of flower (319), maximum no. of fruit (63.6), maximum yield Plant<sup>-1</sup>(7.2 kg) and quality parameters like Maximum TSS (11.56 °Brix) and Maximum acidity (0.71 %) followed by T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm). The lowest observation was recorded in T<sub>3</sub> (Nanozeolite 250 ppm). Biocapsules and nano zeolite with combination of NPK played an important role in fixing the nutrients in soil and protecting the apple plant against pathogen effect. That's why a positive response has to be seen in particular treatment during the research trail.

**Keywords-** biocapsule, nanozeolite, low-chilling apple, subtropical region, HRMN-99

## **INTRODUCTION**

The "King of Temperate Fruit," the Apple (*Malus × domestica* Borkh), originated in Central Asia near Turkey. It is an internationally cultivated deciduous and valuable fruit tree of the rosaceae family. Apples have been farmed for thousands of years in Asia and Europe, and the tree was imported to North America by European colonists. Because of its excellent taste and nutritious content, it is popular all over the world. Apple is one of the most popular fruits grown in temperate climates around the world. Its attractive look, crispy flesh, agreeable flavour, and sweet taste entice customers and bring in premium prices (Ali *et al.*, 2004). Apples are abundant in minerals such calcium, potassium, phosphorus, and magnesium, as well as vitamins C, K, and B-6, folate, protein, thiamin, riboflavin, and pantothenic acid. It also contains a high amount of choline, dietary fibre, omega-3 fatty acids, and biotin. It also

consists of various phytochemicals including flavonoids (e.g., catechins, flavanols, and quercetin) and other phenolic compounds (e.g., epicatechin and procyanidins) found in the skin, core, and pulp of the apple (Hyson 2011). Certain phytochemicals found in apples may also help to reduce the incidence of certain cancers. When eaten raw, apples provide health advantages and are a good source of antioxidants. "An apple a day keeps the doctor away," according to a well-known proverb.

HRMN – 99 Apples are cultivated at a lower elevation of 1800 feet above sea level and do not require chilling hours. Average yield (1 quintal/plant from a 7-year-old plant), ready to harvest in early June (after three years of transplanting), and scab-resistant. During 2015-2017, the National Innovation Foundation-India (NIF) planted roughly 10,000 seedlings in 1190 farmers' fields and 25 organizations across 29 states and 5 union territories to investigate the HRMN-99's compatibility and adaptation in diverse agro-climatic conditions across the country. Farmers in Manipur, Madhya Pradesh, Uttar Pradesh, Bihar, Maharashtra, Gujarat, Dadra and Nagar Haveli, Karnataka, Haryana, Rajasthan, Jammu, Kerala, Uttarakhand, Telangana, Himachal Pradesh, and Delhi have reported successful fruiting, resulting in an increase in the demand for saplings on a large scale (National innovation foundation, Database 2017)

Recently, ICAR (Indian Council of Agricultural Research) scientists have developed the technology to pack bio fertilizers in tiny capsules. This eliminates the need for farmers to carry the sacks of biofertilizers. It consists of a carrier medium rich in live microorganism. When applied to seed, soil or living plants, it increases oil and nutrients or makes them biologically available. The formulation involves culturing of PGPR, pelleting the cells at appropriate growth phase, treating with buffering agent, followed by mixing with protecting agent, inert bulking agent and nutrients. The mixture with a PGPR population of 10<sup>10</sup> CFU/g is then packed in the capsule. The whole process is done at a temperature range between 20-30°C. The total weight of the biocapsule is just 1g. The process of making the formulation is simple and does not require any sophisticated equipment and conditions except basic facilities. The capsule formulation contains PGPR in an immobilized/inactive condition and the cells can be activated by dissolving the capsule in water. This suspension can be diluted and the seeds or seedlings or rhizomes are soaked in the suspension for 30 minutes before sowing/ transplanting into the main field. The remaining suspension can be used as soil drench (Source: ICAR-Indian Institute of Spices Research, Kozhikode).

Nanozeolites are silicate minerals that have pores and channels in their crystal structure. It has a high affinity for cations like Na<sup>+</sup>, K<sup>+</sup>, and Ca<sup>2+</sup> due to its unique increased Cation exchange capacity (CEC). In the soil system, zeolites are responsible for the selective retention of NH<sup>4+</sup> and K<sup>+</sup> ions.

## **METHODS AND MATERIALS**

The present research work was carried out in randomized block design (RBD) with three replications and nine treatment combinations. viz, T<sub>1</sub> (NPK (RDF)), T<sub>2</sub> (Biocapsule 500 ppm), T<sub>3</sub> (Nanozeolite 250 ppm), T<sub>4</sub> (Nanozeolite 250 ppm + Biocapsule 500 ppm), T<sub>5</sub> (Nanozeolite 250 ppm + Biocapsule 250 ppm), T<sub>6</sub> (NPK (RDF) + Biocapsule 500 ppm), T<sub>7</sub> (NPK (RDF) + Nanozeolite 250 ppm), T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm), T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) and about 30 months old Apple plant were taken for research purpose.

### **Experimental Site:**

The experiment was conducted at the Research farm, Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of agriculture, technology and sciences, Prayagraj, Uttar Pradesh for a period lasting from January 2021 to July 2021.

### **Geographical location of the experimental site:**

The location of experimental site is situated at an elevation of 98 meters above mean sea level (MSL) at 25.45° North latitude and 81.84° East longitudes.

### **Climate:**

This region falls under the sub-tropical zone. Prevailing in the South-Eastern part of the U.P. and has both the extremities in temperatures, *i.e.*, winter and summer. In winters, the temperature sometimes is as low as 32°F in December to January, and in summer the temperatures reach up to 115°F in the months of May to June. During winter, frosts and during summer, hot scorching winds are pretty common. The average rainfall is around 1013.4 (mm) with maximum concentration during July to September months with occasional showers in winters.

### **Preparation of Biocapsule solution**

Biocapsules were purchased from the ICAR licensed company CADAGU Agritech. Karnataka. Each biocapsules having 1gm wgt. or 1000 ppm concentration of PGPR<sub>s</sub> and the ability of dissolving in 100 ltr of water. So according to the concentration of capsules are first dissolve in 1 ltr of luke warm water before 12 hours of application and then after that 1 ltr of solution was dissolve in 50 ltr. (For 500 ppm) and 25 ltr. (For 250 ppm) of water. After that Apple plants were drenched with 1 ltr of that solution. The biocapsule drenching were applied at different interval during the research.

### **Preparation of Nanozeolite solution**

Nanozeolite were purchased from the Nanoresearch element Haryana. According to their concentration of ppm they are first dissolve in the 5ml ethyl alcohol solution after that it dissolve in the 1 ltr. of water and then drenched in the Apple plant. The Nanozeolite drenching were applied at different interval during the research.

## **RESULT AND DISSCUSSION**

### ***Growth Parameters***

In case of plant height, Table 1, and graph 1 showed that the Treatment T<sub>8</sub> ((NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) have maximum plant height (2.8 m) after 3 years of transplanting followed by T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) (2.76 m) and minimum plant height (2.47 m) was observed in treatment T<sub>3</sub> (Nanozeolite 250 ppm) after 3 years of transplanting. Increment in plant height might be due to Biocapsules and nanozeolite with the combination of NPK. Biocapsules increases the nutrient fixation in the root and nanozeolite played an important role to make availability of

nutrient in the soil. These results are in support with **Bandana and Chandel (2017)** and **Padhan et. al. (2019)**.

The maximum primary branches (3.66) were observed in treatment T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) followed by treatment T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) which was (3) and the lowest primary branches (2) was recorded in treatment T<sub>3</sub> after 3 years of transplanting. Increased number of primary branches, might be due to the application of Biocapsule and nanozeolite with the combination of NPK. Biocapsule has an antipathogen effect and nanozeolite reduces the disease incidence to the Apple plant. These findings are in accordance with **Treder (2007)** and **Padhan et. al. (2019)**.

The number of secondary branches was noticed maximum (30.33) after 3 years of transplanting with treatment T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm), followed by treatment T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) (28), while minimum number of secondary branches observed with treatment T<sub>3</sub> (Nanozeolite 250 ppm) (14.33) after 3 years of transplanting. Increased number of secondary branches, might be due to the application of Biocapsule and nanozeolite with the combination of NPK. Biocapsule has an antipathogen effect and nanozeolite reduces the disease incidence to the Apple plant. Similar findings were reported by **Treder (2007)** and **Padhan et. al. (2019)**.

In terms of stem diameter, maximum stem diameter was witnessed in T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) (10.66 cm), followed by T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) (10.2 cm) and minimum stem diameter was witnessed in treatment T<sub>3</sub> (Nanozeolite 250 ppm) (6.86 cm) after 3 years of transplanting. Increment in stem diameter, might be due to Biocapsules and nanozeolite with the combination of NPK. Biocapsules increases the nutrient fixation in the root and nanozeolite played an important role to make availability of nutrient in the soil. Similar results were achieved by **Bandana and Chandel (2017)** and **Padhan et. al. (2019)**.

In case of leaf area, treatment T<sub>8</sub> ((NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) have the maximum leaf area which was (40.56 cm<sup>2</sup>) followed by T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) (38.26 cm<sup>2</sup>) and lowest was reported in T<sub>3</sub> (Nanozeolite 250 ppm) (35.4 cm<sup>2</sup>) after 3 years of transplanting. Increment in leaf area, might be due to Biocapsules and nanozeolite with the combination of NPK. Biocapsules increases the nutrient fixation in the root and nanozeolite played an important role to make availability of nutrient in the soil. These findings are in accordance with **Treder (2007)** and **Padhan et. al. (2019)**.

### **Flowering And Fruiting Parameter**

Table 1. and graph 1 Demonstrated that the longest time period from bud burst to first flowering was observed in treatment T<sub>3</sub> (Nanozeolite 250 ppm) was (19.33 days), followed by T<sub>2</sub> (Biocapsule 500 ppm) (18 days) and least time period was observed in treatment T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) which was (12.33 days) after 3 years of transplanting. Biocapsules and nanozeolite with combination of NPK enhanced the reproductive growth that's why this type of result has to be seen. These results are in support with **Bandana and Chandel (2017)** and **Padhan et. al. (2019)**.

The longest time period from bud burst to first fruiting was observed in treatment T<sub>3</sub> (Nanozeolite 250 ppm) was (34 days), followed by T<sub>2</sub> (Biocapsule 500 ppm) (33.66 days) and least time period was observed in treatment T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) which was (27.66 days) after 3 years of transplanting. Biocapsules and nanozeolite with combination of NPK enhanced the reproductive growth that's why this type of result has to be seen. These results are in support with **Bandana and Chandel (2017)** and **Padhan et. al. (2019)**.

In terms of number flowers per plant, Table 2. and graph 2 showed that the maximum number of flowers was witnessed in Treatment T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) which was (319.66), followed by T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) (303.33) and lowest was recorded in T<sub>3</sub> (Nanozeolite 250 ppm) which was (229.33) after 3 years of transplanting. Biocapsules and nanozeolite with combination of NPK enhanced the reproductive growth that's why this type of result has to be seen. These findings are in accordance with **Sharma et. al. (2005)** and **Padhan et. al. (2019)**.

In terms of numbers of fruit per plant, maximum no. of fruits per tree was observed in Treatment T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) (63.66), followed by T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) (59.66) and minimum no of fruits per plant is observed in treatment T<sub>3</sub> (Nanozeolite 250 ppm) which was (31.66) after 3 years of transplanting. These results are in support with **Prilak et. al. (2007)** and **Fediala et. al. (2018)**.

Coming to the fruit weight, the maximum fruit weight was witnessed in T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) (113.33 g), followed by T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) (109.11 g) and minimum fruit weight was witnessed in treatment T<sub>3</sub> (Nanozeolite 250 ppm) which was (73.66 g) after 3 years of transplanting. Biocapsules and nanozeolite with combination of NPK enhanced the reproductive growth that's why this type of result has to be seen. Similar findings were reported by **Verma et. al. (2009)** and **Hany et. al. (2020)**

In terms of polar diameter, the highest polar diameter was noticed in treatment T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) (7.3 cm), followed by T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) (7.03 cm) and lowest polar diameter was noticed in T<sub>3</sub> (Nanozeolite 250 ppm) which was (5 cm) after 3 years of transplanting. Biocapsules and nanozeolite with combination of NPK enhanced the reproductive growth that's why this type of result has to be seen. These findings are in accordance with **Prilak et. al. (2007)** and **Fediala et. al. (2018)**.

In terms of radial diameter, the highest radial diameter was noticed in treatment T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) (6.73 cm), followed by T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) (6.53 cm) and lowest radial diameter was noticed in T<sub>3</sub> (Nanozeolite 250 ppm) which was (4.4 cm) after 3 years of transplanting. Biocapsules and nanozeolite with combination of NPK enhanced the reproductive growth that's why this type of result has to be seen. These results are in support with **Verma et. al. (2009)** and **Fediala et. al. (2018)**.

The maximum fruit yield (kg/plant) was recorded in treatment T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) (7.22 kg), followed by T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) (6.51 kg) and the minimum fruit yield (kg/plant) was recorded in T<sub>3</sub> (Nanozeolite 250 ppm) which was (2.33 kg) after 3 years of transplanting. Similar findings were reported by **Verma et. al. (2009)** and **Prilak et. al. (2007)**.

In terms the fruit yield (t/ha), maximum fruit yield was witnessed in treatment T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) (16.05 t.), followed by T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) (14.48 t.) and the minimum fruit yield was witnessed in treatment T<sub>3</sub> (Nanozeolite 250 ppm) which was (5.19 t.) after 3 years of transplanting. These findings are in accordance with **Prilak et. al. (2007)** and **Verma et. al. (2009)**.

### ***Fruit Quality Parameters***


Table 2. and graph 2 Showed that the maximum Total soluble solid was noticed in treatment T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) (11.56<sup>o</sup>), followed by T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) (10.63<sup>o</sup>) and the minimum T.S.S was noticed in T<sub>3</sub> (Nanozeolite 250 ppm) which was (8.53<sup>o</sup>). These results are in support with **Singh *et. al.* (2011) and Emadpour *et. al.* (2015).**

In case of acidity, the highest acidity was witnessed in the treatment T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) (0.71%), followed by T<sub>9</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) (0.63%) and the lowest acidity was witnessed in T<sub>3</sub> (Nanozeolite 250 ppm) which was (0.49%). Similar findings were reported by **Drake *et. al.* (2002) and Roussos and Gasparatos (2009).**

Coming to the T.S.S: Acid ratio, the highest ratio was found in T<sub>3</sub> (Nanozeolite 250 ppm) (17.30), followed by T<sub>2</sub> (Biocapsule 500 ppm) (17.16) and lowest T.S.S: Acid ratio was found in T<sub>6</sub> (NPK (RDF) + Biocapsule 500 ppm) which was (15.01). These findings are in accordance with **Drake *et. al.* (2002) and Singh *et. al.* (2011).**

**Table.1 Impact of biocapsule and nanozeolite on growth and development of apple.**


<b>Treatment notations</b>	<b>Treatments</b>	<b>Plant height (m)</b>	<b>No. of primary branches</b>	<b>No. of secondary branches</b>	<b>Stem diameter (cm)</b>	<b>Leaf area (cm<sup>2</sup>)</b>	<b>Bud burst to first flowering (days)</b>	<b>bud burst to first fruiting (days)</b>
<b>T<sub>1</sub></b>	NPK (RDF)	2.61	2.66	23.66	8.86	36.73	15.66	31
<b>T<sub>2</sub></b>	Biocapsule 500 ppm	2.47	2	18	7.53	35.73	18	33.66
<b>T<sub>3</sub></b>	Nanozeolite 250 ppm	2.44	1.66	14.33	6.86	35.4	19.33	34
<b>T<sub>4</sub></b>	Nanozeolite 250 ppm + Biocapsule 500 ppm	2.56	2.33	22.33	8.46	36.33	16.33	32.33
<b>T<sub>5</sub></b>	Nanozeolite 250 ppm + Biocapsule 250 ppm	2.52	2.33	20.33	8.1	35.9	17.66	33
<b>T<sub>6</sub></b>	NPK (RDF) + Biocapsule 500 ppm	2.69	3	26	9.8	37.76	14.33	30.33
<b>T<sub>7</sub></b>	NPK (RDF) + Nanozeolite 250 ppm	2.66	3	24.66	9.46	37.06	15	30.66
<b>T<sub>8</sub></b>	NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm	2.8	3.66	30.33	10.66	40.56	12.33	27.66
<b>T<sub>9</sub></b>	NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm	2.74	3	28	10.2	38.26	13.66	29.33
<b>F-test</b>		S	NS	S	S	S	NS	S
<b>S. Ed</b>		0.02	0.55	2.90	0.19	0.40	2.51	1.76
<b>CD at 5%</b>		0.05	1.18	6.15	0.41	0.85	5.34	3.73



**Graph 1. Impact of biocapsule and nanozeolite on growth and development of apple.**

**Table.2 Impact of biocapsule and nanozeolite on yield and quality of apple.**

<b>Treatment notations</b>	<b>Treatments</b>	<b>No of flowers</b>	<b>No of fruits</b>	<b>Fruit weight (g)</b>	<b>polar diameter (cm)</b>	<b>Radial diameter (cm)</b>	<b>Fruit (kg/plant)</b>	<b>Fruit (T/Ha)</b>	<b>T.S.S (°BRIX)</b>	<b>Acidity (%)</b>	<b>TSS: ACID RATIO</b>
<b>T<sub>1</sub></b>	NPK (RDF)	268.33	47.66	96.33	6.4	5.7	4.59	10.21	9.3	0.62	15.23
<b>T<sub>2</sub></b>	Biocapsule 500 ppm	232.33	34.66	77.33	5.2	4.7	2.68	5.96	8.93	0.52	17.16
<b>T<sub>3</sub></b>	Nanozeolite 250 ppm	229.33	31.66	73.66	5	4.4	2.33	5.19	8.53	0.49	17.30
<b>T<sub>4</sub></b>	Nanozeolite 250 ppm + Biocapsule 500 ppm	254.66	44.66	91.22	5.96	5.66	4.06	9.04	9.23	0.58	15.79
<b>T<sub>5</sub></b>	Nanozeolite 250 ppm + Biocapsule 250 ppm	241.33	37	81.44	5.53	5.13	3.01	6.68	9	0.54	16.62
<b>T<sub>6</sub></b>	NPK (RDF) + Biocapsule 500 ppm	289.66	53	100.66	6.7	6.4	5.33	11.85	10.06	0.67	15.01
<b>T<sub>7</sub></b>	NPK (RDF) + Nanozeolite 250 ppm	276.33	51	98.44	6.53	6.33	5.01	11.14	9.96	0.64	15.49
<b>T<sub>8</sub></b>	NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm	319.66	63.66	113.33	7.3	6.73	7.22	16.05	11.56	0.71	16.15
<b>T<sub>9</sub></b>	NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm	303.33	59.66	109.11	7.03	6.53	6.51	14.48	10.63	0.69	15.29
<b>F-test</b>		S	S	S	S	S	S	S	S	S	NS
<b>S. Ed</b>		5.29	3.14	1.09	0.06	0.07	0.29	0.65	0.34	0.04	1.71
<b>CD at 5%</b>		11.21	6.66	2.32	0.14	0.15	0.62	1.37	0.72	0.10	3.63



**Graph 2. Impact of biocapsule and nanozeolite on yield and quality of apple.**



**Fig-1. Apple fruit and flowers**

## **CONCLUSION**

On the basis of the present investigation, it is concluded that the various treatments biocapsule and nanozeolite has been applied to enhance the vegetative and reproductive growth of apple cv. HRMN-99, where treatment T<sub>8</sub> (NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 500 ppm) is found superior among others, followed by T<sub>9</sub> NPK (RDF) + Nanozeolite 250 ppm + Biocapsule 250 ppm) and the lowest was T<sub>3</sub> (Nanozeolite 250 ppm) in every aspect of growth, yield and quality parameters.

## **REFERENCES:**

**Ali, & Raza, H. & Khan, Muhammad Azam & Hussain, Maaz. (2004)** Effect of different periods of ambient Storage on chemical composition of apple fruit. *International Journal of Agriculture and Biology*. **06**(3) 568–571

**Bandana and Chandel J.S. (2017)** Effect of growth regulators, PGPR and nitrogen on growth of apple nursery plants under protected conditions. *Journal of Hill Agriculture*, **8** (2):176-180.

**Drake, S.R., Raese, J.T. and Smith T.J. (2002)** Time of nitrogen application and its influence on Golden Delicious apple yield and fruit quality. *Journal of Plant Nutrition* 25, 143-153.

**Emadpour, M., Ghareyazie, B., Kalaj, Y. R., Entesari, M., & Bouzari, N. (2015)** Effect of the potassium permanganate coated zeolite nanoparticles on the quality characteristic and shelf life of peach and nectarine. *Int J Agric Technol*, 11, 1263-1273.

**Fediala Abd El-Gleel Mosa W., Sas Paszt L., Fraç M., Trzciński P., Treder W., Klamkowski K. (2018)** The role of biofertilizers in improving vegetative growth, yield and fruit quality of apple. *Hort. Sci. (Prague)*, 45: 173-180.

**Hany, M. Rahman, A. and Darwesh R. (2020)** Effect of Integrated Fertilization with Inorganic, Organic Fertilizers in Presence of Enterobacter ludwigii Local Strain on Growth, Yield and Fruit Quality of Anna Apple Trees. *Env. Biodiv. Soil Security* (4) 345-360

**Hyson D. A. (2011)** A comprehensive review of apples and apple components and their relationship to human health. *Advances in nutrition (Bethesda, Md.)*, 2(5), 408–420.

ICAR-Indian Institute of Spices Research, Kozhikode.  
<https://icar.org.in/content/novel-method-storing-and-pgprmicrobes-through-biocapsules>

National innovation foundation. 9<sup>th</sup> national biennial grassroots innovation and outstanding traditional knowledge award 2017.  
[https://nif.org.in/upload/innovation/9th/3936h\\_ariman.pdf](https://nif.org.in/upload/innovation/9th/3936h_ariman.pdf)

**Padhan A., Mishra S. and Bahadur V. (2019)** Effect of growing media on growth, development and establishment of low chilling variety of apple “HRMN-99” under Prayagraj agro climatic conditions. *Journal of Pharmacognosy and Phytochemistry*. 8(3): 1227-1230

**Pirlak L., Turan M., Sahin F. & Esitken A. (2007)** Floral and Foliar Application of Plant Growth Promoting Rhizobacteria (PGPR) to Apples Increases Yield, Growth, and Nutrient Element Contents of Leaves, *Journal of Sustainable Agriculture*, 30:4, 145-155

**Roussos, P.A. and Gasparatos, D. (2009)** Apple tree growth and overall fruit quality under organic and conventional orchard management. *Scientia Horticulturae* 123, 247-252.

**Sharma, S.D., Sharma, N., Sharma, C.L., Sood, R. and Singh, R.P. (2005)** Studies on correlations between endomycorrhizal and Azotobacter population with growth, yield and soil nutrient status of apple orchards in Himanchal Pradesh. *Acta Horticulture* 696, 283-287.

**Singh S.R., Zargar M.Y., Najar G.R., Peer F.A. and Ishaq M.I. (2011)** Integrated Use of Organic and Inorganic Fertilizers with Bio-inoculants on Yield, Soil Fertility and Quality of Apple. *Journal of the Indian Society of Soil Science*, 59 (4) 362-367.

**Treder, W. (2007)** Influence of fertigation with nitrogen and a complete fertilizer on growth and yielding of ‘Gala’ apple trees. *Journal of Fruit and Ornamental Plant Research* 5, 143-154.

**Verma, M. L., Singh, C., & Bhardwaj, S. P. (2009)** Effects of biofertilizers on soil moisture, nutrient status and fruit productivity under organic cultivation of apple in Himachal Pradesh. *Indian Journal of Soil Conservation*, 37(3), 201-205.