

Original Research Article

Effect of Organic Fertilizers on Growth, Yield and quality of Cauliflower (*Brassica oleracea* var. *Botrytis*.)

Abstract

The present experiment was carried out during 2021-22 with an aim to study the different treatments of organic fertilizers on growth, yield and quality of Cauliflower and to work out with economics, in Central Horticulture Research Farm of Department of Horticulture, SHUATS,. The experiment was conducted in Randomized Block Design with 10 treatment_replicated three times. The treatments were T₁ FYM 100% (100:40:100), T₂ Vermicompost (75%) + Poultry manure (25%), T₃ Vermicompost (75%) + Goat manure (25%), T₄ Vermicompost (50%) + Poultry manure (50%), T₅ Vermicompost (50%) + Goat manure (50%), T₆ Vermicompost (25%) + Poultry manure (75%), T₇ Vermicompost (25%) + Goat manure (75%), T₈ {Vermicompost + Poultry manure} (75%) + Goat manure (25%), T₉ {Vermicompost + Poultry manure} (50%) + Goat manure (50%), T₁₀ {Vermicompost + Poultry manure} (25%) + Goat manure (75%). From the present investigation it was concluded that treatment T₅ Vermicompost (50%) + Goat manure (50%) was found best in the terms of Plant height (cm), Leaf Length (cm), Number of leaves per plant, Head size (cm), Head weight (g), Net weight of head per plot, Yield per plot (kg), TSS, Acidity (%), B:C Ratio.

Keywords:- Vermicompost, Poultry manure, Goat Manure, Growth, Yield and Economics.

Introduction

Vegetables play an important role in human nutrition. Regular intake of recommended amount of vegetables 300 g per day per capita leads to sound health. Recently, interest in vegetable production has been increased tremendously as a result of greater appreciation of the food value of vegetables and the place of vegetables in the national requirement. Vegetables are an indispensable group of food, providing vitamins, minerals, protein, carbohydrates and fibers in the diet besides having medicinal value and thus provide nutritional security through these nutrients.

India is the second largest producer of vegetables (next to China) and our share in world vegetable production is 10.7 percent. In India, area under vegetable crop cultivation is 10,100 thousand ha with a production of 185883 thousand MT and productivity of 18.40 MT/ha. Cauliflower is grown over an area of 469 thousand ha with production of about 9103 thousand MT per annum (Anonymous, 2019).

Among the vegetables, cauliflower is one of the most important winter season vegetable among the cole crops which belong to the genus *Brassica* of the family Cruciferae with chromosome number $2n =$

18. The word "cauliflower" derives from the Latin words *caulis* (cabbage) and *floris* (flower). The delicacy cauliflower curd is a modification of inflorescence. It is grown throughout the country for its tender curds which are used as vegetable, soup and for pickling (Choudhary, 1996). The crop is reported to be a native of Southern European the Mediterranean region and was introduced in India in 1822 from England (Chatterjee, 1986).

Cauliflower is a cold weather hardy crop and thrives best in cool and moist climate. This had originated from Cyprus and the first crop of cauliflower had been introduced by India in sixth century A.D. There is a great demand for this vegetable on account of its delicious taste. Due to abortive floral parts which are fleshy and closely crowded, these are used for culinary purpose either alone or mixed with potato. Pickle can also be prepared from the firm curd. Cauliflower has high protein and peculiar regarding in stability of Vitamin C after cooking (Singh, 1997). Raw cauliflower provides good amount of nutrients and vitamins. Hundred grams of cauliflower contains vitamin-C (46.4 mg), vitamin-K 16 µg, folate (57.0µg), vitamin B6 (0.2 mg), fiber (2.5 g), Potassium (303.0 mg), Manganese (0.2 mg) and Vitamin B5 (0.7 mg). As weather is a

limiting factor in the production cauliflower, the plant grows best in cool day time temperature 70-85°F (21-29°C), with plentiful sun shine and moist soil conditions high inorganic matter and sandy soils.

Material and Method

The area of Prayagraj district comes under subtropical belt in the south east of Uttar Pradesh, which experience extremely hot summer and fairly cold winter. The maximum temperature of the location reaches up to 46°C-48° C and seldom falls as low as 4°C- 5°C. The relative humidity ranges between 20 to 94 %. The average rainfall in this area is around 1013.4 mm annually. However, occasional precipitation is also not uncommon during winter months.

Result and Discussion

Plant height at 30 DAS

Data reveals that there was a significant effect of various treatments on plant height at 30 days after sowing (DAS). Among the different levels of organic fertilizer, T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum plant height 28.79 cm, followed by T₁₀ (Vermicompost + Poultry manure) (25%) + Goat manure (75%) with 26.75 cm, whereas

minimum Plant height 20.87 cm was recorded in Control.

Plant height at 60 DAS

Data reveals that there was a significant effect of various treatments on plant height at 60 days after sowing (DAS). Among the different levels of organic fertilizer, T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum plant height 42.35 cm, followed by T₁₀ (Vermicompost + Poultry manure) (25%) + Goat manure (75%) with 40.12cm, whereas minimum Plant height 36.95 cm was recorded in control.

Plant height at 90 DAS

Data reveals that there was a significant effect of various treatments on plant height at 90 days after sowing (DAS). Among the different levels of organic fertilizer, T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum plant height 48.72 cm, followed by T₁₀ (Vermicompost + Poultry manure) (25%) + Goat manure (75%) with 46.32 cm, whereas minimum Plant height 42.89 cm was recorded in Control.

Leaf Length 30 DAS

Data reveals that there was a significant effect of various treatments on leaf length at 30 days after sowing (DAS).

Among the different levels of organic fertilizer, T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum leaf length 7.89 cm, followed by T₁₀ (Vermicompost + Poultry manure} (25%) + Goat manure (75%) with 7.51 cm, whereas minimum leaf length 5.64 cm was recorded in Control.

Leaf Length at 60 DAS

Finding reveals that there was a significant effect of various treatments on leaf length at 60 days after sowing (DAS). Among the different levels of organic fertilizer, T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum leaf length 21.95 cm, followed by T₁₀ (Vermicompost + Poultry manure} (25%) + Goat manure (75%)) with 20.50 cm, whereas minimum leaf length 17.89 cm was recorded in Control.

Leaf Length at 90 DAS

Finding reveals that there was a significant effect of various treatments on leaf length at 90 days after sowing (DAS). Among the different levels of organic fertilizer, T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum leaf length 31.88 cm, followed by T₁₀ (Vermicompost + Poultry manure} (25%) + Goat manure (75%)) with 30.35 cm, whereas minimum leaf length 27.89 cm was recorded in Control.

Number of Leaves at 30 DAS

Finding reveals that there was a significant effect of various treatments on number of leaves at 30 days after sowing (DAS). Among the different levels of organic fertilizer, T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum number of leaves 5.95, followed by T₁₀ (Vermicompost + Poultry manure} (25%) + Goat manure (75%)) with 5.13, whereas minimum number of leaves 3.78 was recorded in Control.

Number of Leaves at 60 DAS

Finding reveals that there was a significant effect of various treatments on number of leaves at 60 days after sowing (DAS). Among the different levels of organic fertilizer, T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum number of leaves 12.62, followed by T₁₀ (Vermicompost + Poultry manure} (25%) + Goat manure (75%)) with 11.46, whereas minimum number of leaves 8.18 was recorded in Control.

Number of Leaves at 90 DAS

Data reveals that there was a significant effect of various treatments on number of

leaves at 90 days after sowing (DAS). Among the different levels of organic fertilizer, T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum number of leaves 16.67, followed by T₁₀ (Vermicompost + Poultry manure} (25%) + Goat manure (75%)) with 15.45, whereas minimum number of leaves 12.12 was recorded in Control.

In case of head size, among the different levels of organic fertilizers T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum head size 20.89 cm, followed by T₁₀ (Vermicompost + Poultry manure} (25%) + Goat manure (75%)) with 19.65 cm, whereas minimum size of head 14.25 cm was recorded in Control.

In case of head weight, among the different levels of organic fertilizers T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum head weight of cauliflower 1520.56 g, followed by T₁₀ (Vermicompost + Poultry manure} (25%) + Goat manure (75%)) with 1495.37 g, whereas minimum weight of cauliflower 800.49 g was recorded in Control.

In case of Net head weight among the different levels of organic fertilizers T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum Net weight of head of

cauliflower 1275.95 g, followed by T₁₀ (Vermicompost + Poultry manure} (25%) + Goat manure (75%)) with 1200.64 g, whereas minimum Net weight of head of cauliflower 550.19 g was recorded in Control.

In case of Yield per plot among the different levels of organic fertilizers T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum yield per plot of cauliflower 9.28 kg, followed by T₁₀ (Vermicompost + Poultry manure} (25%) + Goat manure (75%)) with 8.43 kg, whereas minimum yield per plot of cauliflower 4.85 kg was recorded in Control.

Among the different levels of organic fertilizers T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum total soluble solid of cauliflower 6.58 °Brix, followed by T₁₀ (Vermicompost + Poultry manure} (25%) + Goat manure (75%)) with 5.75 °Brix, whereas minimum total soluble solid of cauliflower 4.20°Brix was recorded in Control.

Among the different levels of organic fertilizers T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum acidity of cauliflower 89.82 mg/100g, followed by T₁₀ (Vermicompost + Poultry manure} (25%) + Goat manure (75%)) with 78.85 mg/100g,

whereas minimum acidity of cauliflower 63.45 mg/100 was recorded in Control.

In case of Economics, among the different levels of organic fertilizers T₅ (Vermicompost (50%) + Goat manure (50%)) recorded maximum B:C Ratio 2.06, followed by T₁₀ (Vermicompost + Poultry manure) (25%) + Goat manure (75%) with 1.38, whereas minimum B:C Ratio of cauliflower 1.14 was calculated in Control.

UNDER PEER REVIEW

	CD	0.5	0.62	0.32	0.56	0.6	0.51	0.49	0.58	1.03
	S.Ed	1.6	2.1	0.81	0.23	0.18	0.16	0.21	0.23	0.51

Table 2:Effect of different organic fertilizers on Head size, Head weight, Net head weight, Yield per plot, TSS, Ascorbic acid and B:C Ratio.

Sym bol	Treatment	Head Size (cm)	Head weight (g)	Net head weight (g)	Yield per plot (kg)	TSS (^oB)	Ascorbic acid (mg/100g)	Benefit cost ratio
T ₁	FYM 100% (100:40:100)	14.25	800.49	550.19	4.85	4.2	63.45	1.27
T ₂	Vermicompost (75%) + Poultry manure (25%)	14.85	900.61	750.36	5.46	4.8	68.75	2.50
T ₃	Vermicompost (75%) + Goat manure (25%)	16.75	1280.49	1150.35	7.59	5.01	70.50	2.76
T ₄	Vermicompost (50%) + Poultry manure (50%)	18.65	1375.91	1190.46	7.92	5.65	72.75	2.83
T ₅	Vermicompost (50%) + Goat manure (50%)	20.89	1520.56	1275.95	9.28	6.58	89.82	3.88
T ₆	Vermicompost (25%) + Poultry manure	15.89	1250.64	1000.549	6.24	5.13	75.45	2.84

	(75%)							
T ₇	Vermicompost (25%) + Goat manure (75%)	18.12	1385.12	1160.479	7.65	4.89	77.56	3.37
T ₈	{ Vermicompost + Poultry manure } (75%) + Goat manure (25%)	17.35	1185.60	975.15	6.61	4.67	69.89	3.07
T ₉	{ Vermicompost + Poultry manure } (50%) + Goat manure (50%)	17.95	1320.28	1007.79	7.73	5.35	71.95	3.10
T ₁₀	{ Vermicompost + Poultry manure } (25%) + Goat manure (75%)	19.65	1495.37	1200.64	8.43	5.75	78.85	3.59
	F Ratio	S	S	S	S	S	S	
	CD	0.85	6.59	6.59	6.59	0.67	5.12	
	S.Ed	0.29	2.89	2.89	2.89	0.23	2.31	

Conclusion

It is concluded from the study that the application of (Vermicompost (50%) + Goat manure (50%)) has increased the yield of cauliflower and also helps to retain the fertility of the soil as avoiding the use of chemical fertilizers. The use of organic manure helps to increase the humus content in the soil and also helpful in up taking of NPK from the soil. Optimum level of organic manure use can lead to the sustainable development.

References

- A.O.A.C 1980.** *Official methods of analysis of the association of official analytical chemists.* Edited by Willum Horwitz, 13th eds. Washington, D.C. USA. p. 746.
- Abdel Nabi, H.M.A., El-Gamily, E.I. and El-Amoushi, A.A.R.A. 2020.** Organic fertilization and foliar application with some microelements and biostimulants effects on productivity and quality of cauliflower. *Journal of Plant Production*, 5(11): 41-47.
- Abduli, M.A., Aimiri, L., Madadian, E., Gitipour, S. and Sedighian, S. 2012.** Efficiency of vermicompost on quantitative and qualitative growth of tomato plants. *International Journal of Environmental Research*, 5(6): 467-472.
- Abdulla, A.A. and Sukhraj, K. 2010.** Effect of vermiwash and vermicompost on soil parameters and productivity of okra (*Abelmoschus esculentus*) in Gayana. *African Journal of Agriculture Research*, 4(8)1794-1798.
- Akhter, S., Mostarin, T., Khatun, K., Haq, M.E., Soniya, I.A., Monira, S., Sharmin, I. and Ghosh, A., 2019.** Effect of organic manure and its application timing on the growth and yield of cauliflower (*Brassica oleracea* var. *botrytis*). *Asian Journal of Agricultural and Horticultural Research*, 4 (6)1-14.
- Alam, S.M., Ullah, M.A., Haider, S.I., Nawab, N.N., Aamir, S.S. and Mahmood, I.A. 2019.** Effect of farm yard manure and planting densities on growth, yield and quality of okra under natural farming. *International Journal of Research in Agriculture and Forestry*, 2(6) 21-25.
- Anonymous, 2019.** *Third Advance Estimates*, National Horticulture

- board Gurgaon. Government of India, www.n.h.b.gov.in.20/3/2021.
- Ansari A. 2008.** Effect of vermicompost and vermiwash on the productivity of spinach (*Spinacia oleracea*), onion (*Allium cepa*) and potato (*Solanum tuberosum*). *World Journal of Sciences*, **4(13)**554-557.
- Anuja, S. and Archana, S. 2012.** Effect of organic nutrient on yield and quality of bitter gourd. *International Journal of Agriculture Science*, **8(5)** 205-208.
- Atal 2017.** *Studies on the effect of different organic nutrient sources on bell pepper (Capsicum annum L.) production.* M.Sc. (Agri.) Thesis, Dr. Yashwant Singh Parmar University of Horticulture and Forestry, Nauni, Solan, Himachal Pradesh, India.
- Bano, B., Kale, R.D. and Gajanan, G.N. 1987.** Culturing of earthworm (*Eudrilus eugeniae*) for cast production and assessment of worm cast as bio-fertilizer. *Journal of Soil Biological Ecology*, **6(7)**98-104.
- Basavaraj, K., Devakumar, N., Latha, B. and Somanatha, A.C. 2015.** Effect of organic liquid manure, jeevamrut and panchagavya on yield of frenchbean (*Phaseolus vulgaris* L.). *Proceedings of National Symposium on Organic Agriculture*, pp.111.
- Bhattarai, B.P. and Maharjan, A. 2013.** Effect of organic nutrient management on the growth and yield of carrot (*Daucus carota* L.) and soil fertility status. *Nepalese Journal of Agriculture Science*, **8(11)**:16-25.
- Bouyoucos, G.J. 1962.** A Hydrometer method improved for making particle and size analysis of soil. *Agronomy Journal*, **6(5)**464-466.
- Brady, N.C. 1983.** *The nature and properties of soil.* Mac Millan Publishing Co. New York and Collier Mac Millan Publisher, London.
- Ceronio, G.M., Engelbrecht, G.M. and Mbatha, A.N. 2012.** Organic fertilizer use and the potential influence on cabbage (*Brassica oleracea* L. var. *capitata*) plant and soil nutrient composition. *ActaHorticulturae*, **1(5)**: 265-272.
- Choudhary, R.S., Anchal-Das, Patnik, U.S. and Das, A. 2003.** Organic farming for vegetable production using vermicompost and FYM in Kokriguda watershed of Orissa.

- Indian Journal of Soil Conservation*, 8(31): 203-206.
- Dasappa, R., Rao, D.M., Chikkanna and Qadri, S.M.H., 2011**, Influence of organic manures on the soil microbial activity in the irrigated mulberry garden. *Green Farming*, 1(2):396-400.
- Devanda, P. 2020**. *Effect of organic manures and liquid formulations on growth, yield and quality of Okra [Abelmoschus esculentus (L.) Moench] cv. "Arka Anamika*. M.Sc. (Horti.) Thesis submitted to the Department of Horticulture, Rajasthan College of Agriculture, MPUAT, Udaipur.
- Gopakkali, P. 2014**. Effect of organic farming practices on growth, yield, quality and economics of onion (*Allium cepa*) in dry zone of Karnataka. *Indian Journal of Agronomy* 11(59) 336-340.
- Gopal, M., Gupta, A., Palaniswami, C., Dhanapal, R. and George, V. T. 2010**. Coconut leaf vermiwash: A bio-liquid from coconut leaf vermicompost for improving the crop production capacities of soil. *Current Science*, 5(98): 1202-1208.
- Gorakhnath and Singh, K. 2009**. Utilization of vermiwash potential on certain summer vegetable crops. *Journal of Central European Agriculture*, 9(10): 417-426.
- Hannah, K.A., Pandian, B.J. and Selva R. 2005**. Panchagavya spray produces tastier tomato fruit. *Agrobios News Letter*, 5(4): 4-5.
- Hartatik, W. and Widowati, L.R. 2013**. Manurefertilizers. *Inorganic fertilizer and bio fertilizers Indonesian Soil Research Center, Bogor. pp. 59-82*.
- Kumar, J., Phookan, D.B., Lal, N., Kumar, H., Sinha, K. and Hazarika, M. 2010**. Effect of organic manures and biofertilizers on nutritional quality of cabbage (*Brassica oleracea* var. *capitata*). *Journal of Eco-friendly Agriculture*, 10(10):114-119.
- Kumar, K. V. and Singaram, P. 2011**. Impact of organic manure and organic spray on soil microbial population and enzyme activity in green chillies. *Journal of Eco-friendly Agriculture*, 6: 10-12.
- Maheshwari, V.N., Shrikumaran, M.P., Rekha, G.S. and Kaleena, P.K.**

- 2016.** Growth promoting effects of panchagavya on dolichus lablab under field experiment. *International Journal of Applied Science*, **5(4)**: 513-518.
- Mahmood, Y.A., Ahmed, F.W., Juma, S.S. and Al Arazah, A.A. 2019.** Effect of solid and liquid organic fertilizer and spray with humic acid and nutrient uptake of nitrogen, phosphorus and potassium on growth, yield of cauliflower. *Plant Archives*,**10(19)**:1504-1509.
- Natarajan, K., 2002.** Panchagavya - A Manual. In: *Panchagavya Boon to organic farming*. Eds. International Book Distributing Co., Lucknow (India), pp.39-40.
- Negi,E., Shailaja, P., Pant, S.C., Kumar, S., Bahuguna, P., Mekap, B. and Nautiyal, B.P. 2017.** Effect of organic manures and bifertilizers on growth, yield, quality and economics of broccoli (*Brassica oleracea var. italica*) cv. 'Green Head' under high-hill conditions of Uttarakhand. *International Journal of Advanced Biological Research*, **7(1)**:96-100.
- Oleson, S.R., Cole, C.V., Watanabe, F.S. and Dean, L.A. 1954.** Estimation of available phosphorus in soil by extraction with sodium bicarbonate. *USDA circular*, No. 939, Washington DC, USA.