

***In vitro* Effect of Alum on Microbes Associated with Foliar Phytoplasma Disease of Noni**

ABSTRACT

Foliar phytoplasma disease of Noni (*Morinda citrifolia*) has a devastating socioeconomic impact on host crop species and a huge problem to farmers and scientists. Therefore, this study investigates the *in vitro* effect of alum on microorganisms associated with foliar phytoplasma disease (FPD) of Noni. Bacteria and fungi were isolated and identified by culture-dependent technique using standard microbiological procedures and susceptibility of species were evaluated by disc and agar well diffusion techniques and inhibition zones (IZs) were measured in millimeter (mm). The microbes identified were *Bacillus subtilis*, *B. tequilensis*, *Brevundimonas vesicularis*, *B. cereus*, *Staphylococcus aureus* and *Serratia* species as well as *Aspergillus flavus*, *A. fumigatus*, *A. niger* and *Penicillium* species. Relative abundance (%) of bacterial species occurred in the order; *S. aureus* > *B. tequilensis* > *B. cereus* and least being *Br. vesicularis* whereas fungi were *A. fumigatus* (75%), *A. niger* and *A. flavus* (50%) respectively and *Penicillium* (25%) species. Susceptibility test efficacy of Alum was concentration dependent with *Serratia* sp. as (6.0mm) and *S. aureus* (4.0mm) whilst *A. flavus* (36.0mm), *A. fumigatus* (32.5mm), *Penicillium* (30.2mm) and *A. niger* (30.0mm) at 5.0% Alum concentrations as the largest IZs respectively. The minimum inhibitory concentration (MIC) of Alum against all the microflora were < 2.0gm/mL except for *S. aureus* which portends a good future prospects in agronomy. However, higher concentrations of Alum compared favourably with Ketoconazole (control) but showed wide disparity with Ofloxacin (control). This study, however, may proffer solution for the prevention and management of FPD or foliar related diseases by periodic spraying or fumigation with Alum.

Keywords: Foliar phytoplasma disease, alum, noni, fumigation, ofloxacin, ketoconazole.

1. INTRODUCTION

Noni (*M. citrifolia*) is a plant with tremendous socioeconomic significance and most popularly cultivated plant of the genus *Morinda* and belongs to the family (Rubiaceae).

Originally a native of Southeast Asia but now grown in a variety of tropical and subtropical environments [1,2,3]. Noni became popular following its usage as food as well as for therapeutic and pharmaceutical purposes [4-7]. As a result, several parts of the plant such as leaves, roots, fruits, stem bark are being studied due to wide spectrum bioactivity and therapeutic importance [8,9].

A plethora of diseases have been associated with Noni shoot such as leaf blight, shot hole, black leaf spot, anthracnose, sooty mould, cracking and rot, stem cancer and blight, black flag, foliar phytoplasma disease, eventually culminating in plant death ([10-13]. FPD of Noni present profound reduction in growth, stunted or dwarfed with puckers upwards and inwards as well as lamina shrinkages from petiole to the tip [12]. However, this pathology features the connection with microorganisms, plant and climate as demonstrated by the “disease triangle concept” [14,15]. Disease development requires an adaptive plant, a harmful microorganism and appropriate natural conditions, absence of ideal conditions for any of these three elements may be negatively impacted. There is paucity of literatures on the microbiology of Noni shoot especially those associated with FPD as well as its prevention, control or management. However, earlier reports have shown that microbial community structures of Noni shoot to consist of *Enterobacter*, *Bacillus*, *Serratia*, *Pelomonas* and *Ochrobactrum* as well as *Phytophthora*, *Sclerotiumrol*, *Colletotrichum* and *Rhizopus* [10,16,17].

Potash alum, a natural substance have been known to possess antimicrobial properties and would be used as a treatment agent against foliar phytoplasma disease. It is nontoxic, affordable and eco-friendly. It has found application on/in several items and processes beneficial to humans such as drug, food processing and preservation, domestic and industrial water treatments, adjuvant to potentiate immune response in humans, anticariogenic agent, cosmetics and pharmaceutical industries as well as for treatment of ‘Ich and white eye’ diseases in fishes [18-22]. The study was designed to investigate the *in vitro* effect of Alum on microbes associated with foliar phytoplasma disease of Noni.

2. MATERIALS AND METHODS

2.1 Collection of samples

Typical features of the affected shoot in the field with FPD are stunted growth, reduced leaf size with puckers (upward and inward) and lamina shrinkages from petiole to the tip (**Plate 1**). Such samples were collected from Dilomat farm and conveyed to the Department of Microbiology, Rivers State University, Port Harcourt for analyses.



Plate 1. Foliar phytoplasma disease of Noni, sandwiched with the uninfected. Infected shoot are shorter with reduced leaves shrinking upward and inward.

2.2 Preparation of Alum and discs

Potassium aluminium sulphate (Vickers Laboratories, Ltd, England) was prepared by reconstituting appropriate quantity in sterile distilled water (w/v) such as 2.0,3.0,4.0 and 5.0g in 100ml to obtain 2- 5% respectively. Sterile discs (6mm diameter) were made of Whatman filter paper using an office perforating device.

2.3 Preparation of samples for microbiological analysis

A 25gm portion of FPD samples were washed, dried and blended with a mechanical grinder in 225ml of sterile normal saline to obtain the homogenate. Further decimal dilutions were

carried out before inoculation of 0.1ml aliquot onto surface-dried pre-sterilised nutrient agar (NA) for bacteria and Sabouraud's dextrose agar (SDA) for fungi. Followed by incubation at 37°C for 24h and 25±2°C for 2 or more days for bacteria and fungi respectively.

2.4 Macroscopy and microscopy of microbial cultures

Discrete microbial colonies were phenotypically identified according to their shape, colour, texture and contour and microscopically under oil-immersion objective, followed by biochemical tests for bacteria. Additionally, x10 and x40 magnification by infiltration with lactophenol cotton blue was carried out for fungi [23,24](Forbes et al., 2007; Cheesbrough, 2016). Cultures of bacteria were purified and preserved on NA at 5°C and each fungus was further grown for viability, confirmed and maintained on SDA slants at 5°C for further assay.

2.5 Antibiogram procedure

Antimicrobial susceptibility tests were carried out by agar well diffusion (AWD) and disc diffusion (DD) methods. After the growth of bacterial and fungal cultures for 18h (~ x 10⁸ overnight cultures) and 2-5days respectively on respective media, they were adjusted to 0.5 McFarland turbidity standard. Cultures were aseptically plated with various Alum concentrations and equidistantly spaced from one another and incubated at the different temperatures (for bacteria and fungi). Antibacterial and antifungal drugs; Ofloxacin (2,0mg) and Ketoconazole were used as controls respectively. Colony diameters were measured. The minimum inhibitory concentration (MIC), i.e., the lowest concentration able to inhibit any visible bacterial growth, was determined by measuring cell growth after 18-24h incubation at 37°C with the various concentrations used [25](Harvey, 2012).

2.6 Statistical Analysis

Means of duplicate measurements were determined for each sample using Microsoft Excel® 2016.

3. RESULTS

Bacterial profile associated with foliar phytoplasma disease (FPD) and their susceptibility to Alum concentrations are presented in **Table 1**. The Inhibitory effect of Alum on bacterial isolates was concentration dependent using Agar well diffusion (AWD (values in bold) and disc diffusion (DD) techniques. At 5.0% Alum concentration had the largest inhibitory zones. *Serratia* species had the larger IZs but the inhibitory effect was much more pronounced with the control whereas the results of other species fluctuated. However, IZs by AWD technique were much larger than those of DD whilst the control showed exceptionally the largest IZs.

Table 1. Inhibitory effect of Alum concentrations on bacteria associated with foliar phytoplasma disease of Noni using to AWD and DD techniques

Bacteria	Concentration of Alum (%) and Mean IZ (mm)									
	AWDT				DDT				Control	
	2.0%	3.0%	4.0%	5.0%	2.0%	3.0%	4.0%	5.0%	2.0mg/mL	
<i>Bacillus subtilis</i>	1.0	3.0	3.0	4.0	0.0	1.5	1.8	2.0	49.5	32.0
<i>B. tequilensis</i>	1.0	2.0	3.0	4.0	0.0	2.0	2.5	3.0	45.6	25.5
<i>St. aureus</i>	0.0	1.5	2.5	3.5	0.0	2.5	3.5	4.0	41.0	22.5
<i>Serratia</i> sp.	2.0	3.0	4.0	6.0	0.0	2.5	3.0	4.0	51.0	22.0
<i>Br. vesicularis</i>	1.0	2.0	3.0	4.0	0.0	1.6	1.8	2.0	43.0	30.5
<i>B. cereus</i>	1.0	1.5	2.0	4.0	0.0	1.8	2.0	2.5	42.0	33.8

Legend: AWDT = Agar well diffusion technique (in **Bold**); DDT = Disc diffusion technique = IZ = Inhibition zone (measured in millimeter (mm)). Control = Ofloxacin.

Each value represents mean of 2 experiments.

The inhibitory effect of Alum against fungal isolates associated with foliar phytoplasma disease of Noni are presented in **Table 2**. Susceptibility of fungal isolates to Alum exhibited similar trend of dose dependence, with 5% concentration having the larger IZs. The inhibitory pattern of different concentrations of Alum revealed more profound effects on the isolates by AWD than with DD (**Table 2**). Of the two genera of fungi; *Aspergillus* and *Penicillium* the most sensitive to Alum at various concentrations was *Aspergillus flavus*,

followed by *A. fumigatus* and *Penicillium* sp with *A. niger* as the least. However, higher concentrations of Alum also depicted comparable results with those of control.

Table 2. Inhibitory effect of Alum concentrations to fungi associated with foliar phytoplasma disease of Noni using AWD and DD techniques

Fungi	Concentration of Alum (%) and Mean IZ (mm)									
	AWDT				DDT				Control	
	2.0%	3.0%	4.0%	5.0%	2.0%	3.0%	4.0%	5.0%	2.0mg/mL	
<i>Aspergillus flavus</i>	28.0	30.0	34.0	36.0	22.0	25.0	30.0	34.0	39.0	24.8
<i>A. fumigatus</i>	22.0	29.0	30.3	32.5	15.0	25.0	29.5	30.0	35.2	25.0
<i>A. niger</i>	15.0	19.4	21.0	30.0	10.0	18.2	20.4	22.0	25.3	20.2
<i>Penicillium</i> sp.	20.0	22.5	25.0	30.2	20.0	21.4	23.0	29.6		35.0

22.53

Legend: AWDT = Agar well diffusion technique (in **Bold**); DDT = Disc diffusion technique; IZ = Inhibition zone (measured in millimeter (mm)). Control = Ketoconazole.


Each value represents mean of 2 experiments.

The minimum inhibitory concentration (MIC) of Alum against the microflora associated with foliar phytoplasma disease (FPD) are depicted in **Table 3**. All the species had MIC of < 2.0gm/mL except *Staphylococcus aureus*.

Table 3. MIC of Alum against bacterial and fungal species associated with FPD of Noni

Bacteria	Alum		Fungi	gm/mL
	gm/mL			
<i>B. subtilis</i>	< 2.0		<i>A. flavus</i>	< 2.0
<i>B. tequilensis</i>	< 2.0		<i>A. fumigatus</i>	< 2.0
<i>S. aureus</i>	> 2.0		<i>A. niger</i>	< 2.0
<i>Serratia</i> sp	< 2.0		<i>Penicillium</i> sp	< 2.0
<i>Br. vesicularis</i>	< 2.0			

Relative abundance (%) of bacteria associated with foliar phytoplasma disease of Noni are presented in **Figure 1**. Basically, four genera bacteria were of identified with the predominant genera as *Bacillus* followed by *Staphylococcus* and *Serratia* with *Brevundimonas* as the least. The most abundant single species was *Staphylococcus aureus* followed by *Bacillus tequilensis* and *B. cereus*, the least being *Brevundimonas vesicularis*.



Figures 1. Relatiive abundance (%) of bacterial isolates associated with foliar phyoplasma disease of Noni

The relative abundance (%) of fungal isolates associated with foliar phytoplasma disease of Noni are shown in **Figure 2**. The data shows that there are two (2) genera of fungi; *Aspergillus* and *Penicillium*. The dominant genus, *Aspergillus*, has *A. fumigatus* (75%) as the most abundant species, followed by *A. niger* and *A. flavus* (50%) respectively, the least being *Penicillium* species (25%).




Figure 2. Relative abundance (%) of fungal isolates associated with foliar phytoplasma disease of Noni.

4. DISCUSSION

Foliar phytoplasma disease (FPD) has been a challenge to farmers and persistent occurrence of such exogenous pathology can have huge negative impact on income and well-being. The inhibition of microbes of which microorganism is one of the major components of the 'disease triangle' can proffer solution to the prevention and/or management of the foliar phytoplasma disease. The microbes associated with FPD belongs to six genera; *Bacillus*, *Staphylococcus*, *Serratia*, *Brevundimonas*, *Aspergillus* and *Penicillium* as revealed in this work.

Literature on foliar phytoplasma disease of Noni has been earlier reported [12](Sarwade *et al.*, 2015) but devoid of the aetiologic agents or microbes associated with the disease. However, two major microbial genera from this study; *Bacillus* (*B. tequilensis*) and *Asperillus* (*A. niger*) have been reported to producing enzymes (e.g., pectinases; polygalacturonase, pectin lyase and pectinesterase) capable of degrading starches, hemicelluloses, celluloses and other cell wall components [26,27,28,29](Maldonado *et al.*, 2002; Hubka *et al.*, 2013;; Shah *et al.*, 2013; Aarti and Khusro, 2015). Beyond the role of degradation, *Aspergillus* is also a known phytopathogen [30](Lee *et al.*, 2016). These microbial attributes can negatively impact the tender morphological architecture of Noni foliage to present the typical condition of foliar phytoplasma disease (**Plate 1**).

The relative abundance (%) of foliar microbes may be attributed to soil, weather and other environmental conditions which in turn impacts the microbial community structure where the plant is grown [16,30,31,32]Jeffrey *et al.*, 1999; [30]Lee *et al.*, 2016; [16]Liu *et al.*, 2014; [32]Brown, 2021). The antimicrobial potentials of Alum has been well documented in

previous studies [22,33,34]Amadi et al., 2019a,b; [22]Ade and Makode, 2021) and validates the present study. The low MIC of Alum of < 2.0 gm/mL suggests future prospects for its agronomic applications. As an eco-friendly natural agent, it can be explored and adopted to ameliorate or combat foliar phytoplasma disease as well as other foliar pathogens. Although, there was wide disparity in relation to Alum's effect compared to the antibiotic (control), its relative comparison at higher doses with Ketoconazole underscores antifungal efficacy.

5. CONCLUSIONS

The research revealed the identification of six microbial genera; four of bacteria (*Bacillus*, *Staphylococcus*, *Serratia* and *Brevundimonas*) and two of fungi (*Aspergillus* and *Penicillium*) associated with foliar phytoplasma disease of Noni. Alum treatment against these microbes exhibited appreciable MIC and antimicrobial efficacy. This presupposes that Alum can be explored and used to ameliorate future challenges in agronomy especially foliar related diseases.

DISCLAIMER

The samples used for this research are commonly used in our area of research and country. There is absolutely no conflict of interest between the author and producers of Noni fruits because we do not intend to use these samples as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the author.

Delete all REFERENCES in yellow. Uninfected Noni foliage was added to PLATE 1 for clarity.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. Wang Z, Dou R, Yang R, Cai K, Li C, Li W. Changes in Phenols, Polysaccharides and Volatile Profiles of Noni (*Morinda citrifolia* L) Juice during Fermentation. *Molecules*, 2021; 26, 2604.
2. Almeida ES, Oliveira D, Hotza D. Properties and Applications of *Morinda citrifolia* (Noni): A Review. *Comprehensive Reviews in Food Science and Food Safety*. 2019;18:883-909.
3. Nelson SC, Elevitch CR. Noni: The Complete Guide for Consumers and Growers, 1st ed.; Permanent Agriculture Resources: Holualoa, HI, USA. 2006; 67-80.
4. Deng S, West BJ, Jensen CJ. A quantitative comparison of phytochemical components in global noni fruits and their commercial products. *Food Chemistry*, 2010;122(1), 267–270. <https://doi.org/10.1016/j.foodchem.2010.01.031>
5. Sibi G, Parul C, Sayak A, Ravikumar KR. (2012). Phytoconstituents and Their Influence on Antimicrobial Properties of *Morinda citrifolia* L. *Research J Medicinal Plants*. 2012; 6(6):441-448.
6. Assi RA, Darwis Y, Abdulbaqi IM, Khan AA, Vuanghao L, Laghari MH. *Morinda citrifolia* (Noni): A comprehensive review on its industrial uses, pharmacological activities, and clinical trials. *Arabian J Chemistry* 2017;10:691-707.
7. Amadi LO, Nwala H. Preliminary investigation on microflora associated with fermentation of noni (*Morinda citrifolia*) fruit and its nutritional components. *International Journal of Biotechnology and Microbiology* 2021;3, (2):01-04.
8. Langbang D, Rekha, Menon T. Role of Aqueous, Methanol and Ethanol leaf Extract of *Morinda citrifolia* (Indian Noni) in inhibiting on clinical *Pseudomonas aeruginosa* isolates. *J Med Sci Clin Res.*, 2018;06(06):984-992.
9. Ali M, Kenganora M, Manjula SN. Health Benefits of *Morinda citrifolia* (Noni): A Review. *Pharmacognosy Journal*, 2016;8(4):321-334.
10. Nelson SC, Abad ZG. *Phytophthora morindae* a new species causing black flag disease on Noni (*Morinda citrifolia*) in Hawaii. *Mycologia* 2010;102(1):122-134.
11. Jackson G, McKenzie E. *Guignardia morindae* and Noni shot-hole disease (313). *Pacific pests, Pathogens and Weeds – Online edn*. Produced with Support from the Australian Centre International Agriculture Research, Project PC/2010/090. 2013.

12. Sarwade PP, Sarwade KP, Chavan SS. (2015). New First Report of Foliar Phytoplasma Disease on Bartondi Plant in India. J Plant Pathol Microbiol., 2015;6(3):260. doi:[10.4172/2157-7471.1000260](https://doi.org/10.4172/2157-7471.1000260).
13. Jumar, Yusriadi. Surtinah. Control of Anthracnose disease in Chili with Several doses of Noni leaf extracts. IOSR J Agric and Vet. Sc., 2020;13(4):33-38.
14. Steven RB. In: Plant Pathology an Advanced Treatise, Vol. 3, J.G. Horsfall and A.E. Dimond, eds. Acaademic Press, NY., USA. 1960;357-429.
15. Francl LJ. The disease triange; A Plant pathological paradigm revisited. The Plant Health Instructor. 2001. DOI: 10.1094/PHI-T-200-0517-01.
16. Liu Y, Cheng C, Yao S, Xu P, Cao Y, Li J, Wang J, Tan W, Cheng C. Composition and diversity of endophytic bacterial communities in noni (*Morinda citrifolia* L.) seeds. Int. J. Agric. Pol. Res., 2014;2(3):098-104.
17. Kavitha PG, Umadevi M. Medicinal Properties and Pests and Diseases of Noni – A Review. Research J Pharmacognosy and Phytochemistry, 2016; 8(1):41-48.
18. Efiuwewewere BJO, Amadi LO. Effects of Preservatives on the Bacteriological, chemical and sensory qualities of mangrove oyster (*Crassostrea gasar*) harvested from the Niger Delta Region, Nigeria. British Journal of Applied Science and Technology; 2015;5(1):76-84.
19. Al-Talib H, Nasir MSIN, Yaziz H, Zulkafli FN, Adani AN, Rashidi NIA, Murugaiah C, Shaari AS. Potassium aluminium sulphate (Alum) inhibits growth of Human axillary malodour-producing skin flora in vitro. J Clinical Health Science, 2016;1(1): 59-63.
20. Amadi LO. A review of Antimicrobial Properties of Alum and Sundry Applications. Acta Scientific Microbiology 2020;3.4:109-117. DOI:[10.31080/ASMI.2020.03.0553](https://doi.org/10.31080/ASMI.2020.03.0553).
21. Al-Al-Hamdoni SA and Al-Rawi AM. Elevation of the Inhibitory Action of Antimicrobials (Ciprofloxacin and Chlorhexidine) by some Natural Materials against Three Periodontal Pathogens Rafidain J Science 2020;29(2):1-9.
22. Ade GV. Makode PM. (2021). Ich and White eye disease management in Fishes using novel components. Int J Res Biosci Agric Technol, 2021;(17): 551-558.
23. Forbes BA, Sahn DE, Weissfeld AS. 2007. Bailey and Scott's Diagnostic Microbiology. International edition. 12th edn. Mosby, Inc., an affiliate of Elsevier, Inc., USA. 2007.
24. Cheesbrough M. District laboratory practices in tropical countries. Volume 2 part 2 Cambridge University Press, UK. 2006.
25. Harvey R. 2012. Lippincott's Illustrated Reviews: Pharmacology. 5th Edition. Lippincott and Williams & Wilkins, Philadelphia, USA. 2012.
26. Maldonado MC, Cáceres S, Galli E, Navarro AR. Regulation of the production of polygalacturonase by *Aspergillus niger*. Folia Microbiol (Praha) 2002;47(4):409-12.
27. Hubka V, Kolařík M, Kubatova A, Peterson SW. (2013). Taxonomic revision of *Eurotium* and transfer of species to *Aspergillus*. Mycologia 2013;105:912-37.
28. Shah KP, Chandok HK, Rathore P, Sharma MV, Yadav M, Nayarisseri SA. Screening, isolation and identification of polygalacturonase producing *Bacillus tequilensis* strain EMBS083 using 16S rRNA gene sequencing. Eur J Biol Sci 2013;5(1):09-13.
29. Aarti C, Khusro A. Discovery of Polygalacturonase Producing *Bacillus tequilensis*

- strain ARMATI using 16S rRNA Gene Sequencing. *Asian J Pharm., Clin Res.*, 2015;8(5):58-62.
30. Lee S, Park MS, Lim YW. (2016). Diversity of Marine-Derived *Aspergillus* from Tidal Mudflats and Sea Sand in Korea. *Mycobiology*, 2016;44(4): 237-247.
31. Jefferey SB, Daniel PR, Estelle RC. (1999). Microbial community structure and function in the spermosphere as affected by soil and seed type. *Can. J. Microbiol.*, 1999;45: 138-144.
32. Brown S. Effect of Climate on Plant Diseases. *J Plant Pathol Microbiol* 2021;12:305.
33. Amadi LO, Ekechi JS, Seth MA. Bioefficacy of Turmeric Rhizome Extracts with Alum on Microbes: An in Vitro Approach. *American Journal of Microbiological Research*, 2019a;7(2): 51-56. doi: 10.12691/ajmr-7-2-3.
34. Amadi LO, Udoh EJ, Thompson RU, Benjamin RG, Nmomo FW. Antimycotic Potential of Alum on Postharvest Deterioration of Tomato. *Microbiology Research Journal Int'l*, 2019b;29(3):1-10.2019. DOI:10.9734/MRJI/2019/v29i330165.