

Original Research Article

Study of Parental Polymorphism and Allelic Variation for Grain Quality and Yield Traits in Rice (*Oryza Sativa* L.) using SSR Markers

Sanket Rathi^{1*}, Sameer Upadhyay¹, P. K. Singh¹, Rajesh Kumar², Pallavi³, Prashant Bisen⁴, Bapsila Loitongbam⁵ and Sanchika Snehi⁶

¹Department of Genetics and Plant Breeding, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi-221005, Uttar Pradesh, India

²Department of Plant Biotechnology, RGSC, Banaras Hindu University, Varanasi-231001, Uttar Pradesh, India

³Karlsruhe Institute of Technology, Germany

⁴Department of Genetics and Plant Breeding, Narayan Institute of Agricultural Sciences, Gopal Narayan Singh University, Jamuhar, Rohtas-821305, Bihar, India

⁵Department of Genetics and Plant Breeding, College of Agriculture, Central Agricultural University (Imphal), Pasighat-791102, Arunachal Pradesh, India

ABSTRACT:

Aim: Identification of polymorphic markers is prerequisite for conducting any QTL mapping experiment because if the parents are polymorphic for the traits of interest, then further selection of plants in the progenies becomes easy. Hence, the objective of the present study was to identify polymorphic markers for grain quality and yield related traits among the parental lines Improved Samba Mahsuri and Badshahog.

Place and Duration of Study: It was carried out at Molecular Breeding Lab, Department of Genetics and Plant Breeding, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi - 221 005, India, during 2019

Methodology: Two parents Improved Samba Mahsuri and Badshabhog were used for the present study. The DNA extraction was done as per the CTAB method suggested by Murray and Thompson. Standard PCR protocol was followed.

Results: For this a total of 576 randomly selected microsatellite SSR markers distributed over all the 12 chromosomes of rice including 26 gene specific markers related to aroma, cooking and eating quality, grain dimension and yield related traits were used. Overall, 96 markers including 4 gene specific markers were found to be polymorphic between the two genotypes indicating a total polymorphism percentage of 16.67%. The highest polymorphism percentage was recorded on chromosome 6 (26.67%) followed by chromosome no 4 (21.43%) and the lowest polymorphism percentage was observed on chromosome 10 (8.93%). The gene specific markers nksbad2, ARO7, BADEX7_5 and SSI were found to be polymorphic.

Conclusion: Based on the present study it may be concluded that the polymorphic markers identified will further be utilized in genotyping of $F_{2:3}$ population, linkage analysis and mapping QTL's for grain quality and yield traits.

Key words: Rice, aroma, cooking and eating quality, parental polymorphism, QTL, SSR.

1. INTRODUCTION:

Rice is an indispensable cereal to the world providing nutrition to about 21% of the global population and two third of the South East Asian population [24]. In 2019, global rice production was 755.47 million tons from 162.05 million ha of area. Among the 117 different rice producing countries, China ranks first with a production of 209.61 MT followed by India with 177.64 MT in 2019. Both these countries account for more than 50% of the total rice production. Asian countries contribute 89.6 % of the total world production (677.27 MT) followed by Africa (38.77 MT), America (35.32 MT), Europe (4.02 MT) and Oceania (0.76 MT) [2]. To meet the ever-increasing demand, several new varieties are bred every year to achieve the highest yield along with resistance to various biotic and abiotic stresses. But, the grain qualities in these varieties were less prioritized. Grain quality is very hard to define as it comprises of several different facets such as grain appearance quality, milling quality, nutritional quality, eating quality, and cooking quality [11,4]. Moreover, the preference and taste varies from one region to another, what may be preferable in one region may not necessarily be preferable in another region. [6]. Nearly 40% of

people around the globe particularly in the USA, Canada, Europe, Middle East and South East Asian countries desire to eat aromatic rice irrespective of grain types.

Most of the quality traits follow quantitative inheritance and have complex genetic architecture because of the considerable influence of genotype, environment, and their interactions [1,18]. Phenotype-based classical breeding approaches are time-consuming, laborious and are inefficient in improving traits that are governed by quantitative trait loci (QTL), such as yield and other grain quality traits which show continuous phenotypic variation and lack discrete phenotypic segregation in the progeny [32,22]. However, several recent developments such as the use of potential donor parents, identification of QTLs regions associated with important grain qualities, and identification of robust marker have increased our understanding of the genes, pathways, and molecular mechanisms determining overall quality traits in rice [11,20]. Breeding programs should be directed in to develop rice varieties with better eating quality and to meet out the demands of the local consumers as well as emerging food processing industry.

Mapping of QTLs for grain quality traits in rice is an important forward genetic approach and using these dissected complex regions in MAS and gene discovery [27]. With the advent of several molecular techniques, the use of molecular markers has increased to a large extent in the recent past. Among the PCR-based markers, the SSR markers have proved to be very effective tools and the first choice of breeders in the study of genetic diversity and organism relationships due to their several advantages such as they are co-dominant, highly polymorphic in nature, evenly distributed in the genome, efficient, less quantity of DNA is required, are highly cost-effective and transferability [15]. These SSR markers can be effectively used in studying the genetic polymorphism *i.e.*, occurrence of multiple alleles at a single locus, where at least two alleles occur with a frequency greater than one percent [12]. Also, they can be used in population structure analysis, gene mapping and tagging, linkage map construction, tracing marker-trait association, Marker Assisted Selection (MAS) and others. [23]. So, developing rice varieties with desired grain quality *viz.*, aroma, nutritional, cooking and eating quality with high yield as well as resistance to various abiotic and biotic stresses by using molecular techniques and breeding tools will help to alleviate several malnutritional problem, as well as will help farmers and food industries personnel to fetch high price of their produce. Thus, the present study was aimed to identify informative polymorphic SSR markers between two diverse parents *viz.*, Improved Samba Mahsuri and local rice germplasm

Badshabhog for yield and quality traits in rice. The research was also planned to study the influence of the type of the repeat motif on polymorphism.

2. MATERIALS AND METHOD:

2.1 Plant Material used in the study:

The experimental plant material for this study comprised of a highly contrasting rice parents for desirable grain quality and yield traits. The first parent is short grain aromatic landrace 'Badshabhog', a popular and preferred line among the aromatic rice breeders and second variety is 'Improved Samba Mahsuri' a fine grain, medium slender non-aromatic rice, popular specially for grain quality. Details of all the traits of both the parents are presented in table 1. These two are popular varieties and being used as a source for desirable grain traits in rice breeding. Both the parents were raised by following all recommended package of practices as per recommendation of the eastern plain zone. Twenty-one days old seedlings of Improved Samba Mahsuri and donor Badshabhog were transplanted in the field for evaluation. Leaf samples were collected 14 days after transplanting and were preserved at -20°C for parental polymorphism survey after DNA extraction.

Table 1. Details of the parents used in parental polymorphism experiment:

| Sl.no | Trait name | Recipient | Donor |
|-------|------------------------|--|---|
| 1 | Name | Improved Samba Mahsuri | Badshabhog |
| 2 | Type | Released variety in 2008 jointly by ICAR-IIRR and CSIR-CCMB Hyderabad, India | Local landrace collected by Institute of Agricultural Sciences, BHU, Varanasi, U.P, India |
| 3 | Parentage | Samba Mahsuri*4/SS1113 | Local landrace |
| 4 | Yield | 4.75-5.0 tonnesha ⁻¹ | 2.5-3 tonnesha ⁻¹ |
| 5 | Days to 50 % flowering | 100-103 days | 120-123 days |
| 6 | Days to maturity | 130-134 days | 150-154 days |
| 7 | Plant height | 100-103 cm | 155-160 cm |
| 8 | Grain length | 8.15 mm | 5.53 mm |
| 9 | Grain breadth | 2.01 mm | 2.11 mm |
| 10 | Kernel length | 5.32 mm | 3.83 mm |
| 11 | Kernel breadth | 1.78 mm | 1.85 mm |
| 12 | Aroma | Non aromatic | Highly aromatic |
| 13 | Amylose content | 23.84% | 18.46% |

2.2 Genomic DNA Isolation and quality check:

The genomic DNA of both the parents were extracted by the CTAB method [16], with some modifications. 100 mg of young leaves was weighed and the genomic DNA was extracted by using extraction buffer (10% CTAB, 1M Tris HCl (pH 8.0), 0.5M EDTA (pH 8.0), 5 M NaCl, and 0.2 % β -mercaptoethanol) preheated at 60° C. The quality and quantity of extracted DNA was estimated on 0.8 % agarose gel along with a standard ladder and compared with band intensity and thickness. DNA quantification and purity were checked by measuring the O.D values at 260 and 280 nm using a NanoDrop ND100 spectrophotometer. The ratio of UV absorbance at A_{260}/A_{280} ranged between 1.89-1.96, and this ratio indicates good-quality DNA. The quantity of DNA in the isolated samples ranged from 1288.50 to 1701.30 ng/ μ l. After quantification, the DNA samples were diluted with TE buffer, so as to make final concentration of the DNA as 50 ng / μ l stored at 4°C.

2.3 Details of SSR markers used in the study:

A total of 576 randomly selected SSR markers including 26 gene specific markers distributed on 12 chromosomes of rice were used to identify polymorphism between Improved Samba Mahsuri and Badshahog. This was confirmed and tested by using graphical genotypes (GGTs) software package 2.0.[29]. GGT 2.0 is particularly a plant-breeding software package that helps to visualize data of markers with known map positions on a genetic map and displays estimated lengths of genomic compositions and distribution of polymorphic markers across the length of chromosome according to their physical positions (Mb) as colored chromosome bar segments. The information regarding chromosomal location, sequences of primers (forward and reverse), physical position (SSR start and SSR end) and number of repeat motifs were obtained from Gramene markers database (<https://www.gramene.org/>). RAP-DB (The Rice Annotation Project Data Base) was also used for those markers whose information was not available in the Gramene database, by using its BLAST tool and submitting query sequence in FASTA format to run a search against DNA database (blastn) (<https://rapdb.dna.affrc.go.jp/tools/blast>). Fig. 1 depicts the distribution of all the identified polymorphic markers across the length of 12 rice chromosome, according to their physical positions (Mb) on a genetic map using GGT software.

2.4 PCR amplification:

PCR was carried out in Eppendorf thermal cycler with a final reaction volume of 15 μ l containing 1.2 μ l of genomic DNA, 1.5 μ l of 10 X Taq assay buffer containing 15 mM MgCl₂, forward and reverse primer each 1 μ l (10 pmol/ μ l), 0.2 μ l of dNTPs (3 mM), 0.2 μ l Taq DNA polymerase(GeNei) and 9.9 μ l of HPLC water. The PCR reaction was performed under the following conditions: Initial denaturation at 94°C for 5 min, followed by 35 cycles of denaturation at 94°C for 45 seconds, annealing at the temperature specific to primer for 30 seconds, extension at 72°C for 30 seconds, followed by the final extension at 72°C for 5 min and hold at 4°C for ∞ . After completion of amplification, PCR products were stored at -20°C

2.5 Agarose gel electrophoresis:

The amplified products were resolved on 2.5% agarose gels in 1X TAE Buffer at 65V for initial 30 minutes followed by 90V for 1.5 hours. Gel was stained by Ethidium bromide (10 mg/ml) @ 2.5 μ l/1000 ml 1X TAE Buffer. The DNA fragments were then visualized under Gel documentation system (Gel DocTM XR+, BIO-RAD, USA) and the banding pattern was observed and recorded for further analysis.


3. RESULTS AND DISCUSSION:

In the present study, the parental polymorphism survey between the parents, Improved Samba Mahsuri and Badshabhog indicated a clear polymorphism and it is very important to understand this variation at molecular level. Among the 576 SSR markers used, 96 markers including 4 gene specific markers were found to be polymorphic between the two genotypes indicating a total polymorphism percentage of 16.67%. The polymorphic banding pattern of some of the markers are shown in gel picture Fig.2. The highest polymorphism percentage was recorded on chromosome 6 (26.67%) followed by chromosome no 4 (21.43%) and the lowest polymorphism percentage was observed on chromosome 10 (8.93%). The chromosome wise polymorphism percentage is stated in table 2. For QTL mapping genetically diverse parents are required to map the traits of interest [9] and a high percentage of polymorphism between the two parents used in the study shows that the two parents are genetically diverse. The identified polymorphic markers can further be effectively used in QTL mapping experiment for preparation of linkage map and identification of QTL's related to grain quality and yield traits in mapping population derived from both the parents.

Table 2. Chromosomal wise polymorphism percentage of SSR markers between the parents Improved Samba Mahsuri and Badshabhog

| Chromosome no. | Total No. of SSR marker used | No. of Polymorphic markers obtained | Polymorphism (%) |
|----------------------|------------------------------|-------------------------------------|------------------------|
| Chromosome 1 | 51 | 10 | 19.61 |
| Chromosome 2 | 47 | 7 | 14.89 |
| Chromosome 3 | 55 | 8 | 14.55 |
| Chromosome 4 | 42 | 9 | 21.43 |
| Chromosome 5 | 52 | 8 | 15.38 |
| Chromosome 6 | 30 | 8 | 26.67 (Highest) |
| Chromosome 7 | 45 | 8 | 17.78 |
| Chromosome 8 | 53 | 11 | 20.75 |
| Chromosome 9 | 48 | 8 | 16.67 |
| Chromosome 10 | 56 | 5 | 8.93 (Lowest) |
| Chromosome 11 | 45 | 7 | 15.56 |
| Chromosome 12 | 52 | 7 | 13.46 |
| Total | 576 | 96 | 16.67 |

Fig. 1. Genetic linkage map showing distribution of polymorphic markers across the 12 chromosomes.






Fig. 2. Gel picture showing parental polymorphism survey with SSR markers among the parents. P1- Improved Samba Mahsuri, P2- Badshabhog, L- Ladder 50 bp.

The recent past studies on parental polymorphism also revealed comparable results. The polymorphism percentage highest for chromosome 6 and lowest on chromosome 10 was also reported by Shivani *et al.*, [23]. Similarly, Hableet *et al.*, [13] reported polymorphism percentage of 29.02% between the two parents Rajendrakasturi and URG-30. The highest polymorphism was reported on chromosome 4 (40.96%) whereas the lowest polymorphism was observed in chromosome 9 (16%). Chanduet *et al.*, [7] found a total polymorphism percentage of 20.75% by using 800 SSR markers between BPT5204 and *O. rufipogon* WR119. A polymorphic percentage of 6.93 was reported by Kulkarni *et al.*, [14] using 1,904 genomic SSR markers among the parents IR58025A and KMR-3R.


In the present study a total of 96 polymorphic markers were identified, out of which 20 SSR markers were reported to be associated with QTLs for different quality traits and the 4 gene specific markers were associated with genes of aroma and eating and cooking quality traits of rice from previous studies. Among the 4 gene specific markers 3 markers were found to tightly linked with the genes associated with aroma content in rice *viz.*, ARO7 [26], BADEX7_5 [21], nksbad2 [1], and one marker SS [3]

was found to be associated with cooking and eating quality traits. The gene specific primer ARO7, BADEX 7_5 and nksbad2 is linked with BAD2 or badh2 gene with 8-bp deletion in the exon 7 which leads to aroma in rice. The gene specific primer SSI is linked to gene encoding granule-bound starch synthase I. The SSR markers RM19, RM217, RM219 flank the QTL associated with cooking and eating quality traits such as amylose content and alkali digestion value and RM19 was also linked with the trait kernel length after cooking. Bazrkar-Khatibani *et al.*, [5]. Xia *et al.*, [31] found that RM488, RM562, RM432 were linked with grain dimension traits such as grain length and grain breadth while RM432 was also found to be associated with amylose content in rice. Dai *et al.*, [10] reported that RM7180, RM3616, RM6836, RM3183 were significantly associated with grain dimension traits such as grain length and grain breadth. Three SSR markers CHR3_14, CHR3_20, CHR8_4 was found to be associated with aroma content in rice Singh *et al.*, [25]. Two SSR markers RM404, RM547 were found to be linked with aroma content and RM17 with amylose content in rice in a study conducted by Vemireddy *et al.*, [30]. Amarawathi *et al.*, [1] reported that RM80 and RM252 was found to be associated with aroma content in rice and RM217 and RM432 with cooking quality traits such as amylose content and alkali digestion value and grain dimension traits such as grain length and grain breadth. One SSR marker RM42 of chromosome 8 was found to be linked with grain aroma content in two studies conducted by Chen *et al.*, [8]. While Swamy *et al.*, [27] reported that RM5688 was found to be associated with amylose content and other cooking quality traits on chromosome 9.

Effect of the nature and form of the motif on polymorphism detection:

Within the set of 96 polymorphic marker, most consisted of dinucleotide motifs (52.08%), followed by trinucleotide motifs (22.92%), subsequently by tetranucleotide motifs (7.29%). Similarly group of either mono-dinucleotide (6.25%), mono-trinucleotide (1.04%), di-trinucleotide (2.08%), di-tetranucleotide (2.08%), tri-pentanucleotide (1.04%) and penta-di-mononucleotide (1.04%) were also found (Fig. 3). These findings are in accordance with the results of Narshimulu *et al.*, and Nicot *et al.*, [17,19]. The dinucleotide motifs showed a much larger polymorphism level than the trinucleotide, tetranucleotide and the other motifs classes. However, the number of repeat motifs does not affect the level of polymorphism rate. Among the most frequent motif *viz.*, dinucleotide motifs, (GA)_n repeats were more frequent (39.34%), followed by (CT)_n with 27.87% and least by (TA)_n with 1.64%. Among the trinucleotide motifs

(CCT)_n repeats were maximum. (ATAG)_n repeats characterized 30% of the tetranucleotide motifs followed by (AGAT)_n 20% whereas the other repeat classes (CATC)_n, (GAGT)_n, (GATG)_n, (TAAT)_n, (TCTT)_n each represented 10%. On the other hand, the number of repeats ranged from 5 to 46 for the different polymorphic SSR markers (Table 3). The findings of the physical position, type of repeat motif and no. of repeats of the polymorphic SSR markers in the present investigation will aid in formation of set of



polymorphic SSR's which can further be used in genotyping of mapping population for QTL mapping and marker-assisted selection in breeding.

Fig. 3. Frequency distribution of different types of SSR repeat motifs.(DN: Dinucleotide, TN: Trinucleotide, TtN: Tetranucleotide, MDN: Mono-Dinucleotide, MTN: Mono-Trinucleotide, DTN: Di-Trinucleotide, DTtN: Di-Tetranucleotide, TPN: Tri-Pentanucleotide, PDMN: Penta-Di-Mononucleotide, Un: Unknown)

CONCLUSION:

Thus, in the present study, it may be concluded that a total of 96 SSR markers including 4 gene specific markers were found to be polymorphic among the two parents indicating a total polymorphism percentage of 16.67%. Dinucleotide repeats were in much larger number as compared with other classes of nucleotides. These suitable polymorphic markers depending upon the nucleotide repeats will further be selected and utilized in genotyping of F_{2:3} population and identification of QTLs for grain quality and yield traits results in development of improved rice cultivars having a more economic value.

ACKNOWLEDGEMENT:

The authors are sincerely thankful to Department of Genetics and Plant Breeding, Institute of Agricultural Sciences B.H.U., Varanasi and UPCAR, Lucknow for providing the resources and facilities for execution of present experiment. This study was supported by Banaras Hindu University, Varanasi under IoE faculty grant project. The authors declare no conflict of interests.

Authors' contributions:

This work was carried out in collaboration among all authors. Author SR and PK designed the study, performed the molecular work and wrote the first draft of the manuscript. Author PKS designed and supervised the study, proof read and edited of manuscript. Authors RK, SU, PR, BP, SS provided inputs on designing the study and interpretation of results. All authors read and approved the final manuscript.

Table 3: Identified polymorphic markers between the two parents along with their complete details is mentioned in table below:

| Sr. no. | Name | Chr no. | Temp | Forward sequence | Reverse sequence | Map Distance | Motif | No. of repeats |
|---------|---------|---------|------|-------------------------|-------------------------|--------------|-------------|----------------|
| 1. | RM3252 | 1 | 57.4 | GGTAACTTTGTTCCCATGC | GGTCAATCATGCATGCAAGC | 0.3 | CT | 13 |
| 2. | RM220 | 1 | 53.1 | GGAAGGTAAGTGTTCAC | GAAATGCTTCCCACATGTCT | 4.42 | CT | 17 |
| 3. | RM259 | 1 | 56.6 | TGGAGTTTGAGAGGAGGG | CTTGTTCATGGTGCCATGT | 7.44 | CT | 17 |
| 4. | RM562 | 1 | 55.3 | CACAACCCACAAACAGCAG | CTTCCCCAAAGTTTTAGCC | 14.62 | AAG | 13 |
| 5. | RM24 | 1 | 58.2 | GAAGTGTGATCACTGTAA | TACAGTGGACGGCGAAGTCG | 18.97 | GA | 29 |
| 6. | RM9 | 1 | 58.2 | GGTGCCATTGTCGTCCTC | ACGGCCCTCATCACCTTC | 23.32 | GA, GT, GA | 15, 1, 2 |
| 7. | RM543 | 1 | 59.3 | CTGCTGCAGACTCTACTGC | AAATATTACCCATCCCCC | 32.78 | GCG | 10 |
| 8. | RM7180 | 1 | 48.3 | GTGTTTATAGGGGTGCCAC | TGTTGGTGGTGCAGGTAAG | 34.1 | ATAG | 6 |
| 9. | RM1339 | 1 | 56.3 | ATCAAAGCATGTAAACCA | CGTAAGATCTCCCTACCAC | 38.19 | AG | 22 |
| 10. | RM14 | 1 | 55.7 | CCGAGGAGAGGAGTTCGAC | GTGCCAATTCCTCGAAA | 41.36 | GA | 18 |
| 11. | RM279 | 2 | 57.3 | GCGGGAGAGGGATCTCCT | GGCTAGGAGTTAACCTCGCG | 2.88 | GA | 16 |
| 12. | RM71 | 2 | 58.3 | CTAGAGGCGAAAACGAGATG | GGGTGGGCGAGGTAATAATG | 8.76 | ATT, T, ATT | 10, 1, 4 |
| 13. | RM452 | 2 | 58.3 | CGATCGAGAGCGTTAAGG | GGGATCAAACCACGTTTCTG | 9.56 | GTC | 9 |
| 14. | RM324 | 2 | 57.3 | CTGATTCCACACACTTGTGC | GATTCCACGTCAGGATCTTC | 11.38 | CAT | 21 |
| 15. | RM263 | 2 | 60.4 | CCCAGGCTAGCTCATGAA | GCTACGTTTGAGCTACCACG | 25.86 | CT | 34 |
| 16. | RM526 | 2 | 57.5 | CCCAAGCAATACGTCCCTAG | ACCTGGTCATGACAAGGAGG | 26.66 | TAAT | 5 |
| 17. | RM14220 | 2 | 61.5 | TCTCCACACAAACTTGGAACACG | GTCTGTATTGTGGGTGCAAGAGG | 35.67 | AC | 14 |
| 18. | CHR3_14 | 3 | 52.6 | GCCTACTCGTCTACCAACA | GGCACTTATCCATTTCCAG | 4.97 | AT | n/a |
| 19. | CHR3_20 | 3 | 55.3 | ATTGACACGAAGAGGACA | GTGCCCGAGGTGAGTGAAGT | 5.95 | GTA | n/a |
| 20. | RM36 | 3 | 58.3 | CAACTATGCACCATTGTGCG | GTACTCCACAAGACCGTACC | 7.12 | GA | 23 |
| 21. | RM16 | 3 | 57.4 | CGCTAGGGCAGCATCTAA | AACACAGCAGGTACGCGC | 23.12 | TCG, GA | 5, 16 |
| 22. | RM3513 | 3 | 61.5 | TACTCCTATCCTGCCATGG | TGTAGTAGACGAGAGGCCCGG | 25.11 | CT | 28 |
| 23. | RM4796 | 3 | 57.4 | CCACGGTAGTTTTGGTCTAC | AGAGGGGAAGAGTGAGAGAG | 27.47 | TA | 26 |
| 24. | RM520 | 3 | 55.3 | AGGAGCAAGAAAAGTTCC | GCCAATGTGTGACGCAATAG | 30.91 | AG | 10 |

| | | | | | | | | |
|-----|------------|---|------|------------------------------|--------------------------------|-------|------------------------|-------------------|
| 25. | RM51 4 | 3 | 56.3 | AGATTGATCTCCCATTCCC C | CACGAGCATATTACTAG TGG | 35.28 | AC | 12 |
| 26. | RM33 5 | 4 | 59.8 | GTACACACCCACATCGAG AAG | GCTCTATGCGAGTATCC ATGG | 0.68 | CTT | 25 |
| 27. | RM26 1 | 4 | 58.3 | CTACTTCTCCCCTTGTGTC G | TGTACCATCGCCAAATC TCC | 6.57 | C, CT | 9, 8 |
| 28. | RM11 42 | 4 | 57.1 | AAGCACACGTAAAACGGA GG | CGTCACTCTCACCACCA CC | 21.39 | AG | 12 |
| 29. | RM13 88 | 4 | 54 | TTCAATGAGGCAAAGGTA AG | ATTGTAGCTTGGACTAG GGG | 25.2 | AG | 46 |
| 30. | RM31 7 | 4 | 59.8 | CATACTTACCAGTTCACCG CC | CTGGAGAGTGTGAGCTA GTTGA | 29.06 | GC, GT | 4, 18 |
| 31. | RM67 48 | 4 | 52.7 | ATTGGGTTTCTCATATTAT G | CCAACACTCCTAACTAG TTC | 30.9 | TAT | 18 |
| 32. | RM38 43 | 4 | 61.5 | ACCCTACTCCCAACAGTCC C | GGGGTCGTACGCTCATG TC | 31.49 | GA | 23 |
| 33. | RM11 13 | 4 | 55.3 | GGGCGCATGTGTATTTCTT C | TGGGGAAAAACCACAA GCC | 34.08 | AG | 12 |
| 34. | RM28 0 | 4 | 57.4 | ACACGATCCACTTTGCGC | TGTGTCTTGAGCAGCCA GG | 34.98 | GA | 16 |
| 35. | RM12 2 | 5 | 61.5 | GAGTCGATGTAATGTCATC AGTGC | GAAGGAGGTATCGCTTT GTTGGAC | 0.31 | GA, A, GA, A, GA | 7, 1, 2, 1, 11 |
| 36. | RM13 | 5 | 56.6 | TCCAACATGGCAAGAGAG AG | GGTGGCATTTCGATTCCA G | 2.01 | GA | 16 |
| 37. | RM54 8 | 5 | 56.4 | TCGGTGAGAACTGAGAG TACG | AAGGAGGCCATCTCAAT GTG | 2.81 | CT | 12 |
| 38. | RM26 7 | 5 | 56.8 | TGCAGACATAGAGAAGGA ATG | AGCAACAGCACAACTTG ATG | 2.88 | GA | 21 |
| 39. | RM28 9 | 5 | 56.6 | TTCCATGGCACACAAGCC | CTGTGCACGAACTTCCA AAG | 7.8 | G, GA | 11, 16 |
| 40. | RM43 0 | 5 | 55 | AAACAACGACGTCCCTGA TC | GTGCCTCCGTGGTTATG AAC | 18.69 | GA | 25 |
| 41. | RM16 3 | 5 | 60.4 | ATCCATGTGCGCCTTTATG AGGA | CGCTACCTCCTTCACTTA CTAGT | 19.18 | GGAGA, GA, C, GA | 4, 11, 1, 20 |
| 42. | RM36 16 | 5 | 56.3 | GTGCGGATTTCTCCTCTCT C | TGCCGGTCCATTTCTAG AAG | 26.28 | GA | 13 |
| 43. | SSI | 6 | 57.4 | GATCCGTTTTTGCTGTGCC C | CCTCCTCTCCGCCGATC CTG | 3.09 | n/a | |
| 44. | RM21 7 | 6 | 54.4 | ATCGCAGCAATGCCTCGT | GGGTGTGAACAAAGAC AC | 4.23 | CT | 20 |
| 45. | RM68 36 | 6 | 53.9 | TGTTGCATATGGTGCTATT TGA | GATACGGCTTCTAGGCC AAA | 9.3 | TCT | 14 |
| 46. | RM56 4 | 6 | 54.3 | TCCTTCTGCATCAATTCTC CTCG | CTCCATAGCCTTGTTAA GTGATGAGC | 11.61 | GT | 14 |
| 47. | RM31 83 | 6 | 56.6 | GCTCCACAGAAAAGCAAA GC | TGCAACAGTAGCTGTAG CCG | 12.44 | CT | 12 |
| 48. | RM13 40 | 6 | 56.3 | TCCAACTAGTGGGAACG C | CTCAACGCCATGAACCT C | 23.34 | AG | 22 |
| 49. | RM52 8 | 6 | 57.3 | GGCATCCAATTTTACCCCT C | AAATGGAGCATGGAGGT CAC | 26.55 | AGAT | 9 |
| 50. | RM30 | 6 | 59.8 | GGTTAGGCATCGTCACGG | TCACCTCACCACACGAC ACG | 26.87 | AG, A, GA | 9, 1, 12 |
| 51. | RM69 65 | 7 | 53.2 | TCATTTGGATCATAAGCTG G | TTGGATGAGATAACCAA TGC | 2.8 | TTC | 11 |

| | | | | | | | | |
|-----|--------------|----|------|------------------------------|-----------------------------|-------|---------------------|--------------|
| 52. | RM48 1 | 7 | 57.3 | TAGCTAGCCGATTGAATG GC | CTCCACCTCCTATGTTGT TG | 2.88 | CAA | 12 |
| 53. | RM54 2 | 7 | 55 | TGAATCAAGCCCCTCACTA C | CTGCAACGAGTAAGGCA GAG | 12.71 | CT | 22 |
| 54. | RM43 2 | 7 | 55.5 | TTCTGTCTCACGCTGGATT G | AGCTGCGTACGTGATGA ATG | 18.95 | CATC | 9 |
| 55. | RM35 1 | 7 | 58.2 | CCATCCTCCACCGCCTCTC G | TGGAGGAAGGAAAGGG GACG | 23.92 | CCG, CGAAG | 9, 4 |
| 56. | RM11 32 | 7 | 61.5 | ATCACCTGAGAAACATCC GG | CTCCTCCCACGTCAAGG TC | 23.98 | AG | 12 |
| 57. | RM23 4 | 7 | 55 | ACAGTATCCAAGGCCCTG G | CACGTGAGACAAAGAC GGAG | 25.47 | CT | 25 |
| 58. | RM35 55 | 7 | 57.4 | TGGAAGTTTCTGGCGATA G | TGGTTGGACTGAAAAGT CCC | 27.89 | GA | 12 |
| 59. | RM40 8 | 8 | 61.5 | CAACGAGCTAACTTCCGTC C | ACTGCTACTTGGGTAGC TGACC | 0.12 | CT | 13 |
| 60. | RM68 63 | 8 | 54.4 | GCTGCAGAATTAAGGAGA AC | TGCTCAAAATAATCAGC TCC | 2.01 | TGC | 9 |
| 61. | RM56 47 | 8 | 58 | ACTCCGACTGCAGTTTTTG C | AACTTGGTCGTGGACAG TGC | 2.89 | AAG | 16 |
| 62. | RM54 7 | 8 | 58.5 | TAGGTTGGCAGACCTTTTC G | GTCAAGATCATCCTCGT AGCG | 5.59 | ATT | 19 |
| 63. | CHR8 _4 | 8 | 56.9 | GATTGAAAGAGAAAGGTG GTT | CTGTGTAACCGAGTTAC GTTT | 9.95 | ATAG | n/a |
| 64. | RM40 4 | 8 | 57.3 | CCAATCATTAACCCCTGAG C | GCCTTCATGCTTCAGAA GAC | 15.43 | GA | 33 |
| 65. | RM42 | 8 | 58.3 | ATCCTACCGCTGACCATGA G | TTTGGTCTACGTGGCGT ACA | 20.09 | AG, AG, T, GA | 6-2, 1, 5 |
| 66. | ARO7 | 8 | 54.4 | ATTTGCCTCCTGAGTCTG | GAGGATGGGGAAGATA AA | 20.26 | n/a | |
| 67. | BADE X7-5 | 8 | 53.1 | TGTTTTCTGTTAGGTTGCA TT | ATCCACAGAAATTTGGA AAC | 20.32 | n/a | |
| 68. | nksbad 2 | 8 | 61 | GGTTGCATTTACTGGGAGT TATG | TCCACAGAAATTTGGAA ACAAAC | 20.38 | n/a | |
| 69. | RM80 | 8 | 57.7 | TTGAAGGCGCTGAAGGAG | CATCAACCTCGTCTTCA CCG | 24.47 | TCT | 25 |
| 70. | RM31 6 | 9 | 56.7 | CTAGTTGGGCATACGATG GC | ACGTTATATGTTACGT CAAC | 1.1 | GT, TG, TTTG, TG | 8-9, 4, 4 |
| 71. | RM56 88 | 9 | 56.3 | GCAGTGTCCAACCATCTGT G | ATCTGGTCACCCTTTGCT TG | 1.71 | AAT | 17 |
| 72. | RM21 9 | 9 | 54.6 | CGTCGGATGATGTAAAGC CT | CATATCGGCATTCGCCT G | 7.9 | CT | 17 |
| 73. | RM73 87 | 9 | 54.6 | GCAGTAGGGAGCATGGAA AG | AAACGAGTCCTCTTCAG GGG | 10.13 | GAGT | 6 |
| 74. | RM10 5 | 9 | 67.7 | GTCGTGACCCATCGGAG CCAC | TGGTTCGAGGTGGGGATC GGGTC | 12.6 | CCT | 6 |
| 75. | RM32 8 | 9 | 57.5 | CATAGTGGAGTATGCAGC TGC | CCTTCTCCCAGTCGTATC TG | 19.72 | CAT | 5 |
| 76. | RM16 0 | 9 | 66.7 | CCCAAATCAGGAAAGTTT CTCAGC | AGTCATCCTTGGCTACC AGATGC | 19.78 | GAA | 23 |
| 77. | RM38 08 | 9 | 55.6 | CGTTAGCGAAACGAACAG TG | CAGTGGCTCGGTAATCG C | 20.54 | GA | 20 |
| 78. | RM47 4 | 10 | 55.3 | AAGATGTACGGGTGGCAT TC | TATGAGCTGGTGAGCAA TGG | 1.81 | AT | 13 |

| | | | | | | | | |
|-----|---------|----|------|---------------------|-----------------------------|-------|------------------|----------------|
| 79. | RM311 | 10 | 50.7 | TGGTAGTATAGGTAACAT | TCCTATACACATACAAA CATAAC | 9.74 | GT, GTAT, GT | 3, 8, 5 |
| 80. | RM258 | 10 | 58 | TGCTGTATGTAGCTCGCAC | TGGCCTTTAAAGCTGTC GC | 18.01 | GA, GGA | 21, 3 |
| 81. | RM171 | 10 | 58.7 | AACGCGAGGACACGTACT | ACGAGATACGTACGCCT TTG | 19.04 | GATG | 5 |
| 82. | RM590 | 10 | 58.5 | CATCTCCGCTCTCCATGC | GGAGTTGGGGTCTTGTT CG | 23.04 | TCT | 10 |
| 83. | RM286 | 11 | 56.9 | GGCTTCATCTTTGGCGAC | CCGGATTCACGAGATAA ACTC | 0.38 | GA | 16 |
| 84. | RM202 | 11 | 55.6 | CAGATTGGAGATGAAGTC | CCAGCAAGCATGTCAAT GTA | 9 | CT | 30 |
| 85. | RM229 | 11 | 57.4 | CACTCACACGAACGACTG | CGCAGGTTCTTGTA TGT | 18.4 | TC, CT, C, CT | 11, 5, 3, 5 |
| 86. | RM21 | 11 | 60.4 | ACAGTATTCCGTAGGCAC | GCTCCATGAGGGTGGTA GAG | 19.6 | GA | 18 |
| 87. | RM206 | 11 | 56.3 | CCCATGCGTTAACTATTC | CGTTCCATCGATCCGTA TGG | 22.01 | CT | 21 |
| 88. | RM1233 | 11 | 53.2 | TTCGTTTCCTTGGTTAGT | ATTGGCTCCTGAAGAAG G | 26.53 | AG | 15 |
| 89. | RM5926 | 11 | 54.4 | ATATACTGTAGGTCCATCC | AGATAGTATAGCGTAGC AGC | 28.43 | ATT | 21 |
| 90. | RM20 | 12 | 57.8 | ATCTTGTCCTGCAGGTCA | GAAACAGAGGCACATTT CATTG | 0.97 | ATT | 14 |
| 91. | RM19 | 12 | 54.4 | CAAAAACAGAGCAGATGA | CTCAAGATGGACGCCAA GA | 2.43 | ATC | 10 |
| 92. | RM1337 | 12 | 55.3 | GTGCAATGCTGAGGAGTA | CTGAGAATCTGGAGTGC TTG | 11.93 | AG | 21 |
| 93. | RM28305 | 12 | 56.5 | GTCATCTTCGCAAATGGTG | GGTCGTCGTGGTGTAT TCTTGG | 19.92 | GA | 31 |
| 94. | RM28367 | 12 | 62 | CGTATCTCCACCTCCCGAG | GCCAAATCTCACGGATC GAAGC | 21.17 | AG | 21 |
| 95. | RM28556 | 12 | 61.1 | CTAGTAGTGCCACTTAACC | GGATCCAAACACCACCT TAGCC | 23.98 | TC | 14 |
| 96. | RM17 | 12 | 59.3 | TGCCCTGTTATTTCTTCTC | GGTGATCCTTCCCATTT CA | 26.95 | GA | 21 |

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge.

REFERENCES:

1. Amarawathi Y, Singh R, Singh AK, Singh VP, Mohapatra T, Sharma TR, & Singh NK. Mapping of quantitative trait loci for basmati quality traits in rice (*Oryza sativa* L.). *Molecular Breeding*. 2008, 21(1): 49-65.
2. Anonymous, 2021. FAOSTAT Crop Statistics 2019. FAO of UN, International Fertilizer Industry Association. Available from <http://www.fao.org/faostat/en/#data/QC>. Accessed on 20th October, 2021.

3. Bao JS, Corke H, Sun M. Microsatellites, single nucleotide polymorphisms and a sequence tagged site in starch-synthesizing genes in relation to starch physicochemical properties in nonwaxy rice (*Oryza sativa* L.). *Theoretical and Applied Genetics*. 2006;113(7):1185-96.
4. Bao J. Genes and QTLs for rice grain quality improvement. *InTech–Open Science Open Mind*. 2014:239-78.
5. Bazrkar-Khatibani L, Fakheri BA, Hosseini-Chaleshtori M, Mahender A, Mahdinejad N, Ali J. Genetic mapping and validation of quantitative trait loci (QTL) for the grain appearance and quality traits in rice (*Oryza sativa* L.) by using recombinant inbred line (RIL) population. *International journal of genomics*. 2019.
6. Calingacion M, Laborte A, Nelson A, Resurreccion A, Concepcion JC, Daygon VD, Mumm R, Reinke R, Dipti S, Bassinello PZ, Manful J. Diversity of global rice markets and the science required for consumer-targeted rice breeding. *PloS one*. 2014;9(1):e85106.
7. Chandu G, Addanki KR, Balakrishnan D, Mangrauthia SK, Sudhakar P, Satya AK, Neelamraju S. SSR markers for grain iron zinc and yield-related traits polymorphic between Samba Mahsuri (BPT5204) and a wild rice *Oryza rufipogon*. *Electronic Journal of Plant Breeding*. 2020;11(03):841-7.
8. Cheng A, Ismail I, Osman M, Hashim H. Mapping of quantitative trait loci for aroma, amylose content and cooked grain elongation traits in rice. *Plant Omics*. 2014;7(3).
9. Collard BC, Jahufer MZ, Brouwer JB, Pang EC. An introduction to markers, quantitative trait loci (QTL) mapping and marker-assisted selection for crop improvement: the basic concepts. *Euphytica*. 2005;142(1):169-96.
10. Dai L, Wang L, Leng Y, Yang Y, Huang L, Chen L, Wang Y, Ren D, Hu J, Zhang G, Zhu L. Quantitative Trait Loci Mapping for Appearance Quality in Short- Grain Rice. *Crop Science*. 2016;56(4):1484-92.
11. Fitzgerald MA, McCouch SR, Hall RD. Not just a grain of rice: the quest for quality. *Trends in plant science*. 2009;14(3):133-9.
12. Forrester JV, Dick AD, McMenamin PG, Roberts F, Pearlman E. *The eye e-book: basic sciences in practice*. Elsevier Health Sciences; 2020.
13. Hable S, Singh SK, Mounika K, Khaire A, Singh DK, Majhi PK. Study of allelic variation at genome wide SSR loci in parents of mapping population for high grain zinc in rice (*Oryza sativa* L.). *Journal of Experimental Biology and Agricultural Sciences*. 2020; 8(5):558-575.
14. Kulkarni SR, Balachandran SM, Ulaganathan K, Balakrishnan D, Praveen M, Prasad AH, Fiyaz RA, Senguttuvel P, Sinha P, Kale RR, Rekha G. Molecular mapping of QTLs for yield related traits in recombinant inbred line (RIL) population derived from the popular rice hybrid KRH-2 and their validation through SNP genotyping. *Scientific reports*. 2020;10(1):1-21.
15. Mason AS. Challenges of genotyping polyploid species. *Plant Genotyping*. 2015:161-8.
16. Murray MG, Thompson WF. Rapid isolation of high molecular weight plant DNA. *Nucleic acids research*. 1980;8(19):4321-6.
17. Narshimulu G, Jamaloddin M, Vemireddy LR, Anuradha G, Siddiq E. Potentiality of evenly distributed hypervariable microsatellite markers in marker-assisted breeding of rice. *Plant Breeding*. 2011;130(3):314-20.
18. Nelson JC, McClung AM, Fjellstrom RG, Moldenhauer KA, Boza E, Jodari F, Oard JH, Linscombe S, Scheffler BE, Yeater KM. Mapping QTL main and interaction influences on milling quality in elite US rice germplasm. *Theoretical and Applied Genetics*. 2011;122(2):291-309.
19. Nicot N, Chiquet V, Gandon B, Amilhat L, Legeai F, Leroy P, Bernard M, Sourdille P. Study of simple sequence repeat (SSR) markers from wheat expressed sequence tags (ESTs). *Theoretical and Applied Genetics*. 2004;109(4):800-5.
20. Pradhan SK, Pandit E, Pawar S, Naveenkumar R, Barik SR, Mohanty SP, Nayak DK, Ghritlahre SK, Rao DS, Reddy JN, Patnaik SS. Linkage disequilibrium mapping for grain Fe and Zn enhancing QTLs useful for nutrient dense rice breeding. *BMC plant biology*. 2020;20(1):1-24.

21. Sakthivel K, Rani NS, Pandey MK, Sivaranjani AK, Neeraja CN, Balachandran SM, Madhav MS, Viraktamath BC, Prasad GS, Sundaram RM. Development of a simple functional marker for fragrance in rice and its validation in Indian Basmati and non-Basmati fragrant rice varieties. *Molecular breeding*. 2009;24(2):185-90.
22. Shabir G, Aslam K, Khan AR, Shahid M, Manzoor H, Noreen S, Khan MA, Baber M, Sabar M, Shah SM, Arif M. Rice molecular markers and genetic mapping: Current status and prospects. *Journal of integrative agriculture*. 2017;16(9):1879-91.
23. Shivani D, Suman K, Madhubabu P, Rathod R, Cheralu C, Shankar VG, Neeraja CN. Parental polymorphism in iron-and zinc-rich rice varieties (Swarna and type 3) Using ssr markers. *Applied Biological Research*. 2020; 22(1): 69-75.
24. Shrivastav SP, Verma OP, Singh V, Lal K. Interrelationships among yield and its contributing traits in rice (*Oryza sativa* L.) under sodic soil. *Electronic Journal of Plant Breeding*. 2020;11(04):1044-52.
25. Singh R, Singh AK, Sharma TR, Singh A, Singh NK. Fine mapping of aroma QTLs in basmati rice (*Oryza sativa* L) on chromosomes 3, 4 and 8. *Journal of plant biochemistry and biotechnology*. 2007;16(2):75-82.
26. Sun SX, Gao FY, Lu XJ, Wu XJ, Wang XD, Ren GJ, Luo H. Genetic analysis and gene fine mapping of aroma in rice (*Oryza sativa* L. Cyperales, Poaceae). *Genetics and Molecular Biology*. 2008; 31:532-8.
27. Swamy BM, Kaladhar K, Shobha Rani N, Prasad GSV, Viraktamath BC, Reddy GA, Sarla N. QTL analysis for grain quality traits in 2 BC2F2 populations derived from crosses between *Oryza sativa* cv Swarna and 2 accessions of *O. nivara*. *Journal of Heredity*. 2012; 103(3): 442-452.
28. Van Berloo, R. GGT: user manual Version 2.0. Wageningen (The Netherlands): Wageningen University. 2007.
29. Van Berloo R. GGT 2.0: versatile software for visualization and analysis of genetic data. *Journal of Heredity*. 2008;99(2):232-6.
30. Vemireddy LR, Noor S, Satyavathi VV, Srividhya A, Kaliappan A, Parimala SR, Bharathi PM, Deborah DA, Rao KS, Shobharani N, Siddiq EA. Discovery and mapping of genomic regions governing economically important traits of Basmati rice. *BMC plant biology*. 2015;15(1):1-9.
31. Xia D, Zhou H, Qiu L, Jiang H, Zhang Q, Gao G, He Y. Mapping and verification of grain shape QTLs based on an advanced backcross population in rice. *Plos one*. 2017;12(11): e0187553.
32. Yano M, Sasaki T. Genetic and molecular dissection of quantitative traits in rice. *Oryza: from molecule to plant*. 1997:145-53.