

# INFLUENCE OF BIODEGRADABLE POLYMER COATED UREA ON NITROGEN UPTAKE AND UTILIZATION OF MAIZE (*Zea mays* L.)

## Abstract

Controlled release nitrogen fertilizers could be an excellent management approach for improving nitrogen fertilizer efficiency. The present study aimed to investigate the effect of coated urea fertilizers to increase nitrogen uptake and utilization of maize. The nitrogen use efficiency of maize from various biodegradable polymer-coated urea fertilizers, such as palm stearin coated urea (PSCU), pine oleoresin coated urea (POCU), and humic acid coated urea (HACU), was determined in a pot culture experiment conducted at the Department of Soil Science and Agricultural Chemistry, Tamil Nadu Agricultural University, Coimbatore, during 2021. The coating materials have been coated on urea with different coating thicknesses, viz., PSCU - 5, 10, 15%, POCU – 2, 4, 6%, and HACU - 5, 10, 15%. Among all the treatments, T<sub>11</sub>: HACU 15% produced highest grain yield (72.0g plant<sup>-1</sup>) followed by T<sub>7</sub>: POCU 4% (69.7 g plant<sup>-1</sup>) and T<sub>4</sub>: PSCU 10% (69.0g plant<sup>-1</sup>). In terms of dry matter production, T<sub>10</sub>: PSCU 10% produced maximum dry matter (186.5g plant<sup>-1</sup>), followed by T<sub>11</sub>: HACU 15% (186.2 g plant<sup>-1</sup>), and T<sub>7</sub>: POCU 4% (185.3g plant<sup>-1</sup>). The nitrogen uptake by the maize plant was higher in T<sub>7</sub>: POCU 4 % (1.62g plant<sup>-1</sup>), followed by T<sub>11</sub>: HACU 15% (1.59 g plant<sup>-1</sup>) and T<sub>4</sub>: PSCU 10% (1.59g plant<sup>-1</sup>). Irrespective of treatments, the highest nitrogen utilization by the maize crop was found in T<sub>7</sub>: POCU 4% (73.9%) followed by T<sub>4</sub>: PSCU 10% (71.1%) and T<sub>11</sub>: HACU 15% (70.9%) treatments. When compared to uncoated urea fertilizer, all coated urea fertilizers outperformed uncoated urea fertilizer in terms of grain yield, dry matter accumulation, and nitrogen uptake. To improve the nitrogen use efficiency, coated urea fertilizers prove to be a promising alternative to uncoated urea fertilizers.

**Keywords:** *urea, coated fertilizers, nitrogen release, nitrogen utilization, nitrogen uptake*

## 1. Introduction

Maize (*Zea mays* L.) is one of the most widely cultivated cereal crops in the world, providing grain, silage, and biofuel (Bono *et al.*, 2020). Nitrogen (N) is a chemical element that has required by most cultures of economic interest, and it has been the most impact on maize output (Fageria and Baligar, 2005; Marini *et al.*, 2015; Arnuti *et al.*, 2017). Nitrogen is

insufficient in nearly all agricultural soils and cropping systems around the world. It is necessary to use prudent nitrogen management to grow crops to meet the ever-increasing demands of human populations (Mohan *et al.*, 2015).

For any agricultural production system, nitrogen is the most important externally provided ingredient. The food supply of half of the world's population is reliant on nitrogenous fertilizers, either directly or indirectly. Rice, wheat, and maize now consume more than 90% of all nitrogenous fertilizers applied to cereals. Underuse of nitrogen is linked to reduced agricultural yields, whereas overuse has a variety of soil and environmental effects. As a result, the response to applied nitrogen and its effectiveness must have been carefully evaluated to achieve maximum potential and sustainable yield (Yadav *et al.*, 2017).

The consumption of reactive nitrogen (Nr) in the form of fertilizer in India has increased quickly over the last six decades, with an annual rise of 6% since 1970 (Sutton *et al.*, 2017 based on FAO, 2016), but the cropping system of nitrogen use efficiency (NUE) has decreased (Moring *et al.*, 2021). This increased fertilizer use has led to increasing Nr losses, one of the clearest signs of which is the yearly emission of ammonia (NH<sub>3</sub>), the most common fertilizer related atmospheric Nr loss, which is on the rise (EDGAR, 2016).

When urea is applied to plants in bare form it is prone to losses due to nitrous emissions, leaching, denitrification, and surface runoff (Azeem *et al.*, 2014; Trinh *et al.*, 2015). Naz *et al.* (2016) reported, this loss leads to economic loss, low plant nutrient use efficiency, and environmental damage due to water eutrophication and nitrous emissions into the stratosphere.

Controlled release urea can be used to prevent granular urea loss due to leaching, ammonia volatilization, and denitrification (Azeem *et al.*, 2016). The thickness of the coating film is important for better-controlled release qualities (Timilsena *et al.*, 2015). The time it takes for a dissolved nutrient to diffuse out of the coated shell and into the bulk water it is immersed in is determined by the diffusional path it must travel (Ito *et al.*, 2003).

The main goal of fertilizer research is to increase the efficiency and availability of fertilizers to plants, either by increasing the availability of existing fertilizers or by producing a new high-efficiency fertilizer. The discovery of controlled-release fertilizers (CRFs) boosted fertilizer efficiency by allowing fertilizer to be released over a longer period. CRFs are made by lowering the solubility and mobility of conventional fertilizers through physical or

chemical modification (Trenkel, 1997). The present study investigates the low-cost biodegradable polymer-coated urea to enhance the nitrogen use efficiency of maize.

## 2. Material and methods

### 2.1. Collection and preparation of soil sample


Soil samples have collected from Field No. 37F of the eastern block, Tamil Nadu Agricultural University, Coimbatore. The experimental soil is a mixed black calcareous, fine, montmorillonite, isohyperthermic which belonged to the Inceptisol order, Periyanaickenpalayam soil series, and classified as Vertic Ustropept taxonomically. The soil was air-dried, processed by quartering method, then sieved with a 2mm sieve. Representative sub examples were dispossessed for further soil physico-chemical analysis by embracing standard analytical procedures are given in **Table 1**.

**Table 1. Physico-chemical characteristics of experiment soil**

Soil parameters (Units)	Values	Methods
pH (1:2.5 soil/water ratio)	8.17	Jackson (1973)
Electrical conductivity (dS m <sup>-1</sup> ) (1:2.5 soil/water ratio)	0.24	Jackson (1973)
Organic carbon (%)	0.62	Walkley and Black (1934)
Available nitrogen (kg ha <sup>-1</sup> )	179.2	Subbiah and Asija (1956)
Available phosphorus (kg ha <sup>-1</sup> )	16.7	Olsen (1954)
Available potassium (kg ha <sup>-1</sup> )	504.1	Stanford and English (1949)
Bulk density (Mg m <sup>-3</sup> )	1.17	Gupta and Dakshinamurthi (1981)
Particle density (Mg m <sup>-3</sup> )	2.45	Gupta and Dakshinamurthi (1981)
CEC (cmol (P <sup>+</sup> ) kg <sup>-1</sup> )	45.6	Jackson (1973)
Zn (mg kg <sup>-1</sup> )	0.56	Lindsay and Norvell (1978)
Fe (mg kg <sup>-1</sup> )	2.05	Lindsay and Norvell (1978)
Cu (mg kg <sup>-1</sup> )	0.68	Lindsay and Norvell (1978)
Mn (mg kg <sup>-1</sup> )	5.97	Lindsay and Norvell (1978)
Textural class	Sandy clay loam	Piper (1966)

### 2.2. Synthesis and characterization of coated urea fertilizers

Three types of coated urea fertilizers were made with various biodegradable polymer coating materials in different thicknesses of coating, such as palm stearin coated urea 5, 10, and 15%, pine oleoresin coated urea 2, 4, and 6%, and humic acid coated urea 5, 10, and 15%, under laboratory conditions using a rotating drum. The coating thickness between urea and coated materials was examined under a scanning electron microscope (SEM) (**Fig. 1**) at the Department of Nano Science and Technology laboratory, TNAU, Coimbatore.



**Figure 1.** Scanning electron micrograph of (a) PSCU 5%, (b) PSCU 10%, (c) PSCU 15%, (d) POCU 2%, (e) POCU 4%, (f) POCU 6%, (g) HACU 5%, (h) HACU 10%, (i) HACU 15% at x1000 magnification.

About 10kg of soil was weighed and filled in each mud pot. After that, maize (COH (M) 6) hybrid seeds were sown. There were eleven treatments and replicated thrice under completely randomized design (CRD). The treatments details of the experiment have given in **Table 2**. The recommended dose of fertilizer ( $1120 \text{ mg N pot}^{-1}$ ) was applied as three splits (25% basal, 50% at 25 DAS, 25% at 45 DAS), the entire dose of Single super Phosphate (SSP) and Muriate of Potash (MOP) applied as basal application. Plant samples were taken at the vegetative, tasseling, and harvest stages of the plant and determined for nitrogen uptake and maize N use efficiency.

**Table 2. Treatment details of the experiment**

<b>Treatments</b>	<b>Abbreviations</b>
T <sub>1</sub> - Control	Control
T <sub>2</sub> - Uncoated urea	UCU
T <sub>3</sub> - Palm stearin coated urea 5%	PSCU 5%
T <sub>4</sub> - Palm stearin coated urea 10%	PSCU 10%
T <sub>5</sub> - Palm stearin coated urea 15%	PSCU 15%
T <sub>6</sub> - Pine oleoresin coated urea 2%	POCU 2%
T <sub>7</sub> - Pine oleoresin coated urea 4%	POCU 4%
T <sub>8</sub> - Pine oleoresin coated urea 6%	POCU 6%
T <sub>9</sub> - Humic acid coated urea 5%	HACU 5%
T <sub>10</sub> - Humic acid coated urea 10%	HACU 10%
T <sub>11</sub> - Humic acid coated urea 15%	HACU 15%

### **Nitrogen uptake and Fertilizer N utilization calculation**

$$N \text{ uptake (g pot}^{-1}\text{)} = \frac{N \text{ concentration in plant (\%)} \times \text{weight of dry matter (g)}}{100}$$

$$\text{Fertilizer N utilization (\%)} = \frac{\text{Fertilizer N uptake of plant (mg)}}{\text{Fertilizer N applied (mg)}} \times 100$$

### **2.3. Statistical analysis**

Using AGRES statistical package, the data from the experiments were subjected to analysis of variance (ANOVA) to determine significance. Significant critical differences (CD) were calculated at the 5% confidence level wherever treatment differences were found.

## **3. Results and discussion**

### **3.1. Dry matter production**

From the vegetative to the harvest stage, maize crop dry matter production increased (**Table 3**). Maize dry matter accumulation under different N treatments revealed a typical sigmoidal tendency across the maize growing season, according to Guo *et al.* (2021). The dry matter production of coated urea and uncoated urea differed significantly in the vegetative stage, ranging from 18.7-37.2g plant<sup>-1</sup>. T<sub>2</sub>: Uncoated urea treatment (37.2g plant<sup>-1</sup>) had the

maximum dry matter production, while T<sub>1</sub>: control treatment had the lowest dry matter output (18.7g plant<sup>-1</sup>). Uncoated urea produced more dry matter than any of the coated urea fertilizers at this stage. Probably, this is a result of uncoated urea's rapid dissolution, which encourages faster crop absorption than coated urea fertilizers at this early stage. Slow-release fertilizers may release too slowly from the polymer coating to be a helpful N supply in the early phases of crop development (Farmaha and Sims, 2013). According to Mullen (2011), nitrogen in the soil is lost by volatilization, immobilization, denitrification, and leaching prior to crop uptake, with the volume and pathways of loss affected by environmental conditions such as soil moisture and temperature. The medium thickness coated urea fertilizers, such as T<sub>4</sub>: PSCU 10% (30.8g plant<sup>-1</sup>), T<sub>7</sub>: POCU 4% (26.4g plant<sup>-1</sup>), and T<sub>10</sub>: HACU 10% (32.3g plant<sup>-1</sup>), and the high thickness coated urea fertilizers, such as T<sub>5</sub>: PSCU 15% (28.4g plant<sup>-1</sup>), T<sub>8</sub>: POCU 6% (25.7g plant<sup>-1</sup>), and T<sub>11</sub>: HACU 15% (29.9g plant<sup>-1</sup>), produced lesser dry matter production than T<sub>2</sub>: uncoated urea (37.2g plant<sup>-1</sup>) treatment during the vegetative stage because higher percentage coated urea takes longer to dissolve than lower percentage coated urea. Blends of slow and controlled-release urea, according to Andrade *et al.* (2021), provide better regulated N release to maize plants, giving adequate quantities of N at each growth stage or throughout time than traditional urea.

In tasselling stage, the dry matter production was ranged from 48.6–85.3g plant<sup>-1</sup>. The highest dry matter production was recorded in T<sub>7</sub>: POCU 4% treatment (85.3g plant<sup>-1</sup>) which is on par with T<sub>4</sub>: PSCU 10%, T<sub>8</sub>: POCU 6% and T<sub>11</sub>: HACU 15% and the lowest dry matter production was recorded in T<sub>1</sub>: control treatment (48.6g plant<sup>-1</sup>). When compared to coated urea fertilizer treatments, the T<sub>2</sub>: uncoated urea treatment produced less dry matter (59.8g plant<sup>-1</sup>). According to Sofia *et al.* (2019), the vegetative biomass of plants was treated by amended polymer composite fertilizers tend to be higher than positive and negative controls. This is since the maize plants' nitrogen requirements were met by the coated fertilizers. Because nitrogen is released slowly from coated urea fertilizers due to the degradation of coating material as time passes, medium thickness coated urea fertilizers such as T<sub>4</sub>: PSCU 10% (82.4 g plant<sup>-1</sup>), T<sub>7</sub>: POCU 4% (85.3 g plant<sup>-1</sup>), and T<sub>10</sub>: HACU 10% (74.9 g plant<sup>-1</sup>) produced more dry matter. The extent of yield improvement with coated fertilizers, according to Guan *et al.* (2014), was largely dependent on the timely release of nutrients coated inside the three layers of fertilizer. Attapulgit coating reduced nutrient loss in the early growing stage as a barrier to chemical fertilizer, allowing maize plants to acquire nutrients in the middle and late developing phases.

In harvest stage, the minimum and maximum stover yields were produced from T<sub>1</sub>: control and T<sub>4</sub>: PSCU 10% treatments, respectively, it was ranged from 88.6-110.8g plant<sup>-1</sup>. Palm stearin coated urea outperformed other coated urea fertilizers but, on par with all the treatments except control. Palm stearin is a wax containing coating material that is hydrophobic in nature, meaning it won't dissolve easily in water. As a result of the gradual release of nitrogen from PSCU, greater stover yield might have produced. According to Mathialagan *et al.* (2020), palm stearin acts as a physical barrier between urea granules and water, preventing rapid urea disintegration. In comparison to uncoated urea, the application of palm stearin coated urea boosted maize dry matter yield from 60 to 20% pot<sup>-1</sup> and increased N uptake by 77%, according to Nasima *et al.* (2010). In terms of grain yield, T<sub>11</sub>: HACU 15% treatment had the highest yield (72.0g plant<sup>-1</sup>) which is on par with T<sub>7</sub>: POCU 4% (69.7g plant<sup>-1</sup>) and T<sub>4</sub>: PSCU 10% (69.0g plant<sup>-1</sup>), the T<sub>1</sub>: control treatment had the lowest yield (32.9 g plant<sup>-1</sup>). This could be because humic-acid has a multitude of acidic functional groups, a great of specific surface area, high cation exchange capacity, and a lot of absorption capacity. According to Chen *et al.* (2017), using humic-acid urea fertiliser improved sweet potato yield by 29.56 % while also increasing biological yield and harvest index. Total dry matter production from coated urea fertilizers differed significantly from uncoated urea fertilizers and it was ranging from 127.5-186.5g plant<sup>-1</sup>. T<sub>4</sub>: PSCU 10% treatment had the maximum dry matter accumulation (186.5g plant<sup>-1</sup>), while T<sub>1</sub>: control treatment had the lowest dry matter production (127.5g plant<sup>-1</sup>). T<sub>2</sub>: uncoated urea produced less dry matter production (152.3g plant<sup>-1</sup>) compared to all the coated urea fertilizer treatments. This could be because the nitrogen supplied from the coated urea meets the crop need whenever it is needed, resulting in higher dry matter output. Grant *et al.* (2012) and Zhang *et al.* (2019) found that the delayed release fertilizer's nitrogen release rate was synced with plant N absorption, resulting in fewer nitrogen losses. The nitrogen release properties of controlled release urea were also synchronised with the N requirements of wheat and maize crops during their whole growth cycles, according to Zheng *et al.* (2017). Among all coated urea fertilizers, the thickest coated urea produced less dry matter. This could be because the nitrogen released from the coated urea was exceedingly sluggish and did not match the crop's requirements. De Almeida *et al.* (2019) found that using urea blends did not increase maize yields, and Garcia *et al.* (2019) found that coated urea did not always match the crop demand, affecting dry mass and nitrogen uptake of crop.

**Table 3. Effect of coated urea fertilizers on dry matter production (g plant<sup>-1</sup>) of maize**

Treatments	Dry matter production (g plant <sup>-1</sup> )				
	Vegetative	Tasselling	Harvest		
			Stover	Grain	Total DMP
T <sub>1</sub> -Control	18.7±0.72 <sup>f</sup>	48.6±1.15 <sup>g</sup>	88.6±0.53 <sup>b</sup>	32.9±0.64 <sup>f</sup>	127.5±6.89 <sup>d</sup>
T <sub>2</sub> -UCU	37.2±1.61 <sup>a</sup>	59.8±1.24 <sup>f</sup>	101.4±1.79 <sup>a</sup>	46.5±0.77 <sup>e</sup>	152.3±8.24 <sup>c</sup>
T <sub>3</sub> -PSCU 5%	33.2±0.10 <sup>b</sup>	63.7±2.59 <sup>ef</sup>	106.8±2.50 <sup>a</sup>	57.5±2.57 <sup>d</sup>	169.7±4.95 <sup>b</sup>
T <sub>4</sub> -PSCU 10%	30.8±0.64 <sup>bcd</sup>	82.4±3.86 <sup>ab</sup>	110.8±6.17 <sup>a</sup>	69.0±0.18 <sup>ab</sup>	186.5±7.96 <sup>a</sup>
T <sub>5</sub> -PSCU 15%	28.4±1.45 <sup>de</sup>	76.5±1.67 <sup>bcd</sup>	108.4±3.67 <sup>a</sup>	61.4±1.05 <sup>cd</sup>	175.4±0.37 <sup>ab</sup>
T <sub>6</sub> -POCU 2%	32.6±0.41 <sup>bc</sup>	66.2±1.34 <sup>ef</sup>	106.5±1.55 <sup>a</sup>	57.5±1.86 <sup>d</sup>	169.6±4.85 <sup>b</sup>
T <sub>7</sub> -POCU 4%	26.4±0.89 <sup>e</sup>	85.3±4.39 <sup>a</sup>	108.1±0.28 <sup>a</sup>	69.7±0.94 <sup>ab</sup>	185.3±6.17 <sup>a</sup>
T <sub>8</sub> -POCU 6%	25.7±1.35 <sup>e</sup>	80.6±1.85 <sup>abc</sup>	107.4±0.95 <sup>a</sup>	63.8±2.89 <sup>bc</sup>	177.8±1.85 <sup>ab</sup>
T <sub>9</sub> -HACU 5%	30.5±0.87 <sup>bcd</sup>	70.3±2.49 <sup>de</sup>	107.7±3.92 <sup>a</sup>	60.1±0.16 <sup>cd</sup>	173.4±3.25 <sup>ab</sup>
T <sub>10</sub> -HACU 10%	32.3±1.50 <sup>bc</sup>	74.9±2.14 <sup>cd</sup>	108.3±2.31 <sup>a</sup>	59.5±3.28 <sup>cd</sup>	174.5±4.18 <sup>ab</sup>
T <sub>11</sub> -HACU 15%	29.9±1.07 <sup>cd</sup>	82.2±1.63 <sup>ab</sup>	109.5±5.64 <sup>a</sup>	72.0±3.82 <sup>a</sup>	186.2±3.78 <sup>a</sup>
<b>Mean</b>	<b>29.6</b>	<b>71.9</b>	<b>105.8</b>	<b>59.1</b>	<b>170.7</b>
<b>SEd</b>	<b>1.51</b>	<b>3.44</b>	<b>4.62</b>	<b>2.92</b>	<b>7.52</b>

Values are mean of three replicates ± standard error (n=3); values followed by the same letter in each column are not significantly different from each other.

### 3.2. Nitrogen uptake

Nitrogen uptake of maize from coated urea fertilizers significantly differed from uncoated urea fertilizer treatment (Table 4). Applied N exhibited much larger uptakes of N than the typical urea plot, according to Xie *et al.* (2019). In vegetative stage, nitrogen uptake ranged from 0.180-0.625g plant<sup>-1</sup>, with T<sub>1</sub>: control and T<sub>2</sub>: uncoated urea treatments recording the lowest and highest uptake, respectively. Because nutrient uptake was extremely quick at this stage, uncoated urea nitrogen uptake was at its peak. More thickness coated urea had the lowest nitrogen uptake, while less thickness coated urea had the highest nitrogen uptake. This could be because the thickness of the coating determines the amount of urea released beyond the coated layer. Nutrient release from coated fertilizer may be influenced by coating thickness, according to Shaviv *et al.* (2003). It is vital to remember that the higher the proportion of coated layer content, the lower the nitrogen leakage (Behin and Sadeghi, 2016).

In tasselling stage, T<sub>1</sub>: control and T<sub>7</sub>: POCU 4% treatments had the lowest and maximum nitrogen uptake, respectively, ranging from 0.403-1.467g plant<sup>-1</sup>. When compared to all the coated urea fertilizer treatments, uncoated urea treatments produced less nitrogen uptake. This could be because uncoated urea released nitrogen early and caused losses such as ammonia volatilization and nitrate leaching, preventing plants from absorbing nitrogen from the fertilizer. Coated urea fertilizers, on the other hand, postpone nitrogen release, allowing the plant to absorb more nitrogen during this stage. Andrade *et al.* (2021) reported, Controlled-release urea and their blends have a delayed N release and promote continuous N uptake after 70 days. Likewise, the applied N rates were also more important than the N fertilizer technologies after 70 days. To help synchronise N release and uptake by maize, an appropriate blend of N fertilizers could be an attractive choice.

In harvest stage, stover nitrogen uptake ranged from 0.470-0.835g plant<sup>-1</sup>, with the lowest and maximum uptakes detected in the T<sub>1</sub>: control and T<sub>5</sub>: PSCU 15% treatments, respectively. When compared to all coated urea fertilizers, the greatest thickness coated urea T<sub>11</sub>: PSCU 15% registered greater nitrogen uptake in stover, which could be owing to the delayed nitrogen release up until this harvest stage, which induced more nitrogen uptake. The lowest and highest grain nitrogen uptake were obtained in the T<sub>1</sub>: control and T<sub>11</sub>: HACU 15% treatments, respectively, which varied from 0.323-0.885g plant<sup>-1</sup>. The absorbed nitrogen from coated urea fertilizers was gladly supplied to the grains at the right period in the crop cycle. According to Noor Affendi, *et al.* (2018), the plots fertilised with Urea coated with Cu and Zn (UCuZn) and UCuZn + nitrification inhibitor had the maximum exported N in stover and grain. Under uncoated urea, grain maize had the lowest exported nitrogen.

The maize crop's total nitrogen uptake ranged from 0.79-1.62g plant<sup>-1</sup>, with the lowest and maximum uptakes recorded in the T<sub>1</sub>: control and T<sub>7</sub>: POCU 4% treatments, respectively. When compared to coated urea fertilizers, T<sub>2</sub>: uncoated urea had lower total nitrogen uptake (1.13g plant<sup>-1</sup>). This could be attributed to higher nitrogen losses from uncoated urea and reduced nitrogen losses from coated urea fertilizers. When compared to regular urea, Ning *et al.* (2012) found that controlled release urea can boost total nitrogen uptake by aboveground organs and soil residual while lowering loss. Coated urea fertilizers may reduce nutrient losses by enhancing nutrient utilization efficiency. Control release fertilizers can be used to meet crop nutrient requirements in a single or divided application while also reducing pollution in the environment (Noor Affendi, *et al.*, 2018). We might deduce from these

findings that the coated fertilizers' release pattern satisfied the crop demand and provided more nitrogen to the plants.

**Table 4. Effect of coated urea fertilizers on nitrogen uptake ( $\text{g plant}^{-1}$ ) of maize**


Treatments	N uptake ( $\text{g plant}^{-1}$ )				Total uptake
	Vegetative	Tasselling	Harvest		
			Stover	Grain	
T <sub>1</sub> -Control	0.180±0.003 <sup>g</sup>	0.403±0.015 <sup>h</sup>	0.470±0.010 <sup>f</sup>	0.323±0.001 <sup>f</sup>	0.79±0.020 <sup>f</sup>
T <sub>2</sub> -UCU	0.625±0.002 <sup>a</sup>	0.712±0.037 <sup>g</sup>	0.608±0.005 <sup>e</sup>	0.521±0.004 <sup>e</sup>	1.13±0.011 <sup>e</sup>
T <sub>3</sub> -PSCU 5%	0.485±0.007 <sup>b</sup>	0.841±0.036 <sup>f</sup>	0.641±0.018 <sup>de</sup>	0.667±0.012 <sup>cd</sup>	1.31±0.015 <sup>cd</sup>
T <sub>4</sub> -PSCU 10%	0.431±0.002 <sup>d</sup>	1.343±0.003 <sup>b</sup>	0.753±0.030 <sup>bc</sup>	0.835±0.018 <sup>a</sup>	1.59±0.078 <sup>a</sup>
T <sub>5</sub> -PSCU 15%	0.375±0.002 <sup>ef</sup>	1.125±0.054 <sup>d</sup>	0.835±0.009 <sup>a</sup>	0.700±0.007 <sup>bcd</sup>	1.53±0.044 <sup>ab</sup>
T <sub>6</sub> -POCU 2%	0.505±0.008 <sup>b</sup>	0.907±0.011 <sup>ef</sup>	0.655±0.005 <sup>de</sup>	0.644±0.022 <sup>d</sup>	1.30±0.012 <sup>d</sup>
T <sub>7</sub> -POCU 4%	0.391±0.005 <sup>e</sup>	1.467±0.012 <sup>a</sup>	0.762±0.042 <sup>bc</sup>	0.858±0.031 <sup>a</sup>	1.62±0.024 <sup>a</sup>
T <sub>8</sub> -POCU 6%	0.357±0.017 <sup>f</sup>	1.225±0.012 <sup>c</sup>	0.773±0.011 <sup>ab</sup>	0.734±0.017 <sup>b</sup>	1.51±0.078 <sup>ab</sup>
T <sub>9</sub> -HACU 5%	0.439±0.015 <sup>bd</sup>	0.942±0.017 <sup>e</sup>	0.592±0.033 <sup>e</sup>	0.697±0.036 <sup>bcd</sup>	1.29±0.029 <sup>d</sup>
T <sub>10</sub> -HACU 10%	0.478±0.023 <sup>c</sup>	1.071±0.019 <sup>d</sup>	0.699±0.001 <sup>cd</sup>	0.720±0.015 <sup>bc</sup>	1.42±0.019 <sup>bc</sup>
T <sub>11</sub> -HACU 15%	0.449±0.011 <sup>cd</sup>	1.299±0.026 <sup>bc</sup>	0.701±0.034 <sup>cd</sup>	0.885±0.039 <sup>a</sup>	1.59±0.031 <sup>a</sup>
<b>Mean</b>	<b>0.429</b>	<b>1.030</b>	<b>0.681</b>	<b>0.689</b>	<b>1.37</b>
<b>SEd</b>	<b>0.015</b>	<b>0.037</b>	<b>0.032</b>	<b>0.031</b>	<b>0.057</b>

Values are mean of three replicates ± standard error (n=3); values followed by the same letter in each column are not significantly different from each other.

### 3.3. Fertilizer Nitrogen utilization

Fertilizer N utilization or nitrogen recovery efficiency (NRE) from the applied fertilizers was calculated based on the nitrogen uptake in treated and control pots of the maize. Fertilizer N utilization was maximum in coated urea fertilizers than uncoated urea fertilizers (**Fig. 2**). The highest fertilizer utilization has recorded in T<sub>7</sub>: POCU 4% (73.9%), which is atpar with T<sub>4</sub>: PSCU 10% (71.1%), and T<sub>11</sub>: HACU 15% (70.9%), while the lower N utilization efficiency has registered in T<sub>2</sub>: Uncoated urea treatment (30.3%). This was due to the slow release of

nitrogen from the coated urea fertilizers to the longer period. Fertilizer N utilization was highest in urea applied with urease inhibitor than in urea applied without urease inhibitor, according to Gans *et al.* (2006). Fertilizer N utilization was 56% in his study, compared to 42.4% for conventional urea. According to Guo *et al.* (2021), the maize crop's nitrogen recovery efficiency ranged between 20.4 and 40.2%, with the maximum NRE recorded in urea blended with slow-release nitrogen fertilizer at a ratio of 3:7 and the lowest NRE recorded with uncoated urea fertilizer treatment. According to Andrade *et al.* (2021), due to lower N availability from N sources throughout time, declines in the NRE in uncoated urea are low throughout the cropping periods.



**Figure 2. Effect of coated urea fertilizers on fertilizer N utilization (%) of maize**

#### 4. Conclusion

Coated urea fertilizers releases nitrogen according to crop demand, has the potential to enhance nitrogen use efficiency while significantly reducing losses due to volatilization, leaching, and denitrification. According to the result of this study, coated urea fertilizer is a good alternative for uncoated urea fertilizers to enhance the nitrogen use efficiency of maize. The impacts of PSCU, POCU, and HACU on maize productivity were clearly proven in this study by releasing nitrogen over a long period of time that corresponded to the nitrogen

requirement at different stages of crop growth. However, further field investigations are necessary to validate the use of this fertilizer on nitrogen use efficiency.

### Acknowledgment

The authors are grateful to the Department of Soil Science and Agricultural Chemistry at Tamil Nadu Agricultural University for providing all of the necessary resources to carry out this study. Also, thanks to the Department of Nano Science and Technology for assisting with the instrumentation work.

**ORCID:** <https://orcid.org/0000-0002-1772-4326>

### References

1. Andrade AB, Guelfi DR, Chagas WF, Cancellier EL, de Souza TL, Oliveira LS, Faquin V, Du C. Fertilizing maize croppings with blends of slow/controlled-release and conventional nitrogen fertilizers. *Journal of Plant Nutrition and Soil Science*. 2021; 184(2):227-37. <https://doi.org/10.1002/jpln.201900609>
- 2.
3. Arnuti F, Cecagno D, Martins AP, Balerini F, Meurer EJ, da Silva PR. Irrigation intensity and management of nitrogen topdressing in corn in a consolidated no-tillage system. *In Colloquium Agrariae*. 2017; 13, 29-40.
4. Azeem B, KuShaari K, Man Z. Effect of coating thickness on release characteristics of controlled release urea produced in fluidized bed using waterborne starch biopolymer as coating material. *Procedia engineering*. 2016; 148:282-289. <https://doi.org/10.1016/j.proeng.2016.06.615>
5. Azeem B, KuShaari K, Man ZB, Basit A, Thanh TH. Review on materials & methods to produce controlled release coated urea fertilizer. *Journal of controlled release*. 2014; 181:11-21. <https://doi.org/10.1016/j.jconrel.2014.02.020>
6. Behin J, Sadeghi N. Utilization of waste lignin to prepare controlled-slow release urea. *International Journal of Recycling of Organic Waste in Agriculture*. 2016; 5(4):289-299. <https://doi.org/10.1007/s40093-016-0139-1>
7. Bono JA, dos SANTOS HW, Pereira SR, dos Reis Neto JF. Nitrogen coated fertilizer with controlled release for the maize crop. *Bioscience Journal*. 2020; 23;36(6). <http://dx.doi.org/10.14393/BJ-v36n6a2020-47704>
8. EDGAR - Emissions Database for Global Atmospheric Research v4.3.1. 2016.
9. Fageria NK, Baligar VC. Enhancing nitrogen use efficiency in crop plants. *Advances in agronomy*. 2005; 88:97-185. [https://doi.org/10.1016/S0065-2113\(05\)88004-6](https://doi.org/10.1016/S0065-2113(05)88004-6)
10. FAO. United Nations Food and Agriculture Organization, Statistics. 2016. Available online at: <http://www.fao.org/faostat/>.

11. Farmaha BS, Sims AL. The influence of polymer-coated urea and urea fertilizer mixtures on spring wheat protein concentrations and economic returns. *Agronomy Journal*. 2013; 105(5):1328-1334. <https://doi.org/10.2134/agronj2012.0454>
12. Garcia PL, Sermarini RA, Trivelin PC. Effect of nitrogen rates applying controlled-release and conventional urea blend in maize. *Journal of Plant Nutrition*. 2019; 42(18):2199-2208. <https://doi.org/10.1080/01904167.2019.1658778>
13. Grant CA, Wu R, Selles F, Harker KN, Clayton GW, Bittman S, Zebarth BJ, Lupwayi NZ. Crop yield and nitrogen concentration with controlled release urea and split applications of nitrogen as compared to non-coated urea applied at seeding. *Field Crops Research*. 2012; 127:170-180. <https://doi.org/10.1016/j.fcr.2011.11.002>
14. Guan Y, Song C, Gan Y, Li FM. Increased maize yield using slow-release attapulgate-coated fertilizers. *Agronomy for sustainable development*. 2014; 34(3):657-665. <https://doi.org/10.1007/s13593-013-0193-2>
15. Guo J, Fan J, Zhang F, Yan S, Zheng J, Wu Y, Li J, Wang Y, Sun X, Liu X, Xiang Y. Blending urea and slow-release nitrogen fertilizer increases dryland maize yield and nitrogen use efficiency while mitigating ammonia volatilization. *Science of The Total Environment*. 2021; 790. <https://doi.org/10.1016/j.scitotenv.2021.148058>
16. Gupta RP, Dakshinamurthi C. Procedures for physical Analysis of soils. IARI, New Delhi 1981.
17. Ito R, Golman B, Shinohara K. Controlled release with coating layer of permeable particles. *Journal of controlled release*. 2003; 92(3):361-368 [https://doi.org/10.1016/S0168-3659\(03\)00363-8](https://doi.org/10.1016/S0168-3659(03)00363-8)
18. Jackson M. Soil Chemical Analysis. Prentic Hall (India) Pvt Ltd New Delhi 1973.
19. Lindsay, Willard L, Norvell W Aa. Development of a DTPA soil test for zinc, iron, manganese, and copper. *Soil Science Society of America Journal*. 1978; 42(3):421-428. <https://doi.org/10.2136/sssaj1978.03615995004200030009x>
20. Marini D, Guimarães VF, Dartora J, Lana MD, Pinto AS. Growth and yield of corn hybrids in response to association with *Azospirillum brasilense* and nitrogen fertilization1. *Revista Ceres*. 2015; 62:117-123. <https://doi.org/10.1590/0034-737X201562010015>
21. Mohan S, Singh M, Kumar R. Effect of nitrogen, phosphorus and zinc fertilization on yield and quality of kharif fodder-A review. *Agricultural Reviews*. 2015; 36(3):1.218-226. <http://dx.doi.org/10.5958/0976-0741.2015.00025.2>
22. Moring A, Hooda S, Raghuram N, Adhya TK, Ahmad A, Bandyopadhyay SK, Barsby T, Beig G, Bentley A, Bhatia A, Dragosits U. Nitrogen challenges and opportunities for agricultural and environmental science in India. *Frontiers in Sustainable Food Systems*. 2021; 5:505347. <https://doi.org/10.3389/fsufs.2021.505347>
23. Mullen RW. Nutrient cycling in soils: Nitrogen. *Soil management: Building a stable base for agriculture*. 2011; 25:67-78. <https://doi.org/10.2136/2011.soilmanagement.c5>
24. Naz MY, Sulaiman SA. Slow-release coating remedy for nitrogen loss from conventional urea: a review. *Journal of Controlled Release*. 2016; 225:109-200. <https://doi.org/10.1016/j.jconrel.2016.01.037>
25. Ning T, Shao GQ, Li ZJ, Han HF, Hu HG, Wang Y, Tian SZ. Effects of urea types and irrigation on crop uptake, soil residual, and loss of nitrogen in maize field on the North China Plain. *Plant, Soil and Environment*. 2012; 58(1):1-8.

26. Noor Affendi NM, Yusop MK, Othman R. Efficiency of coated urea on nutrient uptake and maize production. *Communications in Soil Science and Plant Analysis*. 2018; 49(11):1394-1400. <https://doi.org/10.1080/00103624.2018.1464182>
27. Olsen SR. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. US Department of Agriculture. 1954; 9398:1-19.
28. Piper CS. "Soil and plant analysis, Hans. Pub. Bombay 1966, Asian Ed:368-374.
29. Shaviv A, Raban S, Zaidel E. Modeling controlled nutrient release from polymer coated fertilizers: Diffusion release from single granules. *Environmental science & technology*. 2003; 37(10):2251-2256. <https://doi.org/10.1021/es011462v>
30. Sofia AR, Hala Y, Makkulawu AT, Hiola SF, Karim H, Iriany RN, Sjahril R, Jumadi O. Influence of urea fertilizer applied with polyacrylate polymer, zeolite and Mimba on growth maize. *InIOP Conference Series: Earth and Environmental Science*. 2019; 299 (1):012017). IOP Publishing.
31. Stanford George, Leah English. Use of the flame photometer in rapid soil tests for K and Ca. *Agronomy Journal*. 1949; 41(9):446-447. <https://doi.org/10.2134/agronj1949.00021962004100090012x>
32. Subbiah BV, Asija GL. A rapid method for the estimation of nitrogen in soil. *Current Science*. 1956; 26:259-260.
33. Sutton MA, Drewer J, Moring A, Adhya TK, Ahmed A, Bhatia A, Brownlie W, Dragosits U, Ghude SD, Hillier J, Hooda S. The Indian nitrogen challenge in a global perspective. *InThe Indian Nitrogen Assessment*. 2017; 1:9-28. <https://doi.org/10.1016/B978-0-12-811836-8.00002-1>
34. Timilsena YP, Adhikari R, Casey P, Muster T, Gill H, Adhikari B. Enhanced efficiency fertilizers: a review of formulation and nutrient release patterns. *Journal of the Science of Food and Agriculture*. 2015; 95(6):1131-1142. <https://doi.org/10.1002/jsfa.6812>
35. Trenkel ME. *Controlled-release and stabilized fertilizers in agriculture*. Paris: International fertilizer industry association; 1997.
36. Trinh TH, Kushaari K, Shuib AS, Ismail L, Azeem B. Modelling the release of nitrogen from controlled release fertilizer: Constant and decay release. *Biosystems Engineering*. 2015; 130:34-42. <https://doi.org/10.1016/j.biosystemseng.2014.12.004>
37. Walkley Adous, Black IA. An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil science*. 1934; 37(1):29-38.
38. Xie Y, Tang L, Han Y, Yang L, Xie G, Peng J, Tian C, Zhou X, Liu Q, Rong X, Zhang Y. Reduction in nitrogen fertilizer applications by the use of polymer-coated urea: effect on maize yields and environmental impacts of nitrogen losses. *Journal of the Science of Food and Agriculture*. 2019; 99(5):2259-2266. <https://doi.org/10.1002/jsfa.9421>
39. Yadav MR, Kumar R, Parihar CM, Yadav RK, Jat SL, Ram H, Meena RK, Singh M, Verma AP, Kumar U, Ghosh A. Strategies for improving nitrogen use efficiency: A review. *Agricultural Reviews*. 2017; 38(1):29-40. <http://dx.doi.org/10.18805/ag.v0iOF.7306>
40. Zhang W, Liang Z, He X, Wang X, Shi X, Zou C, Chen X. The effects of controlled release urea on maize productivity and reactive nitrogen losses: a meta-analysis.

Environmental Pollution. 2019; 246:559-565.  
<https://doi.org/10.1016/j.envpol.2018.12.059>

41. Zheng W, Liu Z, Zhang M, Shi Y, Zhu Q, Sun Y, Zhou H, Li C, Yang Y, Geng J. Improving crop yields, nitrogen use efficiencies, and profits by using mixtures of coated controlled-released and uncoated urea in a wheat-maize system. *Field Crops Research*. 2017; 205:106-115. <https://doi.org/10.1016/j.fcr.2017.02.009>

Chen X, Kou M, Tang Z, Zhang A, Li H, Wei M. Responses of root physiological characteristics and yield of sweet potato to humic acid urea fertilizer. *Plos one*. 2017; 12(12):e0189715. <https://doi.org/10.1371/journal.pone.0189715>

Mathialagan R, Mansor N, Shamsuddin MR, Noor Affendi NM, Hamid Nour A. Performance of allicin coated with palm stearin on hydrolyzation of urea applied on soil. *Journal of Plant Nutrition*. 2020; 44(10):1446-1457. <https://doi.org/10.1080/01904167.2020.1862187>

Nasima J, Khanif MY, Hanfi MM, Wan ZW, Dharejo KA. Maize response to biodegradable polymer and urease inhibitor coated urea. *International Journal of Agriculture and Biology*. 2010; 12(5):773-776.

de Almeida RE, Favarin JL, Oliveira FB, Pierozan Junior C, de Oliveira SM, Lago BC, Tezotto T, Otto R, Trivelin PC. Polymer-coated urea in broadcast or furrow application in the corn-palisadegrass intercropping system. *Embrapa Pesca e Aquicultura-Artigo em periodico indexado (ALICE)*. 2019.

Gans W, Herbst F, Merbach W. Nitrogen balance in the system plant-soil after urea fertilization combined with urease inhibitors. *Plant Soil Environment*. 2006; 52(special issue):36-38.