

Analytical Groundwater Quality Assessment for Drinking and Agriculture Purposes in Al-Jouf Region, Kingdom of Saudi Arabia

Ahmed Nazal Al-Shemary*, Khalid Dawi Al-Otabi and Abdulrasoul Mosa Al-Omran

Department of Soil Science, College of Food and Agricultural Sciences, King Saud University, P.O. Box 2460 Riyadh, 11451 Saudi Arabia

ABSTRACT

Kingdom of Saudi Arabia (KSA) primarily relies depends mainly on groundwater for drinking and irrigation purposes. The study was therefore aimed to identify groundwater quality in Al-Jouf Region, KSA using water quality index (DWQI). In addition, investigating the hydro-chemical characteristics that control the groundwater quality. Groundwater samples were collected from 150 groundwater wells at a 300-500 m depth and subjected for chemical analysis. The values of chemical constituents were compared with the KSA and World Health Organization (WHO) standards for drinking and irrigation purposes. The results indicated that, the concentrations of ions were within the ranges of KSA for drinking water and WHO. Based on DWQI data, for drinking water about 23.9 % of the wells were within poor water category (III), while 9.91 % was very poor water within (IV) group, 45.6 % is good water of group (II) and 20.5 % is excellent water within category (I). Regarding the evaluation of water quality, the estimated DWQI values for the 150 well waters in Al-Jouf, region ranged from 40.7 to 319. About 23.9% of wells were considered poor water "class (III)", 9.9% were very poor water "class (IV)", 45.6% were good water for drinking or "class (II)", and 20.53% were excellent water. The result shows that the groundwater possess moderate to high salinity hazards with low to medium sodium hazards. The piper diagram showed that cations were decreasing as follow: $Na^+ > Ca_2^+ > Mg_2^+$, while the anions were decreasing as follow $Cl^- > HCO_3^- > SO_4^{2-} > CO_3^{2-}$. The SAR values varied from 0.68 to 15.43; while Kelly's ratio (KR) ranged between 0.32 to 4.02. The calculated IWQI values of all wells revealed that water was moderate type in which its value was between 22 to -27.

Keywords: Al-Jouf, groundwater, water quality index, salinity hazard.

1. INTRODUCTION

Across many countries, groundwater quality is considered as important source for drinking water and irrigation activities, particularly, in the semi-arid and arid regions. Resultantly, it is estimated that groundwater is the main source for drinking [1]. Kingdom of Saudi Arabia (KSA) is in arid regions, suffers from limited water resources [2]. Exploitation of groundwater led to serious shortage of water and the deterioration of groundwater led to reduction of agricultural land in different regions of KSA [3]. In the rural area of Saudi Arabia such as Al-Jouf region is rural area in KSA, rely mainly on groundwater for drinking and agricultural activities. Al-Jouf area is considered one of the new agricultural regions in KSA with very high potential in agricultural development recently. During the last three decades the region

29 witnessed a huge agricultural project from most of agricultural companies. This stress on
30 agricultural water demand is the main cause of water resources deterioration [2]. In Al-Jouf
31 area, groundwater is currently limiting factor for intensifying the agricultural activities. The
32 irrigation water quality may affect crop production and soil chemical and physical properties
33 [3]. Thus, there is lack of information on irrigation water quality in Al-Jouf area to make
34 necessary management decisions in crop production. Salinity and sodicity, and ion toxicity of
35 the water are very important issues. Sodidity or the presence of too much Na^+ , which causes
36 the poor soil structure [4]. Various factors such as rock-water interaction, lithology, usage of
37 fertilizers and pesticides for agricultural purposes, and climatic conditions largely influence
38 the water quality [5].

39 The drinking water quality index (DWQI) and irrigation water quality index (IWQI) are
40 functions to access water quality and help to take the right decision for the policy makers in
41 reassuring the public and farmers on their water quality [6,7]. The aim of both DWQI and
42 IWQI parameters is to provide a simplified approach for evaluating drinking and irrigation
43 water purposes [8,9]. Numerous studies were conducted using DWQI and IWQI with
44 different methods of calculation of the index and the weight values for each parameter
45 [10,11]. On DWQI, Al-Othman [12] reported that DWQI of ground water in Riyadh region,
46 KSA ranged from 34 to 513 with an average value of 282. In Iran [13] used DWQI to
47 evaluate groundwater quality and their result proved that water is of good quality. Similar
48 results were reported in El-Khairat, Tunisia [14] and in Palakkal District in Kerala, India [15-
49 18]. The proposed index utilizes five hazard groups namely salinity toxicity, infiltration rate,
50 specific ion and heavy metals toxicity [19]. Based on this technique, results indicated that
51 ground water quality in the western Anatolia, Turkey are fairly good and aquifers are mostly
52 suitable for irrigation [17] applied the same procedure in south-central Bangladesh. They
53 reported that groundwater using (IWQI) is moderate to suitable for irrigation. This study
54 aimed to investigate the groundwater quality assessment for drinking and agriculture
55 purposes in Al-Jouf Region, KSA.

56

57

58 **2. MATERIAL AND METHODS**

59

60 **2.1 Study Area Description**

61 Water samples were collected from different agricultural sites in Al-Jouf aera, north of KSA
62 (Fig. 1). Al-Jouf region has an area of 100,212 km^2 , population of 508,475 and the city of
63 Sakaka is the administrative headquarters for the region, which includes three governorates
64 of Al-Qurayyat, Dumat Al-Jandal and Tabarjal.

65 Al-Jouf aera was considered one of the most fertile soils in the KSA, and its famous for olive
66 tree cultivation. Al-Jouf area produces approximately 67% of the domestic production of
67 olive oil in the Kingdom. Also, the cultivation of palm tree, is about 150 thousand tons of old
68 dates additionally, fruits, vegetables and wheat [21]. Rainfall varies greatly among seasons
69 which has an annual average of 50-60 mm. The soils in this area consist of sandy plains of
70 different canals and sandy plains of marshes with the presence of sedimentary plains topped
71 by sandy layers, a sedimentary joint and a lower slope valley, coastal plains, wet coastal
72 sand, and there are rocky plains in some places [22].

73

74




Fig. 1. Location of the study area and sampling

75
76
77
78
79
80

2.2 Chemical Analysis

81 Groundwater samples were collected from different 150 different wells located in Al-Jouf
82 area to estimate groundwater quality. The samples were analyzed for electrical conductivity
83 (EC), pH, cations, anions and B. The EC was measured at 25 °C by using EC- meter in and
84 pH was determined using a pH meter (pH meter—CG 817). While the Ca_2^+ and Mg_2^+
85 was determined by titration method; whereas the Na^+ and K^+ concentration was measured using
86 flame photometer (Corning 400) [34-14]. The HCO_3^- and SO_4^{2-} concentration was
87 determined by titration methos [23], while NO_3^- concentration was determined by the
88 phenoldisulfonic acid method [25] and B was measured using azomethine-H method [26].

89

2.3 Accuracy of Collected Data Ion Balance Errors

91 The ion balance errors were used to check accuracy in the analytical procedures using the
92 following formula (1) as described by American Public Health Association [27]:

93

94

95

96

$$\text{Ion balance} = \frac{\sum \text{cations} - \sum \text{anions}}{\sum \text{cations} + \sum \text{anions}} \times 100 \quad (1)$$

97

98

99
100
101
102
103
104
105
106
107
108
109

2.4 Water Quality Index Calculation

The DWQI calculations include three successive steps were used [6,14,28,29] as indicated below:

DWQI calculations, 12 parameters have been used including weight (w_i) as shown in (Table 1). The most important parameters have a weight of (5) and the lower scale (1). The results indicated that, the maximum weight of (5) has been assigned for NO_3^- , due to its major importance in water quality assessment [30], the minimum weight (2) has been used for Ca_2^+ , Mg_2^+ , and Na^+ .

Table 1. Parameters Relative weight

Chemical parameter	Weights (w_i)	Relative weight (W_i)
pH	3	0.0938
TDS (mg L^{-1})	5	0.1563
Ca_2^+ (mg L^{-1})	2	0.0625
Mg_2^+ (mg L^{-1})	2	0.0625
Na^+ (mg L^{-1})	2	0.0625
K^+ (mg L^{-1})	3	0.0938
HCO_3^- (mg L^{-1})	3	0.0938
Cl^- (mg L^{-1})	3	0.0938
SO_4^{2-} (mg L^{-1})	3	0.0938
NO_3^- (mg L^{-1})	3	0.0938
B (mg L^{-1})	3	0.0938
Total	32	1

110
111
112
113
114
115
116
117

Second the relative weight (W_i) is calculated as equation (2):

$$W_i = \frac{w_i}{\sum_{i=1}^n w_i} \quad (2)$$

Where W_i = the relative weight, w_i is the weight of each parameter and n = the number of parameters. The calculated W_i values of each parameter are shown in (Table 1).

Table 2. Parameters standards for WHO

Parameters	Standards
pH	6.50 –8.50
Hardness	500
TDS (mg L^{-1})	600
* Ca_2^+	75
Mg_2^+	50
Na^+	200

K ⁺	12
HCO ₃ ⁻	120
Cl ⁻	250
SO ₄ ²⁻	250
NO ₃ ⁻	10
B	0.50

118 * (mg/L)

119

120 Third is the quality index (qi) for each parameter as:

121

$$122 \quad q_i = \frac{C_i}{S_i} \times 100 \quad (3)$$

123

124 where qi is the quality index, Ci is the concentration of each chemical parameter mg/L,
125 except for pH, and Si is the WHO standard value, Wi and qi is used to calculate the Sli for
126 each chemical parameter as shown in equations (4) and (5):

$$127 \quad S_{Li} = W_i \times q_i \quad (4)$$

128

$$129 \quad WQI = \sum_{i=1}^n S_{Li} \quad (5)$$

130

131 Sli = the sub parameter; qi = the concentration and n = the parameter number. The
132 calculated WQI values are presented in (Table 3).

133

Table 3. Water quality classification according to WQI

WQI Range	Index	Type of water
<50.5	I	Suitable
50.5–100.1	II	Good
100.1–200.1	III	Poor
200.1–300.1	IV	Very poor
>300.1	V	unsuitable

134

135 2.5 Hydro chemical Characterization

136

137

138

Table 4. Classification of Irrigation water

Potential irrigation problem	Unit	Degree of restriction use			
		None	Slight to moderate	Severe	
Salinity	EC	μS/cm	<750	750–3,000	>3000
	TDS	mg/L	<450	450–2,000	>2000
Permeability	SAR=0–3	and EC=	>750	750–200	<200
	SAR=3–6		>1300	1,300–300	<300
	SAR=6–12		>1800	1,800–500	<500
	SAR=12–20		>2900	2,900–1,300	<1300
	SAR=20–40		>5000	5,000–2,900	<2900
Specific ion toxicity	Na+	SAR	<3	3.00–9.00	>9
	Cl-	mg/L	<140	140–350	>350
	B	mg/L	<0.7	0.70–3.00	>3

*Surface irrigation.

139 The hydro chemical characterization of the untreated groundwater samples was evaluated
140 using Ca^{2+} , Mg^{2+} , Na^+ , K^+ , HCO_3^- , Cl^- , and SO_4^{2-} concentrations. The piper, Schoeller, and
141 Durov diagrams were drawn using Geochemistry Software Aq.QA, version AQC10664, in
142 addition US salinity laboratory [32] diagrams were also used. The following data, salinity
143 hazard, sodium adsorption ratio (SAR), total hardness (as CaCO_3), and Kelly's ratio (KR)
144 were calculated to assessment of the groundwater for irrigation purposes.

145

146 2.6 Irrigation Water Quality Index (IWQI)

147 The IWQI is calculated using equations from (6) to (11) as proposed by [9,16].

$$148 \text{IWQ Index} = \sum_{i=1}^5 G_i \quad (6)$$

149

150 where is an incremental index and G the contribution of the 5 hazard groups. The first
151 category is the salinity hazard of water is estimated using:

152

$$153 G_1 = w_1 r_1 \quad (7)$$

154

155 w = the weight value of this hazard group and r = the parameter index value of the as are
 156 shown in (Table 5).

157

158 Table (4) represents the groundwater quality according to Ayers [19]. It includes various
 159 potential irrigation problems such as salinity on crop water availability, permeability and
 160 specific ion toxicity.

161

162

163

164

165 Table (5) described the infiltration and permeability hazard which was represented by EC
 166 and SAR combination as:

167

$$G_2 = w_2 r_2 \quad (8)$$

168

169

170 Where w is the weight value of this hazard group and r is the rating value of the parameter
 171 as given in (Table 5).

Table 5. Infiltration and permeability hazard index

	SAR-range					Rating	Suitability
	<2-3	3-6	6-12	12-20	>20		
EC-range	>700	>1200	>1900	>2900	>5000	3	High
	700-200	1200-300	1900-500	2900-1300	5000-2900	2	Medium
	<200	<300	<500	<1300	<2900	1	Low

172

173 Table (6) shows the specific ion toxicity involved SAR, Cl⁻ and B concentration ions in the
 174 water as calculated using:

175

$$G_3 = \frac{w_3}{3} \sum_{j=1}^3 r_j \quad (9)$$

176

177

178 Where j= is an incremental index, w = is the weight value and r= is the index value.

179

180

181

$$G_4 = \frac{w_4}{N} \sum_{k=1}^N r_k \quad (10)$$

182

183 Where k = is an incremental index, N = is the total number of heavy metals, w = is the weight
 184 value of this group and r = is the rating value of each parameter as given in (Table 6).

185

Table 6. Classification for heavy element toxicity

Factor	Range	Rating	Suitability
As (mg/L)	< 0.10	3	High
	0.10 ≤ 2.0	2	Medium

	> 2.0	1	Low
Cd (mg/L)	< 0.02	3	High
	0.02 ≤ 0.06	2	Medium
	> 0.06	1	Low
Co (mg/L)	< 0.05	3	High
	0.05 ≤ 5.0	2	Medium
	> 5.0	1	Low
Cu (mg/L)	< 0.20	3	High
	0.2 ≤ 5.0	2	Medium
	> 5.00	1	Low
Fe (mg/L)	< 5.00	4	High
	5.0 ≤ 20.0	3	Medium
	> 20.0	2	Low
Pb (mg/L)	< 5.1	4	High
	5.1 ≤	3	Medium
	> 10.1	2	Low
Mn (mg/L)	< 0.20	3	High
	0.2 ≤ 10	2	Medium
	> 10	1	Low
Mo (mg/L)	< 0.01	3	High
	0.01 ≤ 0.05	2	Medium
	> 0.05	1	Low
Zn (mg/L)	< 2.0	3	High
	2 ≤ 10	2	Medium
	> 10.0	1	Low

187 Table (7) shows the suitability of the irrigation water based on (IWQI). When the IWQ index
188 is <22, the suitability of this water is low, while the index ranging from 22-37 is moderate and
189 when the IWQI exceeds 37, irrigation water will be classified as high suitable.
190

Table 7. Irrigation water quality index (IWQI)

IWQ index	Irrigation Suitability
< 22.5	Low
22.5–37.6	Medium
> 37.6	High

191
192 The miscellaneous effects on sensitive crops that is included NO_3^- , HCO_3^- ions and pH of
193 the water was estimated as:
194

$$195 \quad G_5 = \frac{w_5}{3} \sum_{m=1}^3 r_m \quad (11)$$

196
197 Where m = is an incremental index, w= is the weight value and r is the index value of each
198 parameter.
199 SAR:
200

$$200 \quad \text{SAR} = \frac{\text{Na}}{\sqrt{\frac{\text{Ca} + \text{Mg}}{2}}} \quad (12)$$

201

202

203 Kelly's ratio (KR):

$$KR = \frac{Na}{Ca + Mg} \quad (13)$$

204

205 Residual sodium carbonate (RSC) was calculated as:

206

207

$$RSC = (HCO_3^- + CO_3^{2-}) - (Ca^{2+} + Mg^{2+}) \quad (14)$$

209

210 Magnesium hazard (MH) was computed through the following equation:

$$MH = \frac{Mg^{2+}}{(Ca^{2+} + Mg^{2+})} \times 100 \quad (15)$$

212 Sodium percent was calculated as follow:

$$\%Na = \frac{Na^+}{(Ca^{2+} + Mg^{2+} + Na^+ + K^+)} \times 100 \quad (16)$$

214

215 Soluble sodium percentage (SSP) was expressed below in equation 17:

$$SSP = \frac{Na^+}{Ca^{2+} + Mg^{2+} + Na^+} \times 100 \quad (17)$$

217 Total hardness (TH) was computed via the following equation:

$$TH = Ca^{2+} + Mg^{2+} \times 50 \quad (18)$$

219 Permeability index (PI) was also calculated as follow:

$$PI = \frac{Na^+ + \sqrt{HCO_3^-}}{(Ca^{2+} + Mg^{2+} + Na^+)} \times 100 \quad (19)$$

221 3. Statistical Analysis

222 The data was analyzed using statistics 21 core system and descriptive statistics was used to
223 estimate the relationship between the study parameters.

224

225 4. RESULTS AND DISCUSSION

226 4.1 Water Quality Evaluation for Drinking

227 Tables (8a and 8b) indicated that, salinity index, SAR, Na%, and RSC within the acceptable
228 groundwater quality range for drinking purpose in Al-Jour region proposed by WHO and KSA
229 [2,7]. Besides that, the following parameters.

230
231
232

Table 8a. Descriptive statistics of Al-Jouf groundwater quality

	pH	EC dS ⁻¹	Ca²⁺ mg/L	Mg²⁺ mg/L	Na⁺ mg/L	K⁺ mg/L	Cl⁺ mg/L	HCO₃⁻ mg/L	CO₃⁻² mg/L	SO₄⁻² mg/L	B mg/L	NO₃⁻ mg/L
Maximum	7.92	6.7	76.9	238.8	91.3	18.8	2100	1220	0	400.52	1.51	4.5875
Minimum	7.20	0.28	8	0	14.9	1.3	28	0	0	30.58	0	0.0875
Mean	7.5351	1.6703	28.438	22.685	42.787	6.6193	433.61	257.89	0	102.98	0.339	1.7642
SD	0.1632	1.4224	14.958	26.375	18.057	4.2569	423.15	194.89	0	93.711	0.3958	0.8477
Variance	0.0266	2.0232	223.73	695.65	326.06	18.121	179053	37980	0	8781.7	0.1567	0.7186
SE	0.0133	0.1161	1.2213	2.1535	1.4744	0.3476	34.55	15.912	0	7.6515	0.0323	0.0692
Median	7.55	1.265	24.35	15.24	38.75	4.75	280	183	0	56.67	0.183	1.7125
Skew	-0.0442	1.4533	1.0469	4.5855	0.6208	0.6577	1.5439	1.7502	0	1.5801	2.0476	0.5175

233
234
235
236
237
238
239
240

Table 8b. Descriptive statistics of Al-Jouf groundwater quality

	SAR	KR	RSC	MH	IWQI	DWQI	<i>Na (%)</i>	TH	SSP (%)	PI (%)
Maximum	15.439	4.0174	3.5	0.75	36.973	319.14	75.53	1650	0.814001	0.995973
Minimum	0.6892	0.3214	-24	0	25.421	40.771	22.34	85	0.304792	0.339279
Mean	4.2382	1.0765	-3.5583	0.2827	32.678	96.99	46.71	382.63	0.508586	0.674354
SD	3.0389	0.6101	4.8094	0.1696	3.5621	56.146	-	-	-	-
Variance	9.2347	0.3722	23.13	0.0288	12.688	3152.3	-	-	-	-
SE	0.2481	0.0498	0.3927	0.0138	0.2908	4.5843	-	-	-	-
Median	3.6748	1.0038	-2.05	0.2673	33.196	75.804	-	-	-	-
Skew	1.0851	1.8751	-1.9843	0.2752	0.5832	1.6192	-	-	-	-

241
242
243
244
245
246
247
248
249
250
251
252

253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300


The computed DWQI values for the 150 well in Al-Al-Jouf, region varied from 40.7 to 319.1 (Fig. 2). About 23.8% of studied wells were considered poor water “class (III)”, 9.90% very poor water “class (IV)”, 45.6% good water and 20.53% excellent water. The higher DWQI values for some wells have may be due to high values of salinity, cations and NO_3^- . The relationship between the water quality parameters is shown in (Table 9). The results indicated that, the TDS variations are mainly controlled by SO_4^{2-} ($r=0.91$), Cl^- ($r= 0.91$) and TH ($r= 0.98$). The calculated WQI showed also highly significant interrelation between its values and TDS ($r=0.96$), HCO_3^- ($r=0.76$), Cl^- ($r=0.93$), SO_4^{2-} ($r=0.85$), and TH ($r=0.94$). in addition, TDS have strong positive correlation with Mg^{2+} (0.94), Cl^- (0.95), SO_4^{2-} (0.89) Ca^{2+} (0.77) and Na^+ (0.73) and moderate correlation with NO_3^- (0.67). The results in agreement with those reported by [19,24, 35].

Table 9. Correlation coefficient matrix between water quality parameters and WQI

Parameters	pH	TDS	Ca ²⁺	Mg ²⁺	Na ⁺	K ⁺	HCO ₃ ⁻	Cl ⁻	SO ₄ ⁻²	NO ₃ ⁻	B	TH	WQI
pH	1	-0.1628	-0.186	0.1096	-0.078	0.0293	-0.0247	-0.1906	-0.1658	-0.1145	-0.1466	-0.1849	-0.1323
TDS		1	-0.1603	0.3175	0.3815	-0.6234	0.6697	0.9685	0.9088	0.185	0.4932	0.9588	0.9658
Ca²⁺			1	-0.1351	0.6407	0.5133	-0.343	-0.0988	-0.2443	-0.0502	-0.2313	-0.1016	-0.1567
Mg²⁺				1	0.0117	-0.207	0.2374	0.2856	0.2865	0.1303	0.262	0.4446	0.3737
Na⁺					1	0.256	0.0642	0.4212	0.315	-0.0096	0.1662	0.3153	0.3916
K⁺						1	-0.5328	-0.6009	-0.5493	-0.1476	-0.3252	-0.6573	-0.5611
HCO₃⁻							1	0.5942	0.5579	0.1379	0.2604	0.6501	0.7607
Cl⁻								1	0.8618	0.1703	0.475	0.9479	0.9363
SO₄⁻²									1	0.1417	0.4932	0.8552	0.8563
NO₃⁻										1	-0.0086	0.1888	0.2125
B											1	0.4762	0.4667
TH												1	0.9401
WQI													1

301

302



304
305
306
307
308
309
310
311
312
313
314

Fig. 2. Drinking water quality index (DWQI) in Al-Jouf region, KSA.

315 **4.2 Evaluation of Water Quality for Irrigation Purposes**

316 The major ions chemistry of groundwater samples of Al-Jouf were statistically analyzed and
 317 the results are shown in (Table 8a, 8b). The concentrations of Ca^{++} , Mg^{++} , Na^+ , and K^+ ions
 318 ranged between 8 and 76.9, 0 and 238, 14.9 and 91, 1.30 and 18.9 mg/L with a mean value
 319 of 28.5, 22.7, 42.8, 6.7, mg/L, respectively. The maximum permissible limit of these ions in
 320 irrigation water is 80, 55, 210, and 13 mg/L, respectively [40, 42]. The most of the wells
 321 water are considered suitable for irrigation usage with respect to Ca^{++} , Mg^{++} and Na^+ . the
 322 concentrations of HCO_3^- , Cl^- , SO_4^{2-} , and NO_3^- ions lie in between: 0 - 1220; and 28 – 2100;
 323 and 30.5 - 403.52; and 0.08 - 4.58 mgL^{-1} , respectively with a mean value of 257.9, 432,
 324 102.9, and 1.77 mg/L, respectively. The maximum permissible limit of HCO_3^- , Cl^- , SO_4^{2-} , and
 325 NO_3^- in irrigation water is 120, 250, 250, and 30 mg/L [19, 40, 41, 43]. According to the
 326 standards, most of the wells are suitable for irrigation. The pH values of the water samples
 327 varied from 7.20 to 7.92 with a mean value of 7.53. These mean that all studied water
 328 samples were within safe limit with respect to pH [19]. The TDS varied from 179.2 to 4288
 329 mg/L^{-1} with an average value of 1024 mg/L. Salinity is between excellent and doubtful based
 330 on the classification of TDS suggested by Ayers and Westcot [19]. The suitability of wells is
 331 summarized in (Table 10). The excessive Na^+ content in water sample reduces the
 332 permeability, and reduce the available water for the plant. The water containing excessive
 333 amount of Na^+ may immobilize other nutrient ions particularly Ca^{++} and Mg^{++} , which can
 334 result in deficiencies of these elements in plants [34]. Excess amounts Na^+ , Ca^{2+} , Mg_2^+ and
 335 HCO_3^- in the irrigation water affect soil permeability through widespread irrigation water use
 336 [35]. The SAR values of the groundwater samples varied between 0.6892 to 15.439 with an
 337 average value of 4.285 (Table 9). About 85 % of the SAR values of the water samples were
 338 less than 10 and are classified as excellent for irrigation [32]. In addition, spatial distribution
 339 of SAR in the region were measured and presented in (Fig. 3) as the values increasing in
 340 north part of the region which might create a sodium hazard with intensive agricultural
 341 activities. This area should be monitored closely to prevent any accumulation of Na and
 342 cause further sodium hazard. [14], was also determined the hazardous effect of sodium on
 343 water quality for irrigation usage in KR index. A Kelly's ratio of more than 1 indicates
 344 excessive Na in water. Therefore, water with a KR less than 1 is considered suitable for
 345 irrigation; on the other hand, the ratio more than 1 is unsuitable. The KR ratios in the studied
 346 water ranged between 0.3214 and 4.02 with an average value of 1.08. About 60 % of the
 347 studied waters are considered suitable for irrigation [19].

348 The B concentrations were within permissible limits in the 92% of water samples and the
 349 remaining samples were considered slight to moderate B toxicity. The RSC value ranged
 350 between 3.50 to -24, with an average value of -3.5. About 98 % of the studied waters are
 351 considered suitable for irrigation. Usually, Ca_2^+ and Mg_2^+ are in an equilibrium state in
 352 groundwater. The higher value of Ca_2^+ and Mg_2^+ in water can increase soil pH (therefore soil
 353 converting it to saline soil, resulting in decrease in the P availability. Excess concentration of
 354 Mg_2^+ in groundwater affects the soil quality by converting it into alkaline and decreases the
 355 crop yield. The excess amount of Mg_2^+ ions in waters damage the soil quality which causes
 356 low crop production. The MH in the studied water ranged between 0 to 0.75, with an average
 357 value of 0.28. About 100 % of the studied waters are considered suitable for irrigation. The
 358 Na (%) in the studied water ranged between 22. 0 and 75.8, with an average value of 46.3.

359

Table 10. Classification of water samples in Al-Jouf aera for irrigation based on some characteristics

Parameters	Range	Water	References
------------	-------	-------	------------

		class
TDS	<450	Best
	450-2000	Moderate
	>2000	Hazard
SAR	0-15	Excellent
	15-20	Good
	20-27	Doubtful
	>27	Unsuitable
RSC	<1.25	Good
	1.25-2.5	Doubtful
	>2.5	Unsuitable
KR	<1	Suitable
	1-2	Marginal suitable
	>2	Unsuitable
SSP	<50	Good
	>50	Unsuitable
PI	>75	C-I [5,16,24,34]
	25-75	C-II
	<25	C-III
MH	<50	Suitable
	>50	Harmful
Na%	<25	Unsuitable
	25-45	Excellent
	45-65	Good
	65-85	Permissible
	>85	Doubtful
T.H	<80	Unsuitable
	80-155	Soft
	155-310	Moderately Hard
	>310	Hard
		Very Hard

360
361




Fig. 3. Spatial distribution of SAR values in Al-Jouf area

362
363
364
365
366
367
368
369
370

The SSP in the studied water ranged between 0.81 and 0.30, with an average value of 0.50 and 100 % of the studied waters are considered suitable for irrigation. The (TH) in the studied water ranged between 85 to 1650, with an average value of 382.6 and about 22 % of the studied waters are considered suitable for irrigation. The PI in the studied water ranged between 0.97 and 0.34, with an average value of 0.67 and about 100 % of the studied waters are considered suitable for irrigation.

371 **4.3 Classes of Salinity and Alkalinity Hazard**

372 The salinity and alkalinity hazard class of studied well samples were C2-S1, C3-S1, C3-S2
373 and C4-S3 (Fig. 4). The result indicated that, the groundwater possesses moderate to high
374 salinity hazards with low to medium Na hazards. The excessive amount of salts in some
375 wells of the region can be one of the major problems in the study area. According to [1],
376 groundwater is an essential source for crop rising and food production in this region. He
377 have categorized the groundwater into four types such as low salinity ($EC < 250 \mu S/cm$),
378 medium salinity (250 to $750 \mu S/cm$), high salinity (750 to $2250 \mu S/cm$), and very high salinity
379 ($>2250 \mu S/cm$). Some of the wells, mainly in the north part of the region, which cannot be
380 used for irrigation without special circumstances.

381
382




Fig. 4. Groundwater salinity classification used for irrigation

383
384
385
386
387




Fig. 5. Electrical conductivity spatial distribution in Al-Jouf region, Saudi Arabia

388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403

4.4 Hydro Chemical Aspects

The piper diagram shows that the water types of Al-Jouf well waters are rich in Ca^{2+} , Mg^{2+} , Cl^- and SO_4^{2-} ions (Fig. 6). The piper diagrams provided a convenient method to classify water types collected from different groundwater resources, based on the ionic composition of different water samples [2,7]. The chemical data of well waters of Al-Jouf region are plotted in Gibbs's diagrams (Fig. 7). The distribution of sample points suggests that the chemical weathering of rock-forming minerals and evaporation are influencing the groundwater quality. The rock–water interaction process includes the chemical weathering of rocks, dissolution–precipitation of secondary carbonates and ion exchange between water and clay minerals. The evaporation greatly increases the concentrations of ions formed by chemical weathering, leading to higher salinity. The moving of groundwater sampling points in the Gibbs field towards the evaporation domain from the rock domain suggests that increases the salinity [7].

408

Fig. 7. Diagram showing the mechanism that control of groundwater quality

409

410

4.5 Irrigation Water Quality Index (IWQI)

411


412

413

414

415

IWQI was calculated based on the proposal by Simsek, Gunduz [4] and the results presented in (Fig. 8). Accordingly, all wells water in Al-Jouf region, KSA classified as medium in suitability for irrigation purposes. These findings agreed with those reported by [19].



416

417

418

419

420

Fig. 8. Irrigation water quality index (IWQI) for 150 wells in Al-Jouf region, KSA

CONCLUSIONS

421

422

423

424

425

426

427

428

429

430

431

Water quality indices have been calculated to assess the suitability of groundwater for drinking and irrigation purposes in Al-Jouf, KSA. About 12 parameters have been considered to calculate the DWQI which included: pH, TDS, cations, anions and B concentration. The results indicated that, about 20.5% of well water considered as excellent water, 45.6% classified as good water, 23.8% and 9.9 % were poor and very poor water for drinking purposes, respectively. The results revealed that salinity and alkalinity hazards classes of water samples were 60% of the groundwater is in C2-S1, C3-S1, 25% in C3-S2, and 15% in C4-S3. The hydro chemical analysis showed that the analyzed water samples correspond mainly to Mg, Ca, SO₄ and Cl⁻ ions. The IWQI value was classified as medium in terms of suitability, thus, this could be used for irrigation purposes for all the crops. As a result, the Al-Jouf region could contribute greatly to the sustainable agriculture.

432

ACKNOWLEDGMENTS

433

434

435

The authors are thankful to the deanship of scientific research, college of food and agricultural sciences; King Saud University, Saudi Arabia for financial support and cooperation for the completion of this research study.

436

COMPETING INTERESTS

437 NO conflict of interest.

438 **Authors' Contributions**

439 Ahmed Nazal Al-Shemary, is the first author as graduate student, his contribution was in
440 execution the research work, collecting data, analyzing data, writing first draft of the
441 manuscript and presenting the work in its form in terms of tables and figures. Khalid Dawi Al-
442 Otabi is co-advisor and work closely with first author on collecting data and interpretation of
443 it.

444 Abduroul Mosa Al-Omran is the advisor of the graduate student his contribution was in
445 design the work, analyzing data, interpretation of the data and final version of the manuscript

446

447 **REFERENCES**

448 1. Al-Hadithi M. Application of water quality index to assess suitability of groundwater quality
449 for drinking purposes in Ratmao-Pathri Rao watershed, Haridwar district India. *Journal of*
450 *Scientific and Industrial Research*; 2012; 23:1321-1336.

451 2. Al-Omran A, Al-Barakah F, Altuquq A, Aly A, Nadeem M. Drinking water quality
452 assessment and water quality index of Riyadh, Saudi Arabia. *Water Qual Res J Can*; 2015;
453 50(3):287–296.

454 3. Al-Omran MA, El-Maghraby SE, Aly AA, Al-Wabel IM, Al-Asmari ZA, Nadeem ME.
455 Quality assessment of various bottled waters marketed in Saudi Arabia. *Environ Monit*
456 *Assess*. 2012; 185(8):6397–6406. doi:10.1007/s10661-012-3032-z.

457 4. El-Amier, Y.A.; Kotb, W.K.; Bonanomi, G.; Fakhry, H.; Marraiki, N.A.; Abd-ElGawad, A.M.
458 Hydrochemical Assessment of the Irrigation Water Quality of the El-Salam Canal, Egypt.
459 *Water*; 2021(13):2428. <https://doi.org/10.3390/>.

460 5. Shahid S, Chen X, Hazarika MK. Evaluation of groundwater quality for irrigation in
461 Bangladesh using geographic information system. *J Hydrol Hydromech.*; 2006; 54:3–14.

462 6. Lateef, K. H. Evaluating of ground water quality for drinking water purpose for Tikrit and
463 Samarra cities using water quality index. *European journal of Scientific Research ISSN*
464 *1450-216X Vol. 58 No. 4 pp. 472 481 (2011)*.

465 7. Aly AA, Al-Omran AM, Alwabel M, Almahaini A, Alamari M. Hydrochemical and quality of
466 water resources in Saudi Arabia groundwater: a comparative study of Riyadh and Al-Hasa
467 regions. *Proc Int Acad Ecol Environ Sci.*, 2016; 3(1):42–51.

468 8. Debels, P. Evaluation of water quality in the Chillan River (Central Chile) using
469 physicochemical parameters and a modified water quality index. *Environmental monitoring*
470 *and assessment*. 2005; 110(1): 301-322.

471 9. Saeedi M, Abessi O, Sharifi F, Meraji H. Development of groundwater quality index.
472 *Environ Monit Assess*. 2010; 163(1-4):327–335. <https://doi.org/10.1007/s10661-009-0837-5>.

473 10. Semerjian LA. Quality assessment of various bottled waters marketed in Lebanon.
474 *Environ Monit Assess*. 2011; 172(1-4):275–285. <https://doi.org/10.1007/s10661-010-1333-7>.

- 475 11. Tiwari AK, Singh PK, Mahato MK. GIS-based evaluation of water quality index of
476 groundwater resources in West Bokaro Coalfield, India. *Curr World Environ.* 2014; 9(3):843–
477 850. <https://doi.org/10.12944/CWE.9.3.35>.
- 478 12. Al-Othman, A. A. Evaluation of the suitability of surface water from Riyadh Mainstream
479 Saudi Arabia for a variety of uses. *Arabian Journal of Chemistry*, 2019; 12(8): 2104-2110.
- 480 13. Backman, B. Application of a groundwater contamination index in Finland and Slovakia."
481 *Environmental geology*; 1998; 36(1): 55-64.
- 482 14. Kelly WJ. Permissible composition and concentration of irrigated waters. (1940) 607.
- 483 15. Kumar SS, Marender B, Dhakate R, Sankaran S, Kumar KR (2016) Assessment of
484 groundwater quality for drinking and irrigation use in shallow hard rock aquifer of
485 Pudunagaram, Palakkad District, Kerala. *Appl Water Sci* 6(2):149–167.
486 <https://doi.org/10.1007/s13201-014-0214-6>.
- 487 16. Stigter TY, Ribeiro L, Carvalho Dill AM. Application of a groundwater quality index as an
488 assessment and communication tool in agro-environmental policies—two Portuguese case
489 studies. *J Hydrol.*, 2005; 327(3-4):578–591.
- 490 17. Islam MA, Zahid A, Rahman MM, Rahman MS, Shammi M. Investigation of groundwater
491 quality and its suitability for drinking and agricultural use in the south-central part of the
492 coastal region in Bangladesh. *J Expo Health*, 2017; 9(1):27–41. doi:10. 1007/s12403-016-
493 0220-z.
- 494 18. Abbas A, Nader Y, Amir HM., Ramin N, Majid R, Mahmood Y & Mahmood A. Evaluation
495 of groundwater quality using water quality index and its suitability for assessing water for
496 drinking and irrigation purposes: Case study of Sistan and Baluchistan province (Iran) ,
497 Human and Ecological Risk Assessment: An International Journal, 2019; 25:4, 988 1005,
498 DOI: 10.1080/10807039.2018.1458596.
- 499 19. Ayers, R. S., and D. W. W. Water quality for agriculture. *FAO Irrigation and drainage*
500 *paper 29 Rev. 1.*" Food and Agricultural Organization. Rome 1 (1985): 74.
- 501 20. Derdour, A., L. G., and M. A. Assessment of drinking and irrigation water quality using
502 WQI and SAR method in Mader sub-basin, Ksour Mountains, Algeria. *Sustainable Water*
503 *Resources Management* 2021; 7(1):1-14.
- 504 21. Appelo, C. J., and D. P. *Geochemistry, groundwater and pollution.* CRC press (2004).
- 505 22. Ministry of Agriculture and Water R. In. Edited by Department of Economic Studies and
506 Statistics ASY. 4th ed. ed. Saudi Arabia: Ministry of Agriculture and Water, Riyadh; 1994.
- 507 23. Maiti S. *Handbook of Methods in Environmental Studies, 1: Water and Wastewater*
508 *Analysis.* ABD publishers; 2004).
- 509 24. Tabatabai, M. A. et al. "Sulfur." *Methods of Soil Analysis: Part 3 Chemical Methods* 5
510 (1996): 921-960.
- 511 25. Al-Omran AM, Falatah AM, Al-Matrood SS. Evaluation of irrigation well water quality in
512 Riyadh region, Saudi Arabia. *J King Abdulaziz Univ.* 2005; 16(2):23–40.

- 513 26. Page, A. L., and D. R. K. Methods of soil analysis. American Society of Agronomy,
514 (1982).
- 515 27. American Public Health Association, et al. Standard methods for the examination of
516 water and wastewater. Vol. 2. American Public Health Association., 1912.
- 517 28. Horton, R. K. An index number system for rating water quality. J Water Pollut Control
518 Fed 37.3 (1965): 300-306.
- 519 29. Varouchakis EA, Hristopulos DT. Improvement of groundwater level prediction in
520 sparsely gauged basins using physical laws and local geographic features as auxiliary
521 variables. Adv Water Res., 2013; 52:34–49.
- 522 30. Ramakrishnaiah, C. R., C. S., and G. R. Assessment of water quality index for the
523 groundwater in Tumkur Taluk, Karnataka State, India. E-Journal of chemistry, 2009; 6(2):
524 523-530.
- 525 31. Gibbs, R. J. Mechanisms controlling world water chemistry." Science 170.3962 (1970):
526 1088-1090.
- 527 32. Richards, L. A. et al. Diagnosis and improvement of saline and alkali soils. Vol. 78. No.
528 2. LWW, 1954.
- 529 33. Romanelli A, Lima ML, Londono OMQ, Martinez DE, Massone HE. A Gis-based
530 assessment of groundwater suitability for irrigation purposes in flat areas of the Wet Pampa
531 Plain, Argentina. Environ Manag., 2012; 50:490–503.
- 532 34. Sharifi Z, Safari Sinegani AA. Arsenic and other irrigation water quality indicators of
533 groundwater in an agricultural area of Qorveh Plain, Kurdistan, Iran. Am Eurasian J Agric
534 Environ Sci. 2012; 12(4):548– 555.
- 535 35. Sharaf MA, Hussein MT. Groundwater quality in the Saq aquifer, Saudi Arabia.
536 Hydrological Sciences-Journal-des Sciences Hydrologiques 1997; 4(5):683–696.
- 537 36. Todd, D. K., and L. W. M. Groundwater hydrology. John Wiley & Sons (2004).
- 538 37. Läuchli A., Epstein, E. (1990). Plant responses to saline and sodic conditions," in
539 Agricultural salinity assessment and management, K. K. Tanji, ed., ASCE Manuals and
540 Reports on Engineering Practice No. 71, ASCE, Reston, Va., 113–137.
- 541 38. Ding Z., Majrashi MA., Ghoneim AM., Esmat F. Ali, Mamdouh A Eissa, Rania El Shal
542 (2022). Irrigation and biochar effects on pearl millet and kinetics of ammonia volatilization
543 from saline sandy soils. J Soil Sci Plant Nutr., <https://doi.org/10.1007/s42729-021-00753-0>.
- 544 39. Saleh AH; Gad M; Khalifa MM.; Elsayed S; Moghanm FS.; Ghoneim AM; Danish, S;
545 Datta, R; Moustapha, ME.; Abou El-Safa, MM. Environmental pollution indices and
546 multivariate modeling approaches for assessing the potentially harmful elements in bottom
547 sediments of Qaroun Lake, Egypt. Journal of Marine Science and Engineering, 2021; 9,
548 1443. <https://doi.org/10.3390/jmse9121443>.
- 549 40. Zheli Ding, Esmat F. Ali, Sami Ali Aldhumri, Adel M Ghoneim, Ahmed MS. Kheir, Marwa
550 GM. Ali, Mamdouh A Eissa. Effect of amount of irrigation and type of P fertilizer on potato

551 yield and NH₃ volatilization from alkaline sandy soils. J Soil Sci Plant Nutrition, 2021; 21(2):
552 1565-1576. <https://doi.org/10.1007/s42729-021-00462-8>.

553 41. Mahmoud M. Gaballah, Adel M. Ghoneim, Mohamed I. Ghazy, Hassna M.
554 Mohammed, Raghda M. Sakran, Hafeez Ur Rehman and Noraziyah Abd Aziz
555 Shamsuddin Root traits responses to irrigation intervals in rice (*Oryza sativa*).
556 International Journal of Agriculture & Biology, 2021; 26: 23-30.

557 42. Adimalla N, Qian H. Groundwater chemistry, distribution and potential health risk
558 appraisal of nitrate enriched groundwater: a case study from the semi-urban region of
559 South India. Ecotoxicol Environ Saf. 2021; 207:111277. [https://doi.](https://doi.org/10.1016/j.ecoenv.2020.111277)
560 [org/10.1016/j.ecoenv.2020.111277](https://doi.org/10.1016/j.ecoenv.2020.111277).

561

562

563

564