

## Original Research Article

### EFFECT OF ZN AND FE ENRICHED FYM ON YIELD, NUTRIENT CONTENT AND UPTAKE BY COWPEA IRRIGATED WITH SODIC WATER

#### Abstract

A pot experiment was conducted during *kharif* 2018-19 at S.K.N. College of Agriculture, Jobner (Rajasthan) to study the “Effect of Zn and Fe enriched Farmyard manure on nutrient availability and yield of cowpea [*Vigna unguiculata* (L.)wilczek] under sodic water irrigation” to evaluate the effect of different sodic water and sources of Zn and Fe on yield, nutrient concentration and uptake by cowpea on loamy sand soil. The experiment comprising of 15 treatment combinations replicated three times, was laid out randomized block design with three levels of sodic water (2, 4 and 6 mmol L<sup>-1</sup>) and five levels of sources of Zn and Fe (0, ZnSO<sub>4</sub>.7H<sub>2</sub>O, FeSO<sub>4</sub>.7H<sub>2</sub>O, Zn enriched FYM, Fe enriched FYM) as variables. The total and number of nodules per plant, nodule index, pods per plant and seeds per pod were counted at stage and harvesting respectively. Results revealed that under 6 mmol L<sup>-1</sup> Residual sodium carbonate of irrigation water, the seed and straw yield of, nutrient concentration and uptake viz., P, K, Zn and Fe in seed and straw decreased significantly with all levels of sodic water and maximum reduction was recorded with the application of 6 mmol L<sup>-1</sup> of sodic water, while N in seed and straw, protein content in seed increased significantly. The application of Zn enriched FYM increased seed and straw yield, nutrient concentration and uptake viz., N, P, K, Zn and Fe in seed and straw, protein content in seed.

**KEYWORDS:** Sodic water, Zn enriched FYM, Fe enriched FYM, Seed and Straw yield, Nodule Index

#### Introduction

Pulses are gaining worldwide importance as they are the cheap source of protein in diet and have ability and soil fertility through biological nitrogen fixation as well as of ample amount of residues to the soil (Reddy and Reddy, 2010). India is the largest producer and consumer of pulses in the world accounting for about 29 and 19 per cent of the world's area and production, respectively. The per capita availability of pulses in India is 35.5 g day<sup>-1</sup> as against the minimum requirement of 70 g day<sup>-1</sup>. Cowpea [*Vigna unguiculata* (L.) Wilczek]

commonly known in India as *lobia* is one of the important *khariif* pulse crops grown for vegetable, grain, forage and green manuring. This crop has great importance because of short duration, high yielding and variety. It is a considerable promising pulse crop in farming. Unlike other legumes, cowpea is not so tolerant to sodicity. In many arid and semi-arid regions, of saline and sodic water for irrigation in the absence of appropriate soil-water-crop management often leads to the of salinity and sodicity in the soil profile which adversely the crop productivity. The quality of irrigation water plays a vital role in crop production. The use of sodic water for irrigation adversely affects of soil by influencing the uptake of nutrients and many soil properties (Chauhan *et al.* 1988). This problem becomes more aggravated when the carbonate and bicarbonate of sodic water occur in association with sodium creating the problem of residual sodium carbonate (Doodhwal *et al.* 2020).

Zinc plays a vital role in chlorophyll, protein and nucleic acid and helps in nitrogen and phosphorus by plants as it acts as an activator of dehydrogenase and proteinase enzymes, directly or indirectly in carbohydrates and protein. Zinc is a constituent of tryptophan which is a precursor of auxin hormone (Purushottam *et al.* 2018). Zinc deficiency has been reported to be the most widespread micro-nutritional disorder of the food crops in India as well as the world over. Zinc is a crucial component of the practices recommended for sodic soils reclamation. Deficiencies of Fe, Mn and Cu are much less extensive. The available Zn content of Indian soils varied from traces to 22 mg kg<sup>-1</sup> and 47 per cent of Indian soils were found to be deficient in Zn. Zn deficient soils cause low crop production leading to malnutrition in humans and livestock (Kumar and Dhaliwal, 2021). Continuous use of high sodic water increases the ESP and pH which decreases the availability of Zn (Meena *et al.* 2017). As the soil pH, the ionic form of Zn changed to hydroxide form, which is insoluble and unavailable to plants. Although the high RSC water can be used successfully by applying higher doses of zinc sulphate, Zinc helps in inducing alkalinity tolerance in by enhancing its efficiency in utilizing K, Ca and Mg the adverse effect of sodicity (Raj and Raj, 2019). Thus, reducing of Na/K and Na/Ca ratio in plant tissues to mitigate the adverse effect of alkalinity in is an important aspect (Mishra, 2001). Iron being an essential micronutrient takes part in the metabolic activities of the plant. It acts as an activator of dehydrogenase, proteolase and peptidases enzymes, directly or indirectly involved in the synthesis of carbohydrates and proteins. Iron in chloroplast reflects the presence of cytochromes for performing various photosynthetic reduction processes (Doodhwal *et al.* 2020). Zn and Fe application in the enriched form may enhance the fertilizer use efficiency and increase crop yield. However, no

systematic study has been conducted on Zn and Fe-Enriched Farmyard manure in these soils irrigated with high RSC water in the region of legume crops. The productivity could be sustained through use of organic and inorganic fertilizers. The organic manures for supplement of Zn and Fe deficiency in soils has been emphasized by several workers (Meena et al. 2014). FYM enriched zinc and iron reacts with native reserves of Zn and Fe elements and them available to plants. It has been reported that organic wastes such as presumed cattle and poultry litter and Farmyard manure with or without zinc alleviated zinc deficiency (Amer et al. 2020). The present study was aimed at assessing the effect of Zn and Fe enriched FYM on yield, nutrient content and uptake by cowpea.

### Materials and Methods

**Experimental site and soil:** A pot experiment was conducted in house of Plant Physiology and samples analysis were done in the Department of Soil Science and Agricultural Chemistry, S.K.N. College of Agriculture, Jobner (Rajasthan) during the *kharif* season 2018-2019. Jobner is located at 75.28<sup>0</sup> East longitudes and 26.06<sup>0</sup> North at an altitude of 427 metres above mean sea level. This region comes under Agro Climatic Zone III-a (Semi-arid eastern plain). The climate of the region is typically semi-arid characterized by the aridity and salinity with extremes of temperature both during summer and winter. The average rainfall of this region is about 400 to 500 mm, which is mostly received between July and September (Monsoon season). The loamy sand soil was used for experiment. The experiment comprising of 15 treatment combinations replicated three times, was laid out in randomized block design with three levels of sodic water (2, 4 and 6 mmol L<sup>-1</sup>) and five levels of sources of Zn and Fe (0, ZnSO<sub>4</sub>.7H<sub>2</sub>O @ 12.5 mg/kg soil, FeSO<sub>4</sub>.7H<sub>2</sub>O @ 25 mg/kg soil, Zn enriched FYM 250 mg/kg soil, Fe enriched FYM @ 250 mg/kg soil) as variables.

**Quality of irrigation water:** The different sodic water was prepared artificially by dissolving amount of NaHCO<sub>3</sub>, NaCl, Na<sub>2</sub>SO<sub>4</sub>, CaCl<sub>2</sub> and MgCl<sub>2</sub> in base water (control). The tap water (base water) was used for first irrigation in all pots and later on irrigated 6 times with water of varying RSC levels as per treatment. The composition of prepared water is given in table 1.

**Preparation of Zinc and Fe Enriched FYM:** The locally available organics like farmyard manure (FYM) were used for their enrichment with Zn and Fe. The enrichment process was started 60 days before their use in cowpea. The known quantity of organics was filled in the pre-dug pits of 1.5' x 1.5' x 1.5' size. The FYM were thoroughly mixed with the solutions of ZnSO<sub>4</sub>.7H<sub>2</sub>O and FeSO<sub>4</sub>.7H<sub>2</sub>O equivalent to 25 kg Zn per ha and 50 kg Fe per ha,

respectively supplied through 500 kg of FYM. The moisture percentage of FYM after mixing 70% throughout the enrichment process. The cow dung slurry @ 1% was applied as a starter inoculum of microorganisms to boost up the microbiological activities for process of composting to fix the externally added Zn and Fe through zinc sulphate and iron sulphate, respectively. The pit was covered by sheet for natural chelation during the process of composting. The mixture was turned over weekly and moisture loss was compensated during the process of enrichment. The periodical samples were taken from the pit for of water-soluble Zn and Fe content when the value of water-soluble Zn and Fe appeared to be more or less constant and the enrichment process was considered as complete. It was found that the process was almost completed within 6 to 7 weeks. After enrichment, the required quantity of zinc and Fe enriched FYM @ 250 mg/kg soil was thoroughly mixed in before sowing the crop. As per the crop production guide, the recommended dose of fertilizers (40:20:20 kg N: P<sub>2</sub>O<sub>5</sub>: K<sub>2</sub>O ha<sup>-1</sup>) applied as basal in the form of urea, single super phosphate and murate of potash for all the treatments. The Zinc sulphate (ZnSO<sub>4</sub>) and ferrous sulphate (FeSO<sub>4</sub>) as Enriched Farm Yard Manure (EFYM) form applied as per the treatment structures.

**Plant analysis:** The plant samples were first dry in open sun and then dry at 65<sup>0</sup>C until the attainment of constant weight. The oven-dried plant samples were help of a stainless-steel grinder for subsequent analysis. The chemical analysis of the plant samples was carried out by wet digesting with HNO<sub>3</sub>:HClO<sub>4</sub> (4:1) di-acid mixture as per the procedure outlined by Jackson (1973). For estimation of nitrogen, phosphorus, potassium, zinc and iron content in representative samples of seed and straw taken at the time of threshing were grind to a fine powder. Nutrient content in seed and straw were estimated by using standard methods. Total N was determined using N Auto Analyzer. K was directly measured in flame photometer and P content in plant parts were measured in at 420 nm. For Zn and Fe content digested plant material was measured in AAS. The uptake of nitrogen, phosphorus, potassium, zinc and iron at harvest in seed and straw was estimated by using the following formula.

$$\text{Nutrient content in Seed/ straw (\%)} \times \text{yield (g/pot)}$$

$$\text{N, P, K uptake (mg/pot)} = \text{-----} \times 1000$$

$$100$$

$$\text{Zn and Fe uptake } (\mu\text{g pot}^{-1}) = \frac{\text{Nutrient content in Seed/ straw (ppm)} \times \text{Seed/ straw yield (g pot}^{-1})}{10000} \times 1000$$

**Statistical analysis:** The statistical analysis of the data on the final value of seed and straw yield and soil analysis for Electrical conductivity, pH, Sodium Absorption ratio, organic carbon and DTPA-Zn and Fe were done by method of analysis of variance. To compare the treatment difference, the critical difference (CD) at 5 per cent level of significance was calculated as per method described by Panse and Sukhatme (1967) wherever 'F' test was found significant.

### **Result and Discussion**

**Effect of Sodic water:** The seed and straw yield of cowpea decreased significantly with increasing levels of Sodium Absorption ratio in irrigation water (Table 2). The decrease in the seed yield due to W<sub>4</sub> and W<sub>6</sub> was 16.42 and 48.77 per cent over W<sub>2</sub>, respectively. The W<sub>4</sub> and W<sub>6</sub> decreased the straw yield to the extent of 16.50 and 49.10 per cent over W<sub>2</sub>, respectively. Seed and straw yield of cowpea significantly decreased with increasing of sodic. This may be explained on the basis that increasing level of sodicity of soil increased the SAR and pH of soil resulting decreased availability of N, P, K, Zn, Fe, Ca and Mg but increased the uptake of Na. The higher amount of Na may adversely affect the physiological, metabolic and enzymatic activities and utilization of photosynthates. There are several evidences that cationic (Ca, Mg, Na and K) imbalance could lead to disturbances in photosynthesis and activity of stroma enzymes (Brand and Beckler, 1984; Plaut and Grieve, 1988). The inability of the crop to grow under high SAR is due to the toxicity of Na itself and Ca and K frequently becomes factor for plant growth (Dwivedi and Burrows, 1979). Restricted supply of Ca was reciprocated by a high Na content and was shown to affect the growth of the roots and shoots. Such a reduction in growth under Ca deficiency caused by Na accumulation was attributed to K leakage (Ben-Hauyyim *et al.* 1987).

The maximum N content in seed and straw was recorded under W<sub>6</sub> (6 mmol L<sup>-1</sup> RSC) and it registered 13.27 and 9.40 per cent in seed and straw, it was 32.73 and 16.84 per cent higher over W<sub>2</sub> and W<sub>4</sub>, respectively. The increase in N content in with increasing level of sodic waters may be attributed to a less production of crop resulting in concentration of N. The results find support from the work of Yadav (2001) and Singh *et al.* (2005). The W<sub>4</sub> and W<sub>6</sub> decreased the N uptake by seed to the extent of 13.50 and 42.01 per cent and in straw 5.06 and 32.41 per cent over W<sub>2</sub>, respectively. Similar results were reported by Doodhwal *et al.* 2020.

The application of W<sub>4</sub> and W<sub>6</sub> decreased the P, K, Zn and Fe content in seed and straw (Table 5). The decrease in K content in seed and straw of cowpea due to sodic water

might be due to the antagonistic effect of excess Na on the absorption of K (Doodhwal et al. 2020). A number of subsequent studies have established that increasing sodicity decreased the K and Ca concentration and increased Na concentration in tissue (Mahmood 2011) and (Garhwal *et al.* 2011). Zn and Fe content in grain and straw decreased significantly with increasing levels of RSC of irrigation water. It might be due to the conversion of  $Zn^{2+}$  and  $Fe^{2+}$  to its unavailable form to uptake under a sodic soil environment generated by high RSC of irrigation water. Similar results also reported by Jatav (2000).

The P, K, Zn and Fe uptake decreased significantly with increasing levels of sodic water (Table 5). This might be due to the fact that sodic waters had increased the ESP and pH of soil. The higher sodicity of the soil could have decreased the mobility of P due to presence of  $CO_3^{2-}$  ions. At higher pH, the proportion of  $HPO_4^{2-}$  and  $PO_4^{3-}$  increased over  $H_2PO_4^-$ . The  $OH^-$  ions, thus, decrease the availability of P to the plant. The physiological availability of P in alkali soil is a function of pH and it decreased as the pH increase over the alkaline range (Yadav et al. 2019).

Protein content in seed of cowpea significantly increased with increased level of sodic water. According to Strogonov and Okinina (1961) as earlier, the N taken up by plants gets accumulated in organs as protein and is not available for plant growth leading to increased content of N, which ultimately increased the protein content with increasing levels of sodic water. The results find support from the work of Jatav (2000); Yadav (2001) and Sharma (2003).

**Effect of Sources of Zn and Fe:** The seed and straw yield of cowpea (Table 2) increased significantly with the application of Zn enriched FYM ( $T_3$ ). Application of Zn enriched FYM increase the seed yield by the extent of 68.94, 16.67, 32.58 and 8.07 per cent over  $T_0$ ,  $T_1$ ,  $T_2$  and  $T_4$ , respectively. An investigation of the data further showed that  $T_3$  registered significantly higher straw yield over  $T_0$ ,  $T_1$ ,  $T_2$  and  $T_4$ . Treatment  $T_3$  showed an increase of 78.40, 16.67, 32.58 and 7.97 per cent in straw yield over  $T_0$ ,  $T_1$ ,  $T_2$  and  $T_4$ , respectively. Application of Zn enriched FYM enhanced the seed yield and straw yield over control. The significance difference in seed yield under the influence of FYM and Zn was largely a function of improved growth and the consequent increase in different yield attributes. Further, FYM increases the activities of nitrogen fixation bacteria and increased of humification. Humic acid in FYM enhanced the availability of both native and added in soil and thus plant growths, yield attributes and yield increased. Similar results were also reported by Meena *et al.*, (2006), Yadav *et al.* (2011) and Bandiwaddar *et al.* (2015). The higher seed

and straw yield arising from Zn enriched FYM further substantiated by the significant and the gradual release and steady supply of nutrients from FYM throughout the growth and development of plants maintained the photosynthetic efficiency and production of metabolites at higher level and later on the translocation of photosynthetic to various sinks resulting into higher seed and straw yield. Similar findings were also reported by Prihar (2002), Kumawat (2003), Patel *et al.* (2012) and Kanwar and Sharma (2014).

The N, P, K, and Zn content in seed and straw of cowpea increased significantly with the application of Zn enriched FYM (Table 5). The Fe content in seed and straw increased significantly with the application of Fe enriched FYM. The significant increase in nutrient content in plant ascribed to the beneficial role of FYM in mineralization of native as well as nutrients in soil through added fertilizers in addition of its own nutrient content which enhanced the available nutrients pool of soil. The favourable conditions for microbial as well as chemical activities due to addition of FYM augmented the mineralization of nutrients and ultimately the available nutrient pool of the soil led to higher uptake of nutrient by plant (Satyajeet *et al.*, 2006). These results are in agreement with those of Vasanthi and Kumarswamy (1999) and Nehra and Grewal (2001) who have reported significant increase in available nutrient pool of soil with the use of vermicompost @ 5 t ha<sup>-1</sup>. When the vermicompost is added to the soil, nitrogenous compounds slowly break down and make steady nutrient supply throughout period of crop, leading to higher nutrient availability to plant Rao (2003) and Yadav *et al.*, (2011).

The increase in Zn content by the application of Zn enriched FYM is due to production of a host of complexing agents which form stable organometallic complexes with Zn that facilitates its efficient utilization by plant resulting in yield and its concentration (Stevenson and Ardakani, 1972). Further, this was also supported by the reduced soil pH due to Zn enriched FYM application which is known to encourage solubility and availability of zinc in soil (Jahiruddin and Crasser, 1993 and Sharma *et al.*, 2000). Higher N, P, K, Zn and Fe content with the use of organic materials have also been reported by Singh and Yadav (2006); Yadav *et al.*, (2001) and Tak *et al.*, (2014).

The protein content in cowpea increased significantly with the application of Zn enriched FYM. The favorable effect of Zn and Fe on synthesis and metabolic process augmented the production of photosynthetic and their translocation to different plant which increased the concentration of N in the seed which directly indicates the content of protein.

Similar results were obtained by Ashoka *et al.*, (2009) in baby corn, Parihar *et al.*, (2014) in mustard and Tak *et al.*, (2014) in green gram.

## Conclusion

Cowpea is a pulse crop role in daily diet. In arid and semi-arid of Rajasthan is having the soil sodicity limitation for this crop. experiment results, increasing RSC and pH of soil reduced the crop yield, nutrient content and uptake by plants. Whereas, application of Zn and Fe enriched FYM improved the crop yield as well as mineral nutrition of the crop. This type of study could be used for enhancing production in high sodicity agricultural fields by application of Farmyard manure enriched with Zn and Fe.

## References

- Amer M. M., ELSaka M. S., Zoghdan M. G. and Khalifa T. H. 2020. Amelioration of salt affected soils and improvement of rice and wheat yields by adding compost and foliar spray of zinc, potassium and compost tea. *Asian Journal of Soil Science and Plant Nutrition* 6(3): 47-60; AJSSPN.59946 ISSN: 2456-9682.
- Ashoka, P.; Anand, and S.R.; Mudalagriyappa and Smitha, R. 2009. Effect of macro and micronutrient with organic on growth, quality, and economics of baby corn [*Zea mays* (L.)] in Tungabhadra command area. *Crop Research, Hisar*. **37**: 15-18.
- Bandiwaddar, T.T. and Patil, B.N. 2015. Response of bread wheat (*Triticum aestivum* L.) to varied levels of iron and zinc enriched with organics in maize-wheat cropping sequence. *Karnataka Journal of Agricultural Sciences*, **28**: 467- 473.
- Ben-Hauyyim, G., Kafkafi, U. and Neuman, G.R. 1987. Role of internal potassium in maintaining growth of cultured citrus cell on increasing NaCl and CaCl<sub>2</sub> concentrations. *Plant Physiology*. **85**: 434-439.
- Brand, J.J. and Beckler, D.W. 1984. Evidence for direct role of calcium in photosynthesis. *Journal of Bioenergetics and Biomembranes*. **16**: 239-249.
- Chauhan, R.P.S.; Bhudayal and Chauhan, C.P.S. 1988. Effect of residual sodium carbonate in irrigation water on soil and bread wheat (*Triticum aestivum* L.). *Indian Journal of Agriculture Science*, **58**: 454-458.
- Doodhwal K., Yadav B.L., Yadav M. and Choudhary M. 2018. Effect of soil sodicity and iron levels on soil properties, yield attributes and yield of cowpea. *International Journal of Agriculture Sciences* ISSN: 0975-3710 & E-ISSN: 0975-9107, Volume 10, Issue 13, pp.-6606-6609.

- Doodhwal K., Yadav B. L., Yadav M., Choudhary M. 2020. Impact of sodium and iron levels on yield, nutrient content and uptake by cowpea (*Vigna unguiculata* L.). *Legume Research- An International Journal*, Volume 43 Issue 4: 558-562.
- Dwivedi, C.A. and Burrows, I. 1979. *Plant Physiology in a decade of Research* CSSRI (ICAR) Karnal. pp. 115.
- Garhwal, R.S., Totawat, K.L. and Yadav, B.K. 2011. Effect of RSC water on physico-chemical properties of Typic Haplustalfs with Marigold (*Tagetes erecta*) and Chrysanthemum (*Chrysanthemum* spp.). *Annals of Arid Zone* **50**: 87-92.
- Jahiruddin, M., Chambers, B.J. Cresser, M.S. and Livesey, N.T. 1992. Effects of soil properties on the extraction of zinc. *Geoderma*, **52**:199-208.
- Jatav, M.K. 2000. Efficiency of zinc and iron application on the performance of wheat in treated sodic soil, irrigated with saline sodic water. *M.Sc. (Ag.) Thesis*, MPUAT, Udaipur.
- Kanwar, A. and Sharma S.R. 2014. Effect of organic and inorganic nutrition on fertility status of soil and yield of vegetable cowpea. *Annals of Arid Zone* **53**: 17-20, 2014.
- Kumar B. and Dhaliwal S. S. 2021. Zinc biofortification of dual-purpose cowpea for enhancing the productivity and nutritional quality in a semi-arid regions of India. *Archives of Agronomy and Soil Science*, 1-15.
- Kumawat, P.D. 2003. Response of barley (*Hordium vulgare* L.) to organic manure and nitrogen fertilization. *Ph.D Thesis*, RAU, Bikaner.
- Mahmood, K. 2011. Salinity tolerance in Barley (*Hordeum vulgare* L.): Effects of varying NaCl,  $K^+ / Na^+$  and  $NaHCO_3$  levels on cultivars differing in tolerance. *Pakistan Journal of Botany*, **43**: 1651-1654.
- Meena B.L., Meena R.L., Ambast S.K., Pandey M. 2014. Impact assessment of agriculture technological interventions in tsunami affected South Andaman- a case study. *Bharatiya Krishi Anushandhan Patrika*. 28(3):141–148.
- Meena B.L., Rattan R.K. and Datta S.P. 2017. Solubility relationships of iron and evaluation of its fertility status in degraded soils. *Commun Soil Sci Plant Anal*. 48:1059–1067.
- Meena, M.C., Patel, K.P. and Rathore D.D. 2006. Effect of Zn and Fe enriched FYM on mustard yield and micronutrients availability in lomay sand soil (Typic Haplustept) of Anand. *Journal of the Indian society of soil Science*, **54**: 495-499.
- Mishra, S.K. 2001. Salt tolerance of different cultivars of chickpea as influenced by zinc fertilization. *Journal of Indian Society of Soil Science*, **49**: 135-140.
- Nehra, A.S. and Grewal, K.S. 2001. Influence of integrated use of organic manure and organic fertilizer on soil properties and yield of wheat. *International Conference of Natural Farming and Ecological Balance*, CCSHAU, Hissar, India. Abstracts of paper, pp. 7-10.

- Panase, V.G. and Sukhatme, P.V. 1967. Statistical method for agricultural workers. ICAR, New Delhi.
- Parihar, N.S. 2002. Effect of sulphar, zinc and organic manures on yield and nutrient uptake of wheat. *Ph. D. Thesis*, RAU, Bikaner.
- Parihar, S., Kameriya, P.R. and Choudhary, R. 2014. Response of mustard (*Brassica juncea* L.) to varying levels of Sulphur and fortified vermicompost under loamy sand soil. *Agricultural Science Digest*, **34**: 296 - 298, 2014.
- Patel, A.R., Sadhu, A.C., Chotalia, R.L. and Patel, C.J. 2012. Effect of vermicompost, diammonium phosphate and zinc sulphate on growth, yield attributes and yield of chickpea (*Cicer arietinum* L.). *International Journal of Agricultural Sciences*, **8**: 527-529.
- Plaut, Z. and Grieve, C.M. 1988. Photosynthesis of salt-stressed maize as influenced by Ca : Na ratios in the nutrient solution. *Plant and Soil*, **105**: 283-286.
- Purushottam, B. K., Puhup, C. S. K., Kumar, and Sodi, B. 2018. Effect of irrigation scheduling and zinc application on chlorophyll content, zinc content, uptake and yield of chickpea (*Cicerarietinum* L.). *J. Pharmacognosy Phytochem.* 7 (1): 1834-1837.
- Raj, A. B. ., & Raj, S. K. 2019. Zinc and boron nutrition in pulses: A review. *Journal of Applied and Natural Science*, **11**(3), 673-679.
- Rao, S.S., 2003. Nutrient balance and economics of integrated nutrient management in groundnut (*Arachishypogheal*) - mustard (*Brassicajuncea* L.) cropping sequence. *Madras Agriculture Journal*, **90**:465-471.
- Reddy T.Y. and Reddy Sankara G.H. 2010. Principles of Agronomy. Kalyani Publication, 191-241.
- Satyajeet, Narwal, R.K. and Yadav, V.K. 2006. Effect of vermicompostbiofertilizer and inorganic fertilizer on growth, yield and water use efficiency on pearlmillet-mustard cropping sequence. *Haryana Journal of Agronomy*, **22**: 113-118.
- Sharma, G.K. 2003. Role of soil applied gypsum in reducing toxic effects of boron and SAR of irrigation water on wheat under sodic soil. *Ph.D. Thesis*, RAU, Bikaner.
- Sharma, M.P., Bali, S.V. and Gupta, D.K. 2000. Crop yield and properties of Inceptisol as influenced by residue management under rice-wheat cropping sequence. *Journal of the Indian Society of Soil Science*, **48** : 506 - 509.
- Singh P. K., Pal V., Pandey M. and Singh A. K. 2005. Effect of irrigation water containing RSC on the yield, chemical composition and uptake of nutrients by periwinkle leaves, *Indian Journal of Agricultural Research*, **39**: 116- 121.
- Stevenson, F.J. and Ardakani, M.S. 1972. Micronutrients in Agriculture, *American Society or Agronomy*, Madison, Wisconsin, USA.

- Tak, S., Sharma, S.K. and Reager, M.L. 2014. Effect of vermicompost and zinc on yield attributes, yield and quality of green gram [*Vignaradiatavar. Aureus* (L.) wilczek] in arid western Rajasthan. *International Journal of Agricultural Sciences*, **10**: 138-141.
- Yadav, B.S. 2001. Effect of organic manure and inorganic fertilizers on yield of wheat. *International Conference on Natural Farming and Ecological Balance*, CCS HAU, Hissar, India, March. 7-10, pp. 132.
- Yadav M, Yadav B. L., Yadav S. L., Doodhwal K., Yadav G. N. and Yadav G. 2019. Performance of mungbean (*VignaRadiata* L. Wilczek) under different RSC water and sources of Zinc. *International Journal of Agriculture Sciences* ISSN: 0975-3710 & E-ISSN: 0975-9107, Volume 11, Issue 3, pp.-7857-7859.
- Yadav, R.S., Patel, A.M., Dodia, I.N., Aglodiya, A.V., Patel, G.A. and Augustine, N. 2011. Agronomic bio-fortification of wheat (*Triticum aestivum* L.) through iron and zinc enriched organics. *Journal of Wheat Research*, **3**: 46-51.

**Table 1: Composition of irrigation water**

RSC mmol L <sup>-1</sup>	EC (dSm <sup>-1</sup> )	SAR	Ionic composition (mmol L <sup>-1</sup> )						
			Na <sup>+</sup>	Ca <sup>2+</sup>	Mg <sup>2+</sup>	CO <sub>3</sub> <sup>2-</sup>	HCO <sub>3</sub> <sup>-</sup>	Cl <sup>-</sup>	SO <sub>4</sub> <sup>2-</sup>
2.0 Base water	1.31	8.2	10.0	1.5	1.5	1.0	4.0	6.5	1.5
4.0	3.14	16.7	26.4	2.5	2.5	2.0	7.0	11.4	11.0
6.0	3.14	16.7	26.4	2.5	2.5	2.0	9.0	10.5	9.9

**Table 2: Effect of different sodic water and sources of Zn and Fe on seed and straw yield of cowpea**

Treatments	Seed yield (g/pot)	Straw (g/pot)
<b>Sodic water (RSC-mmol/L)</b>		
W <sub>2</sub>	8.65	15.03
W <sub>4</sub>	7.23	12.55
W <sub>6</sub>	4.43	7.65
SEm±	0.12	0.24
CD (P= 0.05)	0.35	0.68
<b>Sources of Zn and Fe</b>		
T <sub>0</sub> (control)	4.83	8.01
T <sub>1</sub> ( ZnSO.7H <sub>2</sub> O)	7.08	11.39

T <sub>2</sub> (FeSO <sub>4</sub> .7H <sub>2</sub> O)	6.23	10.20
T <sub>3</sub> (Zn enriched FYM)	8.16	15.20
T <sub>4</sub> (Fe enriched FYM)	7.55	13.90
SEm <sub>±</sub>	0.16	0.31
CD (P= 0.05)	0.46	0.88

**Table 3: Interactive effect of sodic water and sources of Zn and Fe on seed yield (g/pot) of cowpea**

Treatment	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>
W <sub>2</sub>	6.17	9.05	7.96	10.43	9.65
W <sub>4</sub>	5.16	7.56	6.65	8.71	8.06
W <sub>6</sub>	3.16	4.63	4.08	5.34	4.94
	SEm <sub>±</sub>	0.29			
	CD (P= 0.05)	0.84			

**Table 4: Interactive effect of sodic water and sources of Zn and Fe on straw yield (g/pot) of cowpea**

Treatment	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>
W <sub>2</sub>	10.25	14.58	13.05	19.45	17.79
W <sub>4</sub>	8.56	12.17	10.90	16.24	14.85
W <sub>6</sub>	5.22	7.42	6.64	9.90	9.05
	SEm <sub>±</sub>	0.53			
	CD (P= 0.05)	1.52			

**Table 5: Effect of different sodic water and sources of Zn and Fe on zinc and iron content and uptake in seed and straw of cowpea**

Treatments	Iron content (ppm)		Iron uptake (µg/pot)		Zinc content (ppm)		Zinc uptake (µg/pot)	
	Seed	Straw	Seed	Straw	Seed	Straw	Seed	Straw
<b>Sodic water (RSC-mmol/L)</b>								
W <sub>2</sub>	99.21	112.31	25.61	22.71	33.56	21.67	86.91	171.47
W <sub>4</sub>	93.83	108.71	23.51	17.43	26.31	20.35	68.70	138.59
W <sub>6</sub>	85.15	102.53	20.92	9.50	14.24	18.07	38.20	79.68
SEm <sub>±</sub>	1.75	2.05	0.45	0.34	0.53	0.38	1.28	2.68
CD (P= 0.05)	5.05	5.89	1.30	0.98	1.52	1.10	3.68	7.73

**Sources of Zn and Fe**

T <sub>0</sub> (control)	80.00	93.97	18.27	9.01	12.65	15.49	39.26	76.01
T <sub>1</sub> ( ZnSO.7H <sub>2</sub> O)	89.71	103.15	22.37	16.17	23.12	19.91	64.54	118.64
T <sub>2</sub> (FeSO4.7H <sub>2</sub> O)	91.85	107.52	21.44	13.64	19.67	18.92	58.15	110.74
T <sub>3</sub> (Zn enriched FYM)	97.24	113.41	28.30	23.58	36.74	23.71	80.63	174.07
T <sub>4</sub> (Fe enriched FYM)	104.85	121.20	26.37	20.33	31.34	22.12	80.44	170.11
SEm±	2.26	2.64	0.58	0.44	0.68	0.49	1.65	3.47
CD (P= 0.05)	6.52	7.61	1.68	1.27	1.96	1.42	4.75	9.98

---

UNDER PEER REVIEW