

Original Research Article

Influence of Spacing and Zinc Application on Growth and Productivity of Baby Corn

(Zea mays L.)

Abstract

The present trial was undertaken at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj, Uttar Pradesh during the 2021 summer season (*Zaid*). The test was a Randomized Block Design having three replicates. A total of nine treatments were designed with different levels of spacing and quantity of zinc application. The cultivation land was uniform with sandy loam soil with pH neutral. The soil had low level of organic carbon (0.72%), medium level of available nitrogen (278.28 kg ha⁻¹) and potassium (233.24 kg ha⁻¹) and higher level of phosphorus (27.80 kg ha⁻¹). Amongst all the treatments, T₉ with 60 X 20 cm spacing and 25 kg/ha ZnSO₄ had the maximum plant height (168.13 cm), number of leaves per plant (13.25), dry plant mass (90.96 g/plant), number of cobs per plant (2.37), length of the cob per plant (18.77 cm), cob weight with husk (47.92 g), cob weight without husk (22.70 g). Further, the maximum crop growth rate (30.45 g/m²/day), cob yield with husk (14.63 t/ha), cob yield without husk (5.09 t/ha) and green fodder yield (28.83 t/ha) was observed for T₃ with 40 X 20 cm spacing and 20 kg/ha ZnSO₄.

Keywords: *Baby Corn, Zinc Application, Productivity, Spacing*

1. Introduction

Baby corn has become one of the most sought-after crops at the global level with great processing and export possibilities. Known by different names such as young corn, mini corn or candle corn, baby corn is maize (*Zea mays* L.) cob or ears harvested early from the female flower without any fertilization while the stalks are undeveloped. At present, China and Thailand are the leading producers and its widespread use has increased in India (Singh *et al.*, 2015) as well since it can be grown at any point in the year and marketable as a vegetable crop (Dass *et al.*, 2008). Different regions of Meghalaya, Uttar Pradesh, Maharashtra, Karnataka, Andhra Pradesh and Haryana are the major players in the production of baby corn (Rani *et al.*, 2017).

Based on the endosperm of grain, maize is defined into eight groups of which baby corn is utilized for vegetable purposes. It is the immature maize cob or ear harvested before the

pollination of the flower or when there is surfacing of the baby corn silk.. For better marketability, yellow-coloured cobs with row array being regular having 10-12 cm spacing and 1-1.5 cm broadness is preferential. Globally, this crop is harvested in Thailand, Taiwan, Sri Lanka, Myanmar, Guatemala and South Africa. In India, for a growth area of 8.49 m per hectare, the production and yield rate are 21.28 m per tonnes and 2507 kg per hectare respectively. Diversification to this high-value crop is beneficial to the Indian farmers due to low financial risk and economic wellbeing ((**Pandey et al., 2002**)).

The metabolic reactions in plants are driven by an important mineral, Zinc. Zinc deficiency results in poor production of chlorophyll, carbohydrates, proteins, auxins and hindrance in growth and development of maize (**Marschner, 1995**). It has an indispensable part in regulation of RNA and DNA structure and certain enzymes such as hydrogenase and carbonic anhydrase (**Tisdale et al., 1995**).

The plant yield is affected based on the density population since optimum spacing promotes resourceful usage of resources such as water, soil, nutrients and sunlight (**Monneveux et al., 2005**). Optimal growth is obtained through ideal spacing though the productivity results vary depending on cultivar and environment (**Bruns and Abbas, 2005**). Plants in closed spacing results in overcrowding and poor yield and plant growth (**Gozobenli et al., 2004**).

2. Materials and Methods

The present trial was undertaken using Super Goldy variety at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj, Uttar Pradesh (25.57°N, 87.19°E, 98 m) during the 2021 summer season (*Zaid*). The test was a Randomized Block Design having three replicates. Different levels of spacing and quantity of zinc application were set and nine treatment were obtained- T₁: 15 kg/ha ZnSO₄ + 40 cm x 20 cm, T₂: 20 kg/ha ZnSO₄ + 40 cm x 20 cm, T₃: 25 kg/ha ZnSO₄ + 40 cm x 20 cm, T₄: 15 kg/ha ZnSO₄ + 50 cm x 20 cm, T₅: 20 kg/ha ZnSO₄ + 50 cm x 20 cm, T₆: 25 kg/ha ZnSO₄ + 50 cm x 20 cm, T₇: 15 kg/ha ZnSO₄ + 60 cm x 20 cm, T₈: 20 kg/ha ZnSO₄ + 60 cm x 20 cm, T₉: 25 kg/ha ZnSO₄ + 60 cm x 20 cm.

The cultivation land was uniform with sandy loam soil with pH neutral. The soil had low level of organic carbon (0.72%), medium level of available nitrogen (278.28 kg ha⁻¹) and

potassium ($233.24 \text{ kg ha}^{-1}$) and higher level of phosphorus (27.80 kg ha^{-1}). The fertilizers used included urea, DAP and MOP at basal level during sowing. The data was recorded for the following growth parameters during harvesting: plant height, number of leaves and dry plant mass. The yield parameters included: number of cobs per plant, cob length (cm), cob mass, green cob productivity (t/ha), green fodder yield (t/ha). The collected data was subjected to analysis of variance (ANOVA) using --- (Gomez K.A. and Gomez A.A. 1984).

3. Results and Discussion

3.1 Growth attributes

3.1.1. Plant Height

Amongst all the treatments, the maximum height was observed for T₉ (168.13 cm) with 25 kg/ha zinc application and 60 X 20 cm spacing. Two treatments, T₈ (167.57 cm) and T₆, (167.13 cm) showed comparable results to T₉. Wider spacing resulted in better plant height as there was less competition for nutrients, water and sunlight whereas closed spacing leads to decrease in stem circumference. Overcrowding brings out less light to the plant population and the lower internodes have restricted extension. The present results are in line with that of **Neupane *et al.* (2011)**. Along with spacing, application of zinc was directly proportional to plant's height. Increase in zinc promotes photosynthesis and formation of chlorophyll and carbohydrates; regulation of stomata and enzyme activities. These results are akin to **Arab *et al.* (2018)**. This was also confirmed by **Rakesh Kumar and Bohra (2014)**. The least plant height was reported in the treatment (T₁).

3.1.2. Number of Leaves per Plant

Amongst all the treatments, the maximum number of leaves per plant was observed from T₉ (13.25) with 25 kg/ha zinc application and 60 X 20 cm spacing. Two treatments, T₈ (13.15) and T₆, (13.07) showed comparable results to T₉. The amount of zinc was directly proportional to the number of leaves per plant. With increment in applied zinc, there is promotion in growth and development hormone auxin, cell division and plant metabolism. The current results are in accordance to that of **Tariq *et al.* (2014)**. The results were also in confirmity with **Azab(2015)**. The lowest number of leaves was obtained in the treatment (T₁).

3.1.3. Dry Plant Mass (g/plant)

Amongst all the treatments, the highest dry plant mass was observed for T₉ (90.96 g/plant) with 25 kg/ha zinc application and with 60 X 20 cm spacing. Two treatments, T₈ (90.81

g/plant) and T₆, (90.46 g/plant) showed comparable results to T₉. Smaller spacing results in reduced photosynthesis due to poor sunlight and CO₂ level whereas wider spacing improves the dry matter accumulation from 20 to 80 DAS. These results are related to that of **Sumeria et al. (2007)**. The quantity of zinc applied also had positive effect on the biomass of the plant. The higher quantity of zinc applied inflated the dry plant mass as well. In addition to this, other growth attributes such as plant height, stem girth and weight of roots affected the dry plant mass (**Palai et al. 2018**). **This is also confirmed by Amutham et al.(2019). The lowest dry wieght was obtained in the treatment (T1).**

3.2. Yield Attributes

3.2.1 Number of Cobs per Plant

Amongst all the treatments, the maximum number of cobs per plant was observed for T₉ (2.37) with 25 kg/ha zinc application and **with** 60 X 20 cm spacing. Two treatments, T₈ (2.34) and T₆, (2.17) showed comparable results to T₉. These findings are in accordance to **Anjum et al. (2017)** where it was stated that zinc plays a vital role in regulation of growth hormones such as auxin, promotes synthesis of carbohydrates, protein as well as pollen which in turn results in higher number of cobs. **The results were also in confirmity with Meena et al. (2013).**

3.2.2. Length of Cob per Plant (cm)

Amongst all the treatments, the maximum cob's length was observed for T₉ with 25 kg/ha zinc application and **with** 60 X 20 cm spacing. Two treatments, T₈ and T₆, showed comparable results to T₉. **This was aslo confirmed by Arab et al.(2018).**

3.2.3. Cob weight (g)

3.2.3.1 With husk

Amongst all the treatments, the highest weigh of cob (with husk) was observed for T₉ (47.92 g) with 25 kg/ha zinc application and **with** 60 X 20 cm spacing. Two treatments, T₈ (47.43 g) and T₆, (46.87 g) showed comparable results to T₉.

3.2.3.2 Without husk

Amongst all the treatments, the highest weigh of cob (without husk) was observed for T₉ (22.70 g) with 25 kg/ha zinc application and **with** 60 X 20 cm spacing. Two treatments, T₈ (22.20 g) and T₆, (21.72 g) showed comparable results to T₉.

The yield and its attributes are the sum of both the photosynthates as well as their translocation from source to their sink and this is affected by the presence of minerals.

Minerals aid in production of enzymes and co-enzymes which in turn have an influence on photosynthetic pathways. One such mineral, Zinc, promotes synthesis of chlorophyll, metabolites, growth-related hormones such as auxin. Zinc is having a productive role over photosynthesis and other metabolic activities elevating growth and development of plant and thus, more yield. The results of our present findings are in accordance to those of Arab *et al.* (2018) and Naik *et al.* (2020).

3.2.4. Cob yield (t/ha)

3.2.4.1. With husk

Amongst all the treatments, the maximum cob yield (with husk) was observed for T₉ (14.63 t/ha) with 25 kg/ha zinc application and with 60 X 20 cm spacing. Two treatments, T₈ (14.24 t/ha) and T₆, (13.67 t/ha) showed comparable results to T₉. This is also confirmed by Mona (2015). Treatment (T7) obtained lowest cob yield with husk

3.2.4.2 Without husk

Amongst all the treatments, the maximum cob yield (without husk) was observed for T₃ (5.09 t/ha) with 25 kg/ha zinc application and with 40 X 20 cm spacing. Another treatment, T₂ (4.77 t/ha) showed comparable results to T₃.

The amount of spacing contributes towards competition of nutrients, light and moisture amongst the plant crops. Lowering the plant density has a positive effect on the yield of seeds. With more competitiveness, reduced sunlight hinders the growth at vegetative phase and ultimately the reproductive stage is not achieved which results in low yield. As concluded by Ariraman *et al.* (2021) the reduced yield could be because lesser plants achieved reproductive phase.

Along with spacing, zinc application influenced the yield of seed as well. Zinc directly effects the synthesis of tryptophan and auxin which sequentially impacts the seed yield and its attributes such as cobs per plant, length and weight of seeds. Additionally, zinc contributes towards nutrient metabolism, biological activity and growth and development since enzyme activity is improved as well. This results in more cob yield. This was also similar to findings noticed by Naik *et al.* (2020). The least cob yield without husk was obtained in the treatment (T7).

3.2.5. Green fodder yield (t/ha)

Amongst all the treatments, the highest green fodder yield was observed for T₃ (28.33 t/ha) with 25 kg/ha zinc application and with 40 X 20 cm spacing. Two treatments, T₂ (27.87 t/ha) and T₆, (27.50 t/ha) showed comparable results to T₃.

Zinc has an important role in plant growth and metabolism. It has a positive effect on physiological processes such as synthesis of chlorophyll, carbohydrates and protein as well as forms a vital part in gas exchange through stomata, plant biomass accumulation and starch utilization. The conversion of ammonia to nitrate is regulated via zinc as well. Parellel findings have been noted by **Tamil Amutham *et al.* (2018)**. **This was also confirmed by Mahid *et al.*(2012)**. Treatment (T7) obtained lowest green fodder yield.

UNDER PEER REVIEW

Table 1: Effect of levels of Zinc and Spacing on growth attributes of Baby corn.

Treatments	Plant height (cm)	No. of leaves/plant	Dry weight (g/plant)
1. 15 kg/ha ZnSO ₄ + 40cm x 20 cm	161.40	12.13	86.02
2. 20 kg/ha ZnSO ₄ + 40cm x 20 cm	162.77	12.25	87.18
3. 25 kg/ha ZnSO ₄ + 40cm x 20 cm	164.27	12.70	88.20
4. 15 kg/ha ZnSO ₄ + 50cm x 20 cm	163.70	12.52	87.84
5. 20 kg/ha ZnSO ₄ + 50cm x 20 cm	166.30	12.93	89.83
6. 25 kg/ha ZnSO ₄ + 50cm x 20 cm	167.13	13.07	90.46
7. 15 kg/ha ZnSO ₄ + 60cm x 20 cm	165.80	12.75	88.67
8. 20 kg/ha ZnSO ₄ + 60cm x 20 cm	167.57	13.15	90.81
9. 25 kg/ha ZnSO ₄ + 60cm x 20 cm	168.13	13.25	90.96
F- test	S	S	S
S. Em (±)	0.34	0.09	0.09
C. D. (P = 0.05)	1.03	0.27	0.27

Treatments	No. of cobs/plant	Length of the cob (cm)		Cob weight (g)		Cob Yield(t/ha)		Green fodder Yield (t/ha)
		With husk	Without husk	With husk	Without husk	With husk	Without husk	
1. 1. 15 kg/ha ZnSO ₄ + 40cm x 20 cm	1.24	13.20	7.30	41.02	17.64	12.39	3.77	26.15
2. 2. 20 kg/ha ZnSO ₄ + 40cm x 20 cm	1.43	14.47	7.60	42.05	18.14	14.24	4.77	27.87
3. 3. 25 kg/ha ZnSO ₄ + 40cm x 20 cm	1.72	15.93	7.83	44.15	19.35	14.63	5.09	28.83
4. 4. 15 kg/ha ZnSO ₄ + 50cm x 20 cm	1.61	15.20	7.67	43.47	18.38	10.98	3.25	24.97
5. 5. 20 kg/ha ZnSO ₄ + 50cm x 20 cm	1.95	17.20	8.20	45.57	20.85	12.89	4.09	26.94
6. 6. 25 kg/ha ZnSO ₄ + 50cm x 20 cm	2.17	17.77	8.27	46.87	21.72	13.67	4.53	27.50
7. 7. 15 kg/ha ZnSO ₄ + 60cm x 20 cm	1.83	16.63	7.90	44.69	20.46	9.60	2.83	23.78
8. 8. 20 kg/ha ZnSO ₄ + 60cm x 20 cm	2.34	18.07	8.50	47.43	22.20	10.01	3.02	24.12
9. 9. 25 kg/ha ZnSO ₄ + 60cm x 20 cm	2.37	18.77	8.58	47.92	22.70	11.82	3.43	25.71
F test	S	S	S	S	S	S	S	S
S. Em (±)	0.07	0.31	0.07	0.39	0.34	0.32	0.20	0.31

CD (P = 0.05) 0.20 0.94 0.21 1.17 1.01 0.97 0.61 0.94

Table 2: Effect of levels of Zinc and Spacing on yield attributes and yield of Baby corn

UNDER PEER REVIEW

4. Conclusion

Amongst all the treatments, T₉ having 60 X 20 cm spacing and 25 kg/ha ZnSO₄ had the maximum plant productivity and its attributes such as plant height, number of leaves per plant, dry plant mass and cob's properties. Further, the maximum crop growth rate, yield and yield attributes such as cob yield and green fodder yield. It can be concluded that wider spacing and increased zinc application has positive effect on plant productivity and growth attributes as well.

REFERENCES

- Anjum, S. A., Saleem, M. F., Shahid, M., Shakoor, A., Safeer, M., Khan, I., Farooq, A., Ali, I. and Nazir, U. 2017. Dynamics of soil and foliar applied boron and zinc to improve maize productivity and profitability. *Pakistan Journal of Agricultural Research* **30**(3): 294- 302.
- Arab, G.M., Dina A. Ghazi and El-Ghamry, A.M. 2018. Effect of zinc application on maize grown on alluvial soils. *Journal of Soil Science and Agricultural Engineering, Mansoura University* **9**(9): 419-426.
- Ariraman, R., Selvakumar, S., Mansingh, M.D.I., Karthikeyan, M. and Vasline, Y.A. 2021. Effect of zinc application on growth, yield parameters, nutrient uptake, yield and economics of maize. *Agricultural Reviews* **10**: 1-6.
- Azab, M.E.E. 2015. Increasing Zn ratio in a compound foliar NPK fertilizer in relation to growth, yield and quality of corn plant. *Journal of Innovation of Pharma Biological Science.*, 2:451:468.
- Bruns HA, Abbas H. K. Ultra-high plant populations and nitrogen fertility effects on corn in the Mississippi Valley. *Agronomy Journal* 2005; **97**(4):1136.

- Dass, S., Yadav, V.K., Kwatra, A., Jat, M.L., Rakshit, S., Kaul, J., Parkash, O., Singh, I., Singh, K.P. and Sekhar, J.C. 2008. Baby corn in India. Technical Bulletin, Directorate of Maize Research, Pusa Campus, New Delhi. 6: 1-45.
- Gozobenli H, Kilinc M, Sener O, Konuskan O. Effects of single and twin row planting on yield and yield components in maize. *Asian Journal of Plant Science* 2004; 3:203-206.
- Marschner, M. (1995). Mineral Nutrition of Higher Plants. 2nd Edition Academic Press, London, New York, ISBN-10:0124735436, pp: 200-255
- Mahdhi SS, Hysain B, Singh L. Influence of seed rate, nitrogen and Zinc on fodder maize (*Zea mays*) in temperate condition of western Himalayas. *Indian journal of Agronomy*. 2012; 57(1):85-88.
- Mona EA. Increasing Zn ratio in a compound foliar NPK fertilizer in relation to growth, yield and quality of corn plant. *Journal of innovations in Pharmaceuticals and biological Sciences*. 2015; 2(4):451-4.
- Monneveux. P, Zaidi P. H, Sanchez. C. Population density and low nitrogen affects yield. Associated Traits in Tropical Maize. *Crop Science* 2005; 45(2):103-106.
- Naik, C., Meena, M. K., Ramesha, Y. M., Amaregouda, A., Ravi, M. V. and Dhanoji, M. M. 2020. Morpho-physiological impact of growth indices to Biofortification on growth and yield of sweet corn (*Zea mays* L. *Saccharata*). *Bulletin of Environment, Pharmacology and Life Sciences* 9(3): 37-43.
- Neupane, M. P., Singh, R.K., Rakesh, K. and Anupma, K. 2011. Response of baby corn (*Zea mays* L.) to nitrogen sources and row spacing. *Environment and Ecology* 29(3):1176-1179.
- Palai, J. B., Sarkar, N.C. and Jena, J. 2018. Effect of zinc on growth, yields, zinc use efficiency and economics in baby corn. *Journal of Pharmacognosy and Phytochemistry* 7(2): 1641-1645.

Pandey, A. K., Mani, V. P., Ved, Prakash, Singh, R. D. and Gupta, H. S. 2002. Effect of varieties and plant densities on yield, yield attributes and economics of baby corn (*Zea mays*). *Indian Journal of Agronomy* **47**(2): 221-226.

Rakesh kumar, Bohra JS. Effect of NPKS and Zn application on growth, yield, economics and quality of baby corn. *Archives Agron. Soil Sci.* 2014; 60(9):1193- 1206.

Reena Rani., Sheoran, R.K., Pooja Gupta Soni., Sakshi Kaith and Arpita Sharma. 2017. Baby corn: A wonderful vegetable. *International Journal of Science, Environment and Technology* **6**(2): 1407-1412.

Shivay YS, Prasad R. Effect of source and methods of zinc application on corn productivity, nitrogen and Zinc concentration s and uptake by high quality protein corn (*Zea mays*). *Egyptian Journal of Biology.* 2014; 16:72-78.

Singh, G., Kumar, S., Singh, R. and Singh, S. S. (2015). Growth and yield of Baby Corn as influenced by varieties, spacings and dates of sowing. *Indian Journal of Agriculture Research* **49**(4): 353-357.

Sumeriya, H. K., Singh, P., Nepalia, V., Sharma, V. and Upadhyay, B. 2007. Response of elite sorghum genotypes to planting geometry and fertility levels. *Research on crops* **8**(2): 312- 315.

Tamil Amutham, G., Karthikeyan, R., Thavaprakaash, N. and Bharathi, C. 2019. Agronomic bio-fortification with zinc on growth and yield of babycorn under irrigated condition. *Journal of Pharmacognosy and Phytochemistry* **8**(3): 434-437.

Tariq, A., Shakeel A., Anjum., Mahmood, A., Randhawa., Ullah, E., Naeem, M., Qamar, R., Ashraf, U. and Nadeem, M. 2014. Influence of zinc nutrition on growth and yield behaviour of maize (*Zea mays* L.) hybrids. *American Journal of Plant Sciences* **5**: 2646-2654.

Tisdale, S.L.; Nelson, W.L. and Beaten, J.D. (1984). Zinc In soil Fertility and Fertilizers. Fourth edition,

New York: Macmillan Publishing Company.

UNDER PEER REVIEW