

Original Research Article

IMPACT OF INORGANIC, ORGANIC AND BIOLOGICAL SOURCES OF NUTRIENTS ON GROWTH AND YIELD OF RICE IN AN INCEPTISOL SOIL OF TAMIL NADU (NAVARAI SEASON)

ABSTRACT

A field experiment was carried out during the Navarai season (2022) using rice variety ADT - 37 to assess the impact of inorganic, organic, and biological sources of nutrients on the growth and yield of rice in an Inceptisol soil of Tamil Nadu. The experimental soil was clay loam in texture and had a pH-8.10, EC-0.7 dSm⁻¹, and available NPK of 186.20, 23.30, and 192.25 kg ha⁻¹, respectively. The experiment comprised of nine treatments viz., T₁ – Control, T₂ - 100% RDF, T₃ - 125% RDF , T₄ - 100% RDF + Press mud @ 10 t ha⁻¹ + 2% Zinc Solubilizing Bacteria, T₅ – 100% RDF + Vermicompost @ 6 t ha⁻¹ + 2% Zinc Solubilizing Bacteria, T₆ – 100% RDF + Coirpith Compost @ 6 t ha⁻¹ + 2% Zinc Solubilizing Bacteria, T₇ – 125% RDF + Press mud @ 10 t ha⁻¹ + 2% Zinc Solubilizing Bacteria, T₈ – 125% RDF + Vermicompost @ 6 t ha⁻¹ + 2% Zinc Solubilizing Bacteria, T₉ – 125% RDF + Coirpith Compost @ 6 t ha⁻¹ + 2% Zinc Solubilizing Bacteria. The **results of the experiment** revealed that application of 125% RDF + Vermicompost @ 6 t ha⁻¹ + 2% Zinc Solubilizing Bacteria (T₈) recorded higher values in growth and yield parameters viz., plant height (106.8 cm), Number of productive tillers hill⁻¹ (14.5), Leaf area index (8.02), chlorophyll content (1.285 mg g⁻¹) Number of panicles m⁻² (585), Number of grains panicle⁻¹ (227) Dry matter production (5720 kg ha⁻¹), Grain yield (6293 kg ha⁻¹) and Straw yield (8943 kg ha⁻¹).

Keywords: Rice, **RDF**, **V**ermicompost, **Z**inc solubilizing bacteria, and **G**rain yield.

INTRODUCTION

Rice occupies a pride place among the food crops cultivated and it has not only meets most of the needs on earth but also symbolizes revolution, industrialization, calorie, and earth. It is the cause of the revolution on earth in nutrition as well as on the food

security front (Pradhan and Moharana, 2015). Rice is cultivated worldwide over an area of about 160.68 million ha⁻¹ with an annual production of about 650.19 million tonnes. In India rice is cultivated over an area of about 39.16 million hectares with an annual production of about 85.59 million tonnes and a productivity of 2.20 tonnes ha⁻¹. (Anonymous, 2015) In Tamil Nadu, rice is grown predominantly among the states in India, which is cultivated in an area of 1.90 million hectares and production of 5.79 million tonnes with a productivity of 3.04 t ha⁻¹. However, yield is still lower when compared to the average productivity of rice-producing countries such as Japan (6.50 t ha⁻¹), China (6.70 t ha⁻¹), Egypt (7.50 t ha⁻¹), and Israel (5.50 t ha⁻¹). In India, during the past three decades, intensive agriculture involving high-yielding varieties of rice has led to the heavy withdrawal of nutrients from the soil. Further, the imbalanced use of chemical fertilizers by farmers has also deteriorated soil health and declined soil organic carbon content, which is a threat to sustainability. Nitrogen is commonly the most limiting nutrient for crop production in the major world's agricultural areas and therefore, adoption of good N management strategies often results in large economic benefits to farmers. The use of organic manures in present agriculture is increasing day by day, because of its utility not only in improving the physical, chemical, and biological properties of soil but also in maintaining good soil health and supplying almost all essential plant nutrients for growth and development of crop plants (Olivares *et al.*, 2020a; Montenegro *et al.*, 2021; Olivares *et al.*, 2022). So, it is time to look for measures to stimulate sustainability in the production of rice on a long-term basis. In this context, vermicomposting of different organic wastes shows promising results in enriching the soil with organic matter, nutrients, etc., thereby maintaining soil fertility. Vermicomposting is a simple process of composting with certain species of earthworms to accelerate the process of waste conversion and to get a better end product (Shweta *et al.*, 2010).

Research pieces of evidence clearly show that vermicompost is a potential organic input that would impart consistent environment and soil Physico-chemical properties, biological activity, and nutrient availability which ultimately improves nutrient uptake, growth and yield components, and yields of cereal crops.

To sustain soil health and to provide adequate plant nutrients, an alternative to chemical fertilizers has to be supplied in integration instead of sudden stoppage of chemicals and total conversion to organic would lead to drastic yield gaps during the early

stages of the crop (Singh *et al.*, 2009). The demand for rice is expected to rise by 1.6 percent per year due to the increasing population, while the area of rice cultivation is expected to reduce 40 million ha in the next 15 to 20 years (Shobarani *et al.*, 2010). Hence it is a need to increase the yield and productivity of rice cultivation using reduced inputs and resources to feed the growing population. Moreover, the increasing cost of chemical fertilizers has reduced the profitability of cultivation. Therefore, alternative use of organic manures like green manuring and crop residues along with inorganic fertilizers not only reduces the demand for inorganic fertilizers but also increases the efficiency of applied nutrients due to the favorable effect on physical, chemical, and biological properties of soil (Prasad *et al.*, 1992).

The effectiveness of integrated nutrient management practice depends on the season, soil type, climate, water management, variety, and cropping pattern. Thus, this paper is written to discuss the impact of inorganic, organic, and biological sources of nutrients in various growth and yield parameters of rice.

Materials and methods

The present investigation was carried out in rice variety ADT -37 in farmers field near Gingee taluk, Villupuram district, Tamil Nadu. It represented clay loam texture, and had a pH-8.10, EC-0.7 dSm⁻¹. The experiment carried out to find out the response of rice (*Oryza sativa*) to different sources of nutrients.

Experimental design

The experimental design was used was a randomized block design with three replications. The treatments were different levels of inorganic, organic, and biological nutrients applied to the soil. Treatments details are given below:

T₁ - Control

T₂ - 100 % RDF

T₃ - 125% RDF

T₄ - 100% RDF + Press mud @ 10 t ha⁻¹ + 2% Zinc Solubilizing Bacteria

- T₅** - 100% RDF + Vermicompost @ 6 t ha⁻¹ + 2% Zinc Solubilizing Bacteria
- T₆** - 100% RDF + Coir pith compost @ 6 t ha⁻¹ + 2% Zinc Solubilizing Bacteria
- T₇** - 125% RDF + Press mud @ 10 t ha⁻¹ + 2% Zinc Solubilizing Bacteria
- T₈** - 125% RDF + Vermicompost @ 6 t ha⁻¹ + 2% Zinc Solubilizing Bacteria
- T₉** - 125% RDF + Coir pith compost @ 6 t ha⁻¹ + 2% Zinc Solubilizing Bacteria

Plot size : 5 x 4 m

Design : Randomized **Block Design**

Replications : Three

Results and discussion

Plant height

Among the various treatments, application of 125% RDF + Vermicompost at 6 t ha⁻¹ + 2% Zinc Solubilizing bacteria (T₈) recorded higher plant height over control and other treatments at tillering stage (54.1 cm), panicle initiation stage (88.3 cm) and harvest stage (106.8 cm). The increase in plant height in response **with** favorable effect of vermicompost on plant height could be attributed to the sustained availability of major and micronutrients resulting in increased plant height. This was also confirmed by Murali and Setty (2000), **Kavitha and Subramanian (2007)**. The least plant height were reported in control plots (T₁).

Number of productive tillers hill⁻¹

Among the various treatments, the application of 125% RDF + Vermicompost at 6 t ha⁻¹ + 2% Zinc Solubilizing bacteria (T₈) recorded the highest number of productive tillers hill⁻¹ of 14.50 significantly higher than other treatments. **This may be due to better growth and enhanced photosynthesis in presence of required nutrients in sufficient amount and also owing to better translocation of photosynthesis to sink because of balance nutrients of NPK in vermicompost (Banik and Sharma, 2009). Higher tiller count might be ascribed to adequate supply of zinc that might have increased the uptake and availability of other essential nutrients which resulted in improvement of plant metabolic process (Naik and Das, 2007).** This result is

corroborating with the reports of Kamaleshwaran and Elayaraja (2021). The lowest number of productive tillers hill^{-1} was recorded in the absolute control T_1 of (6.55) respectively.

Leaf area index

Among the various combinations experimented with, the maximum leaf area index of 8.02 was recorded in the treatment T_8 (125% RDF + Vermicompost at 6 t ha^{-1} + 2% Zinc Solubilizing bacteria) and it was on par with T_7 (125% RDF + Press mud at 10 t ha^{-1} + 2% Zinc Solubilizing bacteria) of 7.77. Meena *et al.*, (2006), Dipankar Kar *et al.*, (2007), Mirza *et al.*, (2010), Morteza Siavoshi *et al.*, (2011) reported increase in LAI on addition of organics. Higher LAI could be attributed to an increase in metabolic activity in the plant which could have promoted meristematic activities causing apical growth. Hence, increase N utilization by plants in response to vermicompost application has increased photosynthesis and ultimately leaf area index. The lowest leaf area index was recorded in the absolute control T_1 of 5.30 respectively.

Chlorophyll content

Among the various treatments, the application of 125% RDF + Vermicompost at 6 t ha^{-1} + 2% Zinc Solubilizing bacteria (T_8) recorded the highest chlorophyll content of 1.285 mg g^{-1} and was on par with the application of 125% RDF + Press mud at 6 t ha^{-1} + 2% Zinc Solubilizing bacteria (T_7) of 1.240 mg g^{-1} and significantly higher than other treatments. Vermicompost stimulates the plant growth possibly through supplying nutrients and increasing chlorophyll which consequently improves the photosynthesis (Ievinsh, 2011). These results are also confirmed by the reports of Ramesh *et al.*, (2018). The lowest chlorophyll content was recorded in the absolute control T_1 of 0.650 mg g^{-1} respectively.

Number of panicles m^{-2}

Among the various combinations experimented with, the maximum number of panicles m^{-2} of 585 was recorded in the treatment T_8 (125% RDF + Vermicompost at 6 t ha^{-1} + 2% Zinc Solubilizing bacteria) and it was on par with T_7 (125% RDF + Press mud at 10 t ha^{-1} + 2% Zinc Solubilizing bacteria) of 568. Highest number of panicle m^{-2} may be due to enhanced and continuous supply of nutrients by the vermicompost.

This was also confirmed by Bejbaruah *et al.*, (2013). The lowest number of panicles m^{-2} was recorded in the absolute control T_1 of 423 respectively.

Number of grains panicle⁻¹

Among the various combinations experimented with, the maximum number of grains panicle⁻¹ of 227 was recorded in the treatment T_8 (125% RDF + Vermicompost at 6 t ha^{-1} + 2% Zinc Solubilizing bacteria) of 227 and it was on par with T_8 (125% RDF + Press mud at 10 t ha^{-1} + 2% Zinc Solubilizing bacteria) of 219. **Enhanced and continuous supply of nutrients by the addition of organics leading to better number of grains panicle⁻¹** (Mohandas *et al.*, 2008). This result is similar to the reports of Deytarafder *et al.*, (2016). The lowest number of grains panicle⁻¹ was recorded in the absolute control T_1 of 140 respectively.

Dry matter production

Among the various treatments, the application of 125% RDF + + Vermicompost at 6 t ha^{-1} + 2% Zinc Solubilizing bacteria (T_8) significantly recorded the highest dry matter production of 5720 kg ha^{-1} and was on par with the application of 125% RDF + Vermicompost at 6 t ha^{-1} + 2% Zinc Solubilizing bacteria (T_8) of 5584 kg ha^{-1} . **Increase in DMP due to zinc and organic manures might be due to intermediates / metabolites of decomposing organic manures that hold zinc in the form available to plant or release of Zn mobilizing compounds such as phyto siderophores from roots and induction of polypeptide involved in Zn uptake and translocation to shoot** (Marschner, 1995; Cakmak and Braun, 2004; Ahmad *et al.*, 2012). The higher DMP might be due to increased leaf area due to sustained and enhanced availability of nutrients from the combined source of vermicompost and inorganic fertilizer till the maturity that would have enhanced better biomass production. The lowest dry matter production was recorded in the absolute control T_1 of 4480 kg ha^{-1} respectively.

Grain yield and straw yield

In respect of grain yield and straw yield, among the various treatments observed, application of 125% RDF + + Vermicompost at 6 t ha^{-1} + 2% Zinc Solubilizing bacteria (T_8) recorded significantly highest value of 6293 kg ha^{-1} and 8943 kg ha^{-1} . This might be due to the higher amount of nutrients supplied through

vermicompost along with inorganic fertilizer, which has increased the availability of nutrients in the soil, thus more uptake of nutrients and increased photosynthetic efficiency as evident from increased LAI resulted in higher grain yield and straw yield. This result is similar to the reports of Ramalakshmi *et al.*, (2012) and Kumar *et al.*, (2014). The control plot recorded the lowest grain yield and straw yield.

UNDER PEER REVIEW

Table 1: Impact of inorganic, organic, and biological sources of nutrients on growth and yield parameters of rice

Treatments details	Plant height			No. of Productive tillers hill ⁻¹	Leaf area index	Chlorophyll Content (mg g ⁻¹)	No. of Panicles m ⁻²	No. of Grains Panicle ⁻¹	Dry matter Production (kg ha ⁻¹)	Grain Yield (kg ha ⁻¹)	Straw yield (kg ha ⁻¹)
	TS	PIS	HS								
T ₁ – Control	41.1	58.2	73.4	6.55	5.30	0.650	423.0	140.0	4480	3904	5750
T ₂ – 100% RDF	42.5	64.2	80.3	8.70	6.05	0.879	464.0	162.0	4728	4260	6262
T ₃ – 125% RDF	42.8	72.6	88.2	9.44	6.32	0.913	481.0	171.0	4874	4552	6607
T ₄ – 100% RDF + Press mud at 10 t ha ⁻¹ + 2% ZSB	45.1	74.8	91.7	11.14	6.85	1.102	519.0	197.0	5163	5143	7471
T ₅ – 100% RDF + Vermicompost at 6 t ha ⁻¹ + 2% ZSB	45.9	76.0	92.7	12.08	7.14	1.141	534.0	205.0	5302	5430	7863
T ₆ – 100% RDF + Coirpith compost at 6 t ha ⁻¹ + 2% ZSB	41.5	69.8	86.5	10.13	6.53	1.069	501.0	186.0	5024	4864	7059
T ₇ – 125% RDF + Press mud at 10 t ha ⁻¹ + 2% ZSB	53.2	86.8	105.2	13.72	7.77	1.240	568.0	219.0	5584	6012	8598
T ₈ – 125% RDF + Vermicompost at 6 t ha ⁻¹ + 2% ZSB	54.1	88.3	106.8	14.50	8.02	1.285	585.0	227.0	5720	6293	8943
T ₉ – 125% RDF + Coirpith compost at 6 t ha ⁻¹ + 2% ZSB	50.1	82.1	99.4	12.94	7.45	1.183	552.0	213.0	5444	5715	8237
S.Ed	0.92	1.55	1.87	0.23	0.13	0.02	10.19	3.88	98.25	105.10	150.74
CD (p=0.05)	1.96	3.28	3.97	0.50	0.29	0.04	21.60	8.22	208.28	222.81	319.56

Note: ZSB – Zinc Solubilizing Bacteria, TS – Tillering stage, PIS – Panicle initiation stage, HS – Harvest stage

The combined use of organic fertilizers and microorganisms that promote plant growth are two strategies that contribute to soil biodiversity (Olivares *et al.*, 2020a; Olivares, 2016). There are works in which microorganisms with properties to promote plant growth have been incorporated into the substrates used as fertilizers with satisfactory results. Similarly, Chirinos and Olivares (2013) mention outstanding effects on growth and productivity in crops, when using biofertilizers, significantly reducing the use of chemical fertilizers. State-of-the-art technologies must focus on the sustainable maintenance of a system, in such a way that the use and application of resources must be conceived to conserve the environment (Olivares, 2022; Olivares and Hernández, 2020; Olivares *et al.*, 2020b).

CONCLUSION

Based on the above experimental results, it could be concluded that the cultivation of rice with an application of 125% RDF + Vermicompost at 6t ha⁻¹ + 2% Zinc Solubilizing Bacteria was found to be agronomically and economically viable, practice for augmenting higher production. Hence this can be recommended by the rice-growing farmers in Tamil Nadu.

REFERENCES

Ahmad, H.R., Aziz, T., Hussain, S., Akram, M., Sabir, M., Kashif, S.J.R. and Harifi, M.M. 2012. Zinc enriched farm yard manure improves grain yield and grain Zn concentration in rice grown on saline sodic soil. **Int. J. Agric. Biol.**, 14: 787-792.

Anonymous. 2015. Economics survey of India, Economic Division Ministry of Finance, Government of India, New Delhi pp 45-55.

Banik, P and Sharma, R. 2009. Effect of organic and inorganic sources of nutrients on the winter crops rice cropping system in sub-humid tropics of India, **Archives of Agronomy and Soil Science**, 55(3): 285-294.

Bejbaruah, R., Sharma, R.C. and Banik, P. 2013. Split application of vermicompost to rice (*Oryza sativa* L.): its effect on productivity, yield components, and N dynamics. **Org. Agr.** 3, 123–128.

Cakman, I and Braun, H.J. 2004. Genotypic variation for zinc deficiency. In: Reynolds, M.P., Ortiz Monasterio, J.J. and Menab, A. (eds.) Application of physiology in wheat breeding, 183-199, CIMMYT, Mexico.

Chirinos, J. and Olivares, B. 2013. Biological Effectiveness of Plant Extracts in the in vitro Control of the Phytopathogenic Bacteria Xanthomona. *Revista Multiciencias* 13(2): 115-121.

Deytarafder, K., Banerjee, M. and Malik, G. C. 2016. Effect of nutrient management on productivity of rice under a system of rice intensification (SRI) and its residual effect on Toria. *International journal of bio-resource, environment and agricultural sciences (IJBEAS)*, 2(3): 377-382.

Dipankar Kar, Ghosh, D. and Srivastava, P.C. 2007. Efficacy of evaluation of different zinc-organo complexes in supplying zinc to maize (*Zea mays* L.). *J. Indian Soc. Soil Sci.*, 55(1): 67-72.

Ievinsh, G. 2011. Vermicompost treatment differentially affects seed germination, seedling growth and physiological status of vegetable crop species, *Plant Growth Regulation*, 65(1): 169-181.

Kamaleshwaran, R. and D. Elayaraja. 2021. Influence of vermicompost and FYM on soil fertility, rice productivity, and its nutrient uptake. *International Journal of Agriculture and Environmental Research*, 7(4): 575-583.

Kavitha, R. and Subramanian, P. 2007. Effect of enriched municipal solid waste compost application on growth, plant nutrient uptake and yield of rice, *J. Agron.*, 6(4): 586-592.

Kumar, A., Meena, R. N., Yadav, L. and Gilotia, Y. K. 2014. Effect of organic and inorganic sources of nutrient on yield, yield attributes and nutrient uptake of rice. Rice, Cv. Prh-10. *International Quarterly Journal of Life Science*, 9(2): 595-597.

Marchner, H. 1995. Mineral nutrition of higher plants. *Academic, San Diego*, 889.

Meena, M.C., Patel, K.P. and Rathod, D.D. 2006. Effect of Zn and Fe enriched FYM on mustard yield and micronutrient availability in loamy sand soil (Typic Haplustept) of Anand. *J. Indian Soc. Soil Sci.*, 54(4): 495-499.

Mirza, H., Ahamed, K.U., Rahmatullah, N.M., Akhtar, N., Nahar, K. and Rahman, M.L. 2010. Plant growth characters and productivity of wetland rice (*Oryza sativa* L.) as affected by application of different manures. **Emir. J. Food Agric.**, 22(1): 46-58.

Mohandas, S. and Appavu, K. 2008. Direct and residual effect of combined application of basic slag with green leaf manures on soil available nutrients and yield of rice. **Madras Agric. J.**, 87(1-3): 53-56.

Montenegro, E; Pitti, J; Olivares, B. 2021. Identification of the main subsistence crops of the Teribe: a case study based on multivariate techniques. **Idesia.**, 39 - 3, pp. 83 - 94. <http://dx.doi.org/10.4067/S0718-34292021000300083>

Morteza Siavozhi, Alireza Nasiri and Shankar L. Laware. 2011. Effect of organic fertilizers on growth and yield components in rice (*Oryza sativa* L.). **J. Agrl. Sci.**, 3(3): 217-224.

Murali, M. K., and Setty, R. A. 2000. Effects of levels of NPK, vermicompost, and growth regulators (Triaccontanol) on growth and yield of scented rice. **Mysore Journal of Agricultural Sciences**, 34(4): 335-339.

Naik, S.K. and Das, D.K. 2007. Effect of split application of zinc on yield of rice (*Oryza sativa* L.) in an Inceptisol. **Arch. Agron. Soil Sci.**, 53: 305-313.

Olivares, B. 2016. Description of soil management in agricultural production systems of sector hammock in Anzoátegui, Venezuela. **La Granja: Revista de Ciencias de la Vida.**, 23(1): 14–24. <https://doi.org/10.17163/lgr.n23.2016.02>

Olivares, B. 2022. Determination of the potential influence of soil in the differentiation of productivity and in the classification of susceptible areas to banana wilt in Venezuela. **UCOPress: Spain**, pp. 89-111, <https://helvia.uco.es/handle/10396/22355>

Olivares, B., Hernández, R. 2020. Application of multivariate techniques in the agricultural land's aptitude in Carabobo, Venezuela. **Tropical and Subtropical Agroecosystems**, 23(2): 1-12.

Olivares, B., Pitti, J., Montenegro, E. 2020b. Socioeconomic characterization of Bocas del Toro in Panama: an application of multivariate techniques. **Revista Brasileira de Gestao e Desenvolvimento Regional**, 16(3): 59-71.

Olivares, BO., Araya-Alman, M., Acevedo-Opazo, C. *et al.*, 2020a. Relationship Between Soil Properties and Banana Productivity in the Two Main Cultivation Areas in

Venezuela. **J Soil Sci Plant Nutr**, 20 (3): 2512-2524. <https://doi.org/10.1007/s42729-020-00317-8>

Olivares, BO., Calero, J., Rey, JC., Lobo, D., Landa, BB., Gómez, JA. 2022. Correlation of banana productivity levels and soil morphological properties using regularized optimal scaling regression. **Catena**, 208, 105718. <https://doi.org/10.1016/j.catena.2021.105718>

Pradhan S, and Moharana S. 2015. Effect of organic nutrient management on growth rate and crop productivity in sustainable rice-rice system. **Journal Crop and Weed**, 11(2):28-33.

Prasad, R., S.N. Sharma, S. Singh, and R. Lakshmanan. 1992. Agronomic practices for increasing nitrogen use efficiency and sustained crop production. **In: National Symposium for “Resource Management for Sustained Production”**, 25-28 February, Indian society of Agronomy, Rajasthan Agricultural University. pp.8.

Ramalakshmi, C.S., P.C. Rao, T. Sreelatha, M. Madhavi, G. Padmaja, P.V. Rao, and A. Sireesha. 2012. Nitrogen use efficiency and production efficiency of rice under a rice-pulse cropping system with integrated nutrient management. **J. Rice Res.**, 5(1&2): 42-51.

Ramesh, N., Saraswathi, E., Rajesh, G., Babu Rao, K and Rajanikanth, G. 2018. Influence of fertilizers and vermicompost on *Oryza sativa* L. **Research Journal of Life Sciences, Bioinformatics, Pharmaceutical, and Chemical Sciences**, 4(5): 240-249.

Shobarani, N., G.S.V. Prasad, A.S.R. Prasad, B. Sailaja, P. Muthuraman, S. Numeera and B.C. Viraktamath. 2010. **Rice Almanac India**. In: Technical Bulletin No 5, DRR, Rajendranagar, Hyderabad. pp 6-7.

Shweta, B.L. Rahul Kumar Singh, and Verma Deepshikha. 2010. Integrating microbial composting and vermicomposting for effective utilization of by-products of sugarcane processing industries. **Bioremediation J.**, 14(3): 158-167.

Singh, D.K., B.S. Mahapatra, S. Gupta, and R. Singh. 2009. Impact of nutrient management practices in rice-based cropping system on productivity and soil health. **Journal of Eco friendly Agriculture**, 4 (2):122-124.