

Original Research Article

Effect of Nitrogen and Phosphorus on growth and yield of Cowpea [*Vigna unguiculata* L.]

ABSTRACT

A field study was conducted during *the growing* season of 2021 at the crop research farm of the Department of Agronomy at Sam Higginbottom University of Agriculture, Technology, and Sciences, Prayagraj in the North Eastern plains of Eastern Uttar Pradesh with the objective of studying the effects of nitrogen and phosphorus on growth, yield, and economics of cowpea (*Vigna unguiculata* L.) Var. Gomati VU 89 under a randomised block design, comprising of 9 treatments, of which treatments (T₁-T₉) with different combinations of nitrogen and phosphorus, which are replicated thrice. The experimental results revealed that 25 kg N/ha + 50 kg P/ha recorded Maximum plant height (69.02 cm), number of branches/plant (12.33), number of nodules/plant (4.0), highest plant dry weight (15.32 gm), crop growth rate (4.80 g/m²/day), test weight (18.57 g), number of pods/plant (20.13), number of seeds/pod (15.1), seed yield (1243.97 kg/ha), harvest index (32.0%). T₉ 25 kg N/ha + 50 kg P/ha resulted in a higher net return (34.05 x 10³/ha), gross return (63.44 x 10³/ha), and benefit-cost ratio (2.16).

Key words: Economics, Growth, Cowpea, Nitrogen, Phosphorus, Yield

INTRODUCTION

India has become self-sufficient with respect to the production of food grains but still lags behind the production of pulses. Moreover, burgeoning population pressure and increasing degree of protein mal-nutrition aggravated the problem and call for stepping up the pulses production. Pivotal role of pulses in human diet and system of farming all over the world is well known. Pulses are the major and cheap source of lysine, which is a rich quality protein, providing supplement to cereal-based diet.

“Pulses being the important source of dietary protein have unique ability of maintaining and restoring soil fertility through biological nitrogen fixation along with addition of ample amount of plant residues to the soil. In India, the pulses occupy nearly 29.46-million-hectare area with a production of 22.95 million tonnes and productivity of 779kg/ha” (Anonymous,2019). Among the different pulses, cowpea [*Vigna unguiculata* L.] is the important crop grown in arid and semi-arid regions of the state. Cowpea belongs to the family Leguminosae, and commonly referred to as lobia. Cowpea is kharif pulse crop grown for vegetables, grain, forage and green manuring. It has good promise as an alternative pulse crop in dry land farming. It also works as smother crop keeping weed infestation low.

Cowpea is highly responsive to fertilizer and manures. The adequate application of fertilizers has played a significant role in providing nutrients for growth and development of cowpea. Nitrogen plays an important role in physiological processes of plant and act as constituents of protoplasm. Nitrogen vitally associated with activity of each living cell. Nitrogen is essential constituents of compounds like amino-acids, enzyme, co-enzyme and alkaloids. Though, they have the capacity to fix atmospheric nitrogen symbiotically. The nitrogen application at lower dosed in initial stage before the plant develops sufficient root system, have been beneficial as it provides the nitrogen need of young seedling.

“Phosphorus, an important macronutrient, is among the most needed elements for crop production in most tropical soils, which tend to be deficient in phosphorus availability. In plant tissues, it is present in much smaller amount than nitrogen and potassium although it is the key plant nutrient involved in energy transfer in the plant chemical reactions” (Prasad, R 2007) and “in the biosynthesis of chlorophyll, where phosphorus as pyridoxal phosphate must be present for the biosynthesis of chlorophyll. Phosphorus is known to promote the development of roots there by flourishing the nitrogen fixation in legumes. This increased amount of nitrogen fixed might be utilized by the host plant for its own growth” (Rajasree and Pillai 2001).

OBJECTIVES

1. To study the Effect of Nitrogen and Phosphorus on growth and yield of Cowpea.
2. To calculate the economics of different treatment combinations

MATERIALS AND METHODS

A field study was conducted during kharif season of 2021, at Crop research farm of Department of Agronomy at Sam Higginbottom University of Agriculture, Technology, and Sciences, Prayagraj, Uttar Pradesh, India. This area is located at 25° 24' 42" N latitude, 81° 50' 56" E longitude and 98 m altitude above the mean sea level (MSL). To determine the impact of nitrogen and phosphorus on growth and yield of cowpea (*Vigna unguiculata* L.), the experiment was set up using a Randomized Block Design with nine treatments that were reproduced three times. The treatment net plots are 3m x 3m in size. The treatment are

categorized as with recommended dose of potash 20 kg ha⁻¹ through Muriate of Potash, in addition with Nitrogen and Phosphorus when applied in combinations as follows, (T₁) 15kg/ha nitrogen + 30kg/ha phosphorus, (T₂) 15kg/ha nitrogen + 40kg/ha phosphorus, (T₃) 15kg/ha nitrogen + 50kg/ha phosphorus, (T₄) 20kg/ha nitrogen + 30kg/ha phosphorus, (T₅) 20kg/ha nitrogen + 40kg/ha phosphorus, (T₆) 20kg/ha nitrogen + 50kg/ha phosphorus, (T₇) 25 kg/ha nitrogen + 30kg/ha phosphorus, (T₈) 25 kg/ha nitrogen + 40kg/ha phosphorus, (T₉) 25 kg/ha nitrogen + 50kg/ha phosphorus. The cowpea crop was harvested treatment wise at harvesting maturity stage. Growth parameters viz. plant height (cm), number of branches, dry matter accumulation g plant⁻¹ were recorded manually on five randomly selected representative plants from each plot of each replication separately and after harvesting, seeds were separated from each net plot and were dried under sun for three days. After that, the seeds were winnowed and cleaned, and the seed yield per hectare was calculated and expressed in tonnes per hectare. Each net plot's stover yield was measured and expressed in tonnes per hectare after complete drying in the sun for 10 days. The data was generated and analyzed using the Gomez and Gomez statistical approach (1984). The benefit:cost ratio was calculated using the price of seed with straw and the overall cost of crop cultivation.

RESULTS AND DISCUSSIONS

Effect on growth parameters:

Plant height

Plant height measurements increased as crop growth progressed, as seen in Table 1. At harvest the treatment T₉ (25 kg N/ha + 50 kg P/ha) recorded maximum height of 69.02 cm. At harvesting stage maximum plant height was measured in T₉ and treatments T₈ was found statistically at par to T₉. The highest plant height in treatment T₉ may be ascribed due to the continuous supply of nutrients throughout all growth stages with beneficial association between Nitrogen and Phosphorus along with Potassium. Increasing levels of nitrogen application up to 40 kg N ha recorded significantly higher plant height throughout the crop growth period. This was reported by Jalbi *et al.* (2015) and Hamid *et al.* (2011). The probable reason for increase in plant height in the application of 60 kg/ha P might have resulted in increased carbohydrate accumulation and their remobilization to reproductive parts of the plants, being the closest sink and hence, resulted in increased plant growth. It is also critical in biological energy transfer processes that are vital for life and growth of the plant. Beneficial effect of P plant growth reported by Khandelwal *et al.* (2012).

Number of branches/plant

At harvesting stage maximum number of branches (12.3) are produced by T₉ (25 kg N/ha + 50 kg P/ha) and T₆ and T₈ are statistically at par to maximum. The findings of the present study indicate that growth attributes of crop such as number of branches had marked variation under various nitrogen levels. Highest level of nitrogen fertilizer 40 kg ha⁻¹ produced significantly higher growth attributes due to increased plant nutrient availability. These findings are supported by Kaiser *et al.* (2010). The probable reason for increase in branches/plant in the application of 60 kg/ha P might have resulted due to balance increased in phosphorus. In similar way that adequate P levels promote root growth and stimulate branches/tillers of plant and also, the effect of nutrients on productivity of cowpea genotype Pusa Komal was recorded by Sharma *et al.* (2015) in the experiment application of 80kg P₂O₅/ha.

Dry matter accumulation

The treatment T₉ (25 kg N/ha + 50 kg P/ha) recorded maximum dry matter accumulation of 15.32 (g) at the harvesting stage and T₆, T₈ treatments are found statistically at par to maximum

dry matter accumulation. Because all other vegetative qualities contained it, dry matter accumulation is more crucial. Nitrogen is one of the most important nutrients for crop growth and development because it influences dry matter production and photosynthetic efficiency through regulating leaf area development and maintenance. A similar statement was also found by Asaduzzaman (2008), Saini and Thakur (2008) and Kumar *et al.* (2009). Application of phosphorus P 60 kg/ha significantly increased the dry matter yield of cowpea. This is reported by Rakesh Kumar *et al.* (2016).

Yield and Yield Attributes:

Number of pods/plant

The statistical examination of the number of pods per plant revealed a significant effect. Treatment 25 kg N/ha + 50 kg P/ha recorded significant and highest number of pods per plant (19.37). However, 25 kg N/ha + 40 kg P/ha recorded statistical parity with 25 kg N/ha + 50 kg P/ha. This could be attributable to the crop's accelerated nitrogen metabolism and extended moisture retention, especially during the moisture stress stage, which may have aided in bearing a higher number of pods per plant at harvest. These findings are consistent with those of Singh, R.P., and Dasharath Singh (2017), who found that foliar spray increases incremental production in pulses. **Seed yield**

Different combinations of Nitrogen and Phosphorus with Potassium have a substantial impact on seed production. The highest seed yield was obtained with the treatment 25 kg N/ha + 50 kg P/ha (1243.9), however no other treatments are statistically at par with the treatment 25 kg N/ha + 50 kg P/ha. Application of nitrogen and phosphorus improves the growth and development of grain yield through increasing nutrient status in plant and their translocation towards sink. Phosphorus played vital role in energy transformation, metabolic process of plant and better development of root system which resulted in increasing grain yield. Significant improvement due to nitrogen and phosphorus as observed under present investigation are in close conformity with findings of Gawai *et al.* (2006) and Shaik *et al.* (1998).

Straw yield

The application of Nitrogen and Phosphorus had also significantly influenced the straw production of the cowpea crop. Highest straw yield (2656.21 kg/ha) was recorded 25 kg N/ha + 50 kg P/ha however, 20 kg N/ha + 50 kg P/ha, 25 kg N/ha + 40 kg P/ha were found to be statistically on par with 25 kg N/ha + 50 kg P/ha. Application of nitrogen and phosphorus improves the growth and development of straw yield through increasing nutrient status in plant and their translocation towards sink. Phosphorus played vital role in energy transformation, metabolic process of plant and better development of root system which resulted in increasing straw yield. Significant improvement due to nitrogen and phosphorus as observed under present investigation are in close conformity with findings of Gawai *et al.* (2006) and Shaik *et al.* (1998).

Economics:

Among the different combination of nutrient source 25 kg N/ha + 50 kg P/ha recorded higher net return (34.05×10^3 ₹/ha), gross return (63.44×10^3 ₹/ha) and benefit: cost ratio (2.16).

CONCLUSION

Treatment 25 kg N/ha + 50 kg P/ha recorded highest growth parameters and seed yield (1243.97 kg/ha), gross return (63.44×10^3 ₹/ha), net return (34.05×10^3 ₹/ha) and benefit: cost ratio (2.16) which may be more preferable for farmers since it is economically more profitable and hence, can be recommended to the farmers after further trails.

REFERENCES

- Anil Kumar, Reena, Brij Nanadan, jai Kumar and Jamwal B.S. (2008). Effect of Phosphorus and seed rate on growth and productivity of large seeded kabuli chickpea (*Cicer arietinum* L.) in subtropical kandi areas of Jamu and Kashmir. *Journal of Food legumes* **21**(4):231-233.
- Dhakad Anita, R. S Rajput, P.K.Mishra and S.K. Sarwagi 2005. Effect of planting geometry and N management on growth and dry matter production of wheat + chickpea intercropping system. *Annals of agric Res, Newseries* **25**(4):621-623.
- Gawai P. P. and V. S. Pawar 2006. Integrated nutrient management in sorghum (*Sorghum bicolor*) Chickpea (*Cicer arietinum* L.) cropping sequence under irrigated condition. *Indian J. of Agron.*, **51**(1):17-20.
- Hamid, A. 2011. Nitrogen and carbon effect on the growth and yield performance of mungbean (*Vigna radiata* L. Wilczek.) *Journal of Agronomy and Crop Science*. **161** (1):11-16.
- Hamsa, A. and Puttaiah, E.T. 2012. Residual effect of zinc and boron on growth and yield of French bean (*Phaseolus vulgaris* L.)-rice (*Oryza sativa* L.) cropping system. *International Journal of Environmental Sciences*, **3**:167-171.
- Haruna, I.M. and Usman, A. 2013. Agronomic efficiency of cowpea varieties [*Vigna unguiculata* (L.) Walp.] under varying phosphorus rates in Lafia, Nasarawa State. *Asian Journal of Crop Science*, **5**(2): 209-215.
- Jalali, M.N., Choudhary, A.k., Manglr, M.Q., Omari, M.A., Hamayon, H., Ghafari, S.R and Dass. 2015. Response of different nitrogen levels on growth and yield of mungbean (*Vigna radiata* L. Wilczek) in semi-arid region jackson. *Soil chemical analysis*, Prentice Hall private Limited, New Delhi.
- Jamro, Shinde CP, Singh. 2010. Effect of various levels of nitrogen, phosphorus and sulphur on the yield and quality of mustard in blackgram mustard cropping sequence. Department of Soil Science & Agricultural Chemistry, College of Agriculture, Gwalior. *Crop Research Hisar*. **10**(3):265-270.
- Jat, S.R., Patel, B.J., Shivran, A.C., Kuri, B.R. and Jat, G. 2013. Effect of P and S levels on growth and yield of cowpea under rainfed conditions. *Annals of Plant and Soil Research* **15**: 114-117.
- Karikari, B., Arkorful, E. and Addy, S. 2015. Growth, nodulation and yield response of cowpea to phosphorus fertilizer application in Ghana. *Journal of Agronomy*, **14**:234-240
- Kumar, D. and Masuthi, A. 2005. Effect of phosphorus, zinc and boron on fruit set, seed yield and yield and quality of vegetable cowpea [*Vigna unguiculata* (L.) Walp.].
- Kumawat, S.R., Khistriya, M.K., Yadav, S.L. and Kumar, M. 2014. Effect of sulphur and phosphorus on growth and yield attributes on summer green gram [*Vigna radiata* (L.) Wilczek]. *International Journal of Agricultural Sciences*, **10**(2): 770-773.
- Lad, N.G., Patange, M.J. and Dhage, S.J. 2014. Effect of nitrogen and phosphorous levels on growth, yield attributing characters, yield and economics of French bean (*Phaseolus vulgaris* L.). *International Journal of Current Microbiology and Applied Sciences*, **3**(12): 822-827.
- Leelavathi, G.S.N.S., Subbiah, G.V. and Pillai, R.N.2011. Effect of different levels of nitrogen on the yield of greengram (*Vigna radiata* L., Wilczek). *Andhra Agricultural Journal, India* **38** (1): 93-94.
- Magani, I.E. and Kuchinda, C. 2009. Effect of phosphorus fertilizer on growth, yield and crude protein content of cowpea [*Vigna unguiculata*(L.) Walp.] in Nigeria. *Journal of Applied Biosciences*, **23**:1387-1393.

- Meena L, R. K. Singh and R. C. Gautam 2002. Effect of moisture conservation practices, phosphorus and bacterial inoculation of chickpea, under rainfed condition. *Indian J. Agron.*, **47**(3):398-404.
- Meena, L. R., R.K. Singh and R. C. Gautam, 2005. Response of chickpea (*Cicer arietinum* L.) to moisture conservation practices, phosphorus levels and bacterial inoculation under rainfed condition. *International J. Trop. Agric.*, **22**(1-4):49-60.
- Meena, L.R. and Chand, R. 2014. Response of fodder cowpea to varying levels of nitrogen and phosphorus under rainfed condition of rajasthan. *The Indian Journal of Small Ruminants* 20: 121-123.
- Mozumder S.N.2007. Effect of nitrogen and Rhizobium fertilizer on two varieties of summer mungbean. *Indian Journal of Agronomy* **46** (3)500-505.
- Namakka, A., Hamidu, H.A., Lakun, I.H., and Ahmad, S.Y. 2016. Effect of phosphorus fertilizer application on growth and yield of Sampea-17 cowpea variety [*Vigna unguiculata* (L.) Walp.] in Zaria, Nigeria. *African Journal of Agricultural Science and Technology*, **4**(2): 591-595.
- Neeraj Kumar verma, B.K. pandey, 2008. Studies on the effect of fertilizer doses and row spacing on growth and yield of chickpea (*Cicer arietinum* L.) *Agric. Sci. Dig* **28**(2):139-140.
- Owla, R.L., Chavan B.N. and Singh, U. 2007. Effect of plant densities and phosphorus levels on growth and yield of green gram (*Vigna radiata*L.). *Agriculture research*, **28**(1): 95-97.
- Prasad, R.N., Singh, R.K., Yadava, R.B. and Chaurasia, S.N.S. 2013. Growth and nodulation of cowpea [*Vigna unguiculata*(L.) Walp.] as influenced by phosphorus levels and bio-inoculants. *Vegetable Science*, **40**(2): 204-206.
- Rakesh Kumar, Deepak Kumar Rathore, Magan Singh, Parveen Kumar and Anil Khippal. 2016. Effect of Phosphorus and Zinc nutrition on growth and yield of fodder cowpea. *Legume Research*, **39** (2):262-267.
- Raman, V., Ramankrishna, M., Purushotham, K. and Reddy, K.B. 2011. Effect of bio-fertilizers on growth, yield and quality of French bean (*Phaseolus vulgaris*). *Vegetable Science*, **38**(1): 35-38.
- Raut R.S. and R. N. Sable, 2003. Studies on the yield maximization of chickpea cv. Vijay through fertilizer and growth regulator under irrigated condition *Jr. of Maharashtra Agric uni.*, **28**(3);311-312.
- Ravindra Kumar Sharma, Effect of Phosphorus, Sulphur and Phosphate solubilizing bacteria on productivity and nutrient uptake of chickpea. *Ann. Agric. Res. New series*, **32**(3 & 4);116-119.
- Rudreshappa, T.S. and Halikatti, S.I. 2002. Response of green gram to nitrogen and phosphorus levels in paddy fallows. *Karnataka Journals of Agriculture Sciences*, **15**(1): 4-7.
- Saini and Athakur. 2008. Effect of nitrogen, phosphorus and sulphur on the micronutrient content of blackgram. *Crop Research Hisar* **9** (1):54-58.
- Shaik, R.S. and H. B. Mungse, 1998. Influence of planting system, plant population and fertilizer levels on physiological parameters of late sown chickpea. *J. Maharashtra Agric Univ.*, **23**(1)73-74.
- Shekara, B.G., Sowmyalatha, B.S. and Bharatkumar, C. 2012. Effect of phosphorus levels on forage yield of forage cowpea. *Journal of Horticulture Letters*, **2**:12-13.
- Singh, A., Baoule, A.L., Ahmed, H.G., Dikko, A.U., Aliyu, U., Sokoto, M.B., Alhassan, J., Musa, M. and Haliru, B. 2011. Influence of phosphorus on the performance of cowpea [*Vigna unguiculata* (L.) Walp.] varieties in the Sudan Savanna of Nigeria. *Agricultural Sciences*, **2**:313-317.

Shekara, B.G., Lohithaswa, H., Govindappa, C.M. and Pavan, R. 2010. Response of fodder cowpea genotypes to varied levels of phosphorus. *Forage Research*, 36: 91-93.

Yadav P. K., Vishram ram, A.D. Singh and ram Ruchi 20113. Effect of row ratio and fertilizer in chickpea (*Cicer arietinum* L.) and Mustard (*Brassica juncea*) intercropping system. *Indian J. of Agron.* **58**(2);198-202.

UNDER PEER REVIEW

Table 1. Effect of Nitrogen and Phosphorus on growth parameters of cowpea var. 'Gomati (VU 89)' at harvest

S.No	T.No.	Treatments	Plant height (cm)	No. of branches plant ⁻¹	Dry matter accumulation (g plant ⁻¹)
1	T ₁	15 kg N/ha + 30 kg P/ha	62.99	9.80	14.52
2	T ₂	15 kg N/ha + 40 kg P/ha	64.50	10.27	14.66
3	T ₃	15 kg N/ha + 50 kg P/ha	65.90	10.53	14.71
4	T ₄	20 kg N/ha + 30 kg P/ha	65.39	10.40	14.69
5	T ₅	20 kg N/ha + 40 kg P/ha	66.75	11.27	14.81
6	T ₆	20 kg N/ha + 50 kg P/ha	67.45	11.47	15.03
7	T ₇	25 kg N/ha + 30 kg P/ha	66.14	10.80	14.78
8	T ₈	25 kg N/ha + 40 kg P/ha	68.13	12.00	15.24
9	T ₉	25 kg N/ha + 50 kg P/ha	69.02	12.33	15.32
		SEm (±)	0.33	0.29	0.10
		CD (P 0.05)	1.00	0.85	0.31

Table 2. Effect of Nitrogen and Phosphorus on yield and yield attributing characters of cowpea var. 'Gomati (VU 89)'

S. No	T. No	Treatments	No. of pods/plant	No. of seed/plant	Seed Yield (t/ha)	Straw Yield (t/ha)
-------	-------	------------	-------------------	-------------------	-------------------	--------------------

1	T ₁	15 kg N/ha + 30 kg P/ha	16.13	12.13	0.79	1.67
2	T ₂	15 kg N/ha + 40 kg P/ha	16.50	12.50	0.74	1.76
3	T ₃	15 kg N/ha + 50 kg P/ha	17.10	13.77	0.88	2.03
4	T ₄	20 kg N/ha + 30 kg P/ha	17.03	13.29	0.84	1.96
5	T ₅	20 kg N/ha + 40 kg P/ha	18.57	14.43	1.05	2.32
6	T ₆	20 kg N/ha + 50 kg P/ha	18.77	14.63	1.10	2.40
7	T ₇	25 kg N/ha + 30 kg P/ha	17.87	13.87	0.97	2.17
8	T ₈	25 kg N/ha + 40 kg P/ha	19.37	14.70	1.14	2.48
9	T ₉	25 kg N/ha + 50 kg P/ha	20.13	15.13	1.24	2.66
SEm (±)			0.29	0.32	0.03	0.10
CD (P 0.05)			0.89	0.94	0.10	0.30

Table 3. Effect of Nitrogen and Phosphorus on economics of cowpea var. ‘Gomati VU 89’

S.No	T.No.	Treatments	Cost of cultivation [#] (x 10 ³ ₹ ha ⁻¹)	Gross return (x 10 ³ ₹ ha ⁻¹)	Net return (x 10 ³ ₹ ha ⁻¹)	Benefit: Cost ratio
1	T ₁	15 kg N/ha + 30 kg P/ha	28,270.00	64,000.00	35,730.00	1.26
2	T ₂	15 kg N/ha + 40 kg P/ha	28,820.00	59,200.00	30,380.00	1.05
3	T ₃	15 kg N/ha + 50 kg P/ha	29,360.00	70,400.00	41,040.00	1.39

4	T ₄	20 kg N/ha + 30 kg P/ha	28,290.00	67,200.00	38,910.00	1.37
5	T ₅	20 kg N/ha + 40 kg P/ha	28,830.00	84,000.00	55,170.00	1.91
6	T ₆	20 kg N/ha + 50 kg P/ha	29,370.00	88,000.00	58,630.00	1.99
7	T ₇	25 kg N/ha + 30 kg P/ha	28,300.00	77,600.00	49,300.00	1.74
8	T ₈	25 kg N/ha + 40 kg P/ha	28,840.00	91,200.00	62,360.00	2.16
9	T ₉	25 kg N/ha + 50 kg P/ha	29,390.00	99,200.00	69,810.00	2.37

#Data not subjected to statistical analysis.