

Original Research Article

Effect of city-compost on plant growth, fruit yield and quality of strawberry cv. Festival in Nagaland

ABSTRACT

In this experiment, the present study on effect of different doses of city-compost were conducted including the dose of Nitrogen, phosphorous and potassium through chemical fertilizer as control ($T_1 = \text{NPK @ } 100:60:150 \text{ Kg ha}^{-1}$) and it comprises of eight treatments as a total ($T_2 = \text{City-compost @ } 1.25 \text{ t ha}^{-1}$, $T_3 = \text{City-compost @ } 2.5 \text{ t ha}^{-1}$, $T_4 = \text{City-compost @ } 3.5 \text{ t ha}^{-1}$, $T_5 = \text{City-compost @ } 4 \text{ t ha}^{-1}$, $T_6 = \text{City-compost @ } 5 \text{ t ha}^{-1}$, $T_7 = \text{City-compost @ } 5.5 \text{ t ha}^{-1}$, $T_8 = \text{City-compost @ } 6 \text{ t ha}^{-1}$). Under growth parameters T_1 gave the maximum plant spread, leaf area, number of runners and number of fruits per plant (6.33). Under yield parameters fruit size in terms of length was obtained highest in T_4 (4.15 cm) and in diameters T_1 gave the maximum size (2.81 cm). Maximum yield were found under T_1 (81.43 g plant⁻¹). Under quality parameters, total soluble solids (TSS) (8.03 °Brix), non-reducing sugar (2.87 %), total sugar (5.20 %) and vitamin C (41.67 mg/100ml) were recorded maximum in T_4 as compared to other treatments.

Key words: City-compost, fertilizers, Nagaland, growth, yield, quality.

Introduction:

Strawberry (*Fragaria X ananassa* Duch.) is a soft fruit which belongs to the family Rosaceae and genus *Fragaria* is a unique fruit with highly desirable taste, flavour and excellent source of vitamins, potassium, fibre and sugars (Sharma and Sharma, 2004). Compared to other berry fruits, strawberries contain a higher percentage of vitamin C, phenolics and flavonoids (Hakkinen and Torronen, 2000). Though it is a major fruit of the temperate region, with the advent of day neutral cultivars, it grows well in tropical and subtropical regions of the world. In India its cultivation is confined only to hilly tracts of Himachal Pradesh, Uttaranchal, parts of Uttar Pradesh and Kashmir valley, but with the introduction of day neutral cultivars its cultivation has spread to tropical and subtropical regions too. Its cultivation can be extended to other suitable areas having assured irrigation and transport facilities. The North East States, particularly Nagaland, Meghalaya, Arunachal Pradesh, Sikkim and Mizoram provide ample opportunity for successful strawberry cultivation due to its mild climatic condition. Strawberry gives the quickest returns in shortest possible time. The strawberry fruit is the first of the season's home-grown supplies to reach the markets and while the principal demand is for desserts, it is also used for jam and ice-cream, canning and quick freezing (Hughes *et al.* 1969).

It is commercially propagated by runners and generally one plant produces 7-10 runners but under proper management, it can go up to 15 runners per plant. Daylight period of 12 hours or less and moderate temperatures are important for flower-bud formation, but different cultivar may have a different day length and temperature requirement. Strawberry can be grown in any type of soil-poor sand to heavy clay-provided proper moisture, organic matter and drainage is present.

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

The importance of [the](#) study on effect of city-compost has not been done under foothill of Nagaland so far. City-compost is a finely divided peat like biodegradable material with high porosity, aeration, water holding capacity, most available micro and macro nutrients and rich microbial activity which makes it an excellent soil conditioner. Keeping in view, the present study "Effect of city-compost on plant growth, fruit yield and quality of strawberry in Nagaland" [was undertaken to quantify and test the performance strawberry cultivar festival under sub-humid sub-tropical part in Nagaland with the following objectives: to study the effect of various doses of city-compost on plant growth, fruit yield, quality of fruit and to study the economics of various treatments of strawberry cultivation.](#)

Formatted: Font: Not Italic, Highlight

Methods and materials:

The experiment was conducted at Horticultural Experimental Farm, [School of Agriculture Sciences and Rural Development \(SASRD\), Nagaland University, Medziphema Campus, in 2019.](#) The experimental farm is situated in the humid and sub-tropical region with an average rainfall ranging from 2000 to 2500 mm annually, the mean temperature ranges from 21°C to 32°C during summer and rarely goes below 8°C in winter. The experimental plot was laid out in randomized block design with different doses of city-compost along with paddy straw as mulch material and one control (inorganic fertilizers [as NPK](#)) and replicated [three times](#). T₁: Inorganic fertilizer (N₁₀₀: P₆₀: K₁₅₀ Kg ha⁻¹) as control, T₂: City-compost @ 1.25 t ha⁻¹, T₃: City-compost @ 2.5 t ha⁻¹, T₄: City-compost @ 3.5 t ha⁻¹, T₅: City-compost @ 4 t ha⁻¹, T₆: City-compost @ 5 t ha⁻¹, T₇: City-compost @ 5.5 t ha⁻¹, T₈: City-compost @ 6 t ha⁻¹. Growth parameters like plant spread, leaf area, number of runners and yield parameters like days taken to flowering, days taken to fruit set, number of fruits, days taken for ripening, fruit size, weight, yield, marketable fruit yield and projected yield was calculated by the relationships given [in equations 1 and 2 below:](#)

Formatted: Highlight

$$\text{Marketable fruit yield (g/plot)} = \frac{\text{Total fruit yield}}{\text{average number of } \frac{\text{fruits}}{\text{plant}}} \quad (1)$$

$$\text{Projected yield (q/ha)} = \frac{\text{Average yield per plant}}{\text{plant density per hectare}} \quad (2)$$

Mean data obtained during the period of investigation were statistically analyzed by the analysis of variance method (Panse and Sukhatme, 1995). The significant of different source of variance were tested by error mean square, using Fisher Snedecor 'F' test of probability at 0.5 % level of significance.

Results and discussion:

[The data pertaining to the plant growth, development, yield and quality of fruits under the different treatments obtained during the course of investigation are represented in below tables:](#) Plants under T₁ gave the maximum spread (13.53cm and 12.67cm, 14.00cm and 13.37cm, 14.40cm and 13.67cm) at 15, 30 and 45 days [respectively](#) after planting and minimum spread was recorded in T₆. Data pertaining to leaf area presented in Table 1 shows the slight differences among different doses of city-compost. The highest leaf area was

Formatted: Font: Not Italic, Highlight

Formatted: Font: Not Italic, Highlight

recorded in T₁ (27.33 cm², 30.26 cm², 33.88 cm² and 36.1 cm²) at 15, 30, 45 and 60 days after planting (DAP) and the least is found under T₂. The various dose of city-compost did not influence to produce any runners at 60 and 90 DAP. However, at 120 and 150 DAP it was found varying significantly among the treatments. It was observed that of all the treatments T₁ produced maximum runners per plant (4.67 and 5.33) at 120 and 150 DAP respectively. The treatment T₂ was recorded with the minimum number of runners on both the dates of observation. The results are also in agreement with findings of Abadi *et al.*, 2011 who reported on effect of municipal compost application on micro elements concentration in soil and tissue of plant of mint. Lima *et al.*, 2004 in corn confirmed the beneficial effectiveness of municipal solid waste compost and sewage sludge in maintaining plant growth and yield. With regard to plant growth parameters, the application of chemical fertilizers showed highest significant variance, it might be due to the fact that inorganic fertilizers source from Urea, SSP and MOP provides sufficient quantity of NPK which are the major elements for the production of photosynthates and accumulation of chlorophyll in the leaves.

Formatted: Font: Not Italic, Not Superscript/ Subscript

Table 1: Effect of various doses of city-compost on plant growth parameters of Strawberry cv. Festival

Formatted: Indent: Left: 0"

Treatments	Plant Spread (cm)						Leaf area(cm ²)				Number of runners	
	North-South			East-West			15DAP	30DAP	45DAP	60DAP	120DAP	150DAP
	15DAP	30DAP	45DAP	15DAP	30DAP	45DAP						
T ₁	13.53	14.00	14.40	12.67	13.17	13.67	27.33	30.26	33.88	36.18	4.67	5.33
T ₂	12.53	12.97	13.37	11.54	11.93	12.33	19.99	21.05	22.20	23.10	2.33	3.00
T ₃	13.17	13.53	13.93	12.30	12.80	13.27	20.74	21.19	23.34	24.11	2.67	3.33
T ₄	12.93	13.33	13.80	11.90	12.60	13.00	24.47	26.07	27.16	29.39	3.00	3.67
T ₅	12.37	13.03	13.47	11.30	11.87	13.07	21.55	22.27	24.04	25.23	3.33	4.00
T ₆	11.60	12.00	12.47	11.19	11.70	12.10	22.15	23.21	24.26	25.89	3.67	4.67
T ₇	12.07	12.67	13.07	12.07	12.67	13.07	22.92	23.92	24.94	26.04	4.00	5.00
T ₈	12.33	12.83	13.27	11.80	12.40	12.83	23.24	24.84	25.79	26.98	4.33	5.00
Sem±	0.31	0.31	0.30	0.29	0.30	0.30	0.36	0.54	0.62	0.65	0.32	0.36
CD (P=0.5)	0.95	0.94	0.91	0.89	0.91	0.92	1.10	1.63	1.89	1.98	0.96	1.09

Formatted Table

*DAP: Days After Planting

The perusal of the data regarding flowering revealed that the number of days taken to flower showed variation among the different treatments. T₄ took the least duration (76.67days) for flower initiation among all treatments and maximum was in T₁ (91.27). The data regarding days taken to fruit set did not show significant variance among the treatments. In general, T₂ took least duration (15.07 days) and longest period in T₁ (16.93 days). Number of fruits per plant were highest under T₁ (6.33) and least recorded in T₂ (4.00). The results of days taken for fruit ripening under T₁ showed least number of days (24.20) and longest has been recorded in T₂ (28.87). In terms of fruit diameter, the maximum fruit size was obtained from plants under T₁ (2.81cm) and the minimum in T₂ (2.37 cm). The weight of the fruit varied significantly among different treatments was shown in Table 2. Maximum fruit weight was recorded under treatment T₄ (13.47_g) while the minimum in T₂ (8.87_g). The data regarding highest yield per plant was recorded in T₁ (81.43_g/plant) and lowest in T₂ (34.42_g/plant). The increase in yield of strawberry plants in the present experiment could be explained by increase the microbial population resulting from adding city-compost in soils. These microorganisms can produce substances which affect fruit weight and

number of fruits. The data of highest marketable fruit yield were recorded in T₁ (725.67 g/m²) while the lowest was recorded in T₂ (307.89 g/m²). A perusal of the data of projected yield revealed that T₁ (75.82 q/ha) has shown highest and lowest in T₂ (32.89 q/ha). Kavitha and Subramanian (2007) in rice, Kasturi *et al.*, 2011 in fenugreek, Tzortzakis *et al.*, 2012 in pepper confirmed the effect of enriched municipal waste, solid waste on growth and yield.

Table 2: Effect of various doses of city-compost on flower, fruit and yield attributes of Strawberry cv. Festival

Formatted: Indent: Left: 0", First line: 0"

Treatments	Days taken to flowering	Days taken to fruit set	Number of fruits/plot	Days for fruit set to ripening	Fruit diameter (cm)	Fruit weight (gm.)	Yield (g/plot)	Marketable Fruit yield (g/plot)	Projected yield (q/ha)
T ₁	91.27	16.63	6.33	24.20	2.81	13.07	18.43	725.67	75.82
T ₂	83.47	15.07	4.00	28.87	2.37	8.87	34.42	307.89	32.89
T ₃	89.53	16.27	4.93	27.13	2.33	10.47	50.41	454.56	47.38
T ₄	76.67	16.27	5.67	27.33	2.42	13.47	75.78	682.64	70.67
T ₅	78.33	16.47	5.33	26.27	2.59	13.20	70.63	635.89	64.13
T ₆	79.07	16.47	5.53	28.13	2.43	13.00	67.03	603.76	62.12
T ₇	82.60	16.73	5.07	25.47	2.35	12.47	63.67	573.54	58.38
T ₈	83.60	16.73	5.15	27.53	2.59	11.80	59.14	533.09	54.33
Sem±	3.09	0.90	0.39	0.54	0.08	0.84	1.91	10.31	1.31
CD (P=0.5)	9.37	NS	1.20	1.63	0.24	2.54	5.79	31.27	3.97

Formatted Table

A perusal of data related to quality parameters are reported in Table 3. The treatments T₅ and T₇ (4.33 days) were shown maximum storage life under refrigerated conditions and T₂, T₄ and T₈ (3.33 days) were shown lowest shelf life. The fruits of T₆ showed maximum shelf life (2 days) under room conditions and minimum was in T₈ (1.00 days). The data regarding TSS recorded highest in T₄ (8.03 °B) and lowest in T₂ (6.20 °B). The lowest titratable acidity has been recorded in T₂ (0.55 %) and highest in T₁ (0.72 %). The highest reducing sugar percentage was found in T₇ (2.87 %) and minimum in T₂ (2.33 %). The different doses of city-compost had influence on non-reducing sugar content of fruits which showed variation among the different treatments. The highest was recorded in T₄ (2.87 %) while the least was recorded in T₇ (1.30 %). The fruits under T₄ gave the highest total sugar content (5.20 %) and lowest in T₂ (3.93 %). The ascorbic acid content was estimated highest in T₄ (41.67 mg/100ml juice) and T₂ (28.33 mg/100ml juice) recorded lowest. Increased TSS and total sugars at higher levels of compost might have resulted since humic acid played a specific regulatory role and accumulate higher amounts of carbohydrates in the fruits which might have resulted in to higher TSS and sugar content in fruits. Shehata *et al.*, 2011 and Mahaden, 2009 conducted an experiment on strawberry and revealed that organic fertilizers influence significantly on fruit yield and quality parameters.

Table 3: Effect of various doses of city-compost on fruit quality parameters of Strawberry cv. Festival

Treatments	Shelf life in days		Total soluble solids (°B)	Titratable acidity (%)	Reducing sugar (%)	Non-reducing sugar (%)	Total Sugar (%)	Vitamin C (mg/100ml)
	Under refrigerator	Under room condition						
T ₁	4.00	1.67	7.55	0.72	2.63	1.73	4.37	40.00
T ₂	3.33	1.33	6.20	0.55	2.33	1.60	3.93	28.33
T ₃	4.00	1.67	6.46	0.56	2.43	1.63	4.07	30.00
T ₄	3.33	1.67	8.03	0.57	2.33	2.87	5.20	41.67
T ₅	4.33	1.67	8.00	0.59	2.50	1.97	4.4	40.00
T ₆	3.67	2.00	7.78	0.62	2.23	2.37	4.60	38.33
T ₇	4.33	1.33	7.04	0.64	2.87	1.30	4.17	35.00
T ₈	3.33	1.00	6.97	0.67	2.47	1.83	4.30	33.33
<i>Sem</i> ±	0.2	0.21	0.40	0.03	0.10	0.12	0.08	2.84
CD (P=0.5)	NS	NS	1.20	0.10	1.30	0.36	0.24	2.62

Conclusion:

The data regarding growth parameters and quality parameters showed much variation among the different treatments. Under growth parameters T₁ gave the maximum plant spread, leaf area, number of runners and number of fruits per plant (6.33). Maximum yields were also found under T₁ (81.43 g plant⁻¹). Under quality parameters, total soluble solids (TSS) (8.03 °Brix), non-reducing sugar (2.87 %), total sugar (5.20 %) and vitamin C (41.67 mg/100ml) were recorded maximum in T₄ as compared to other treatments.

Formatted: Font: 12 pt, Not Italic, English (U.S.), Highlight

Formatted: Normal, Indent: Left: 0", First line: 0.5", Right: 0"

Formatted: Font: 12 pt, Highlight

Formatted: Font: 12 pt

Formatted: Font: 12 pt, Bold, English (U.S.), Not Superscript/ Subscript

References:

- Abadi, Z. A., Sepanlou, M. G. and Bahmanyar, M. A. (2011). The effect of municipal compost application on the amount of micro elements and their absorption in soil and medicinal plant of mint (*Menthas*). *African Journal of Biotechnology*. 10(77): 17716-17725.
- Hakkine, S. N. and Torronen, A. R. (2000). Content of flavonols and selected phenolic acids in strawberries and *Vaccinium* species: Influence of cultivar, cultivation site technique. *Food research International*. 33:517-524.
- Hughes, H. M., Duggon, J. B. and Banwell, M. G. (1969). Strawberry bulletin no. 95, HMSO 10, 6th edition, Ministry og Agri. Fish Food, United Kingdom.
- Kasthuri, H., Shanthi, K., Shivakumar, S., Rajakumar, S. Son, H. K. and Song, Y. C. (2011). Influence of municipal solid waste compost (MSWC) on the growth and

yield of Green Gram (*Vigna radiate* (L) wilczek), Fenugreek (*Trigonella foenumgraecum* L.) and on the soil quality. *Iran Journal Environmental Health Science Eng.* 8(3): 285-294.

Kavitha, R. and Subramanian, P. (2007). Effect of enriched municipal solid waste compost application on growth, plant nutrient uptake and yield of rice. *Journal of Agronomy.* 6(4):586-592.

Lima, J. S., De queiroz, J. G. E. and Freitas, H. B. (2004). Effect of selected and non-selected urban waste compost on the initial growth of corn. *Resources, conservation and recycling.* 42:309:315.

Mahadeen, A. Y. 2009. Influence of organic and chemical fertilization on fruit yield and quality of plastic-house grown Strawberry. *Jordan journal of Agricultural Sciences.* 5(2):167-169.

Panse, V. C. and Suhatme, P. V. 1995. *Statistical method for agriculture workers.* ICAR, New Delhi. 119-210.

Sharma, R. R. and Sharma, V. P. 2004. *The Strawberry.* ICAR, New Delhi, India.

Shehata, S. A., Gharib, A. A., El-Mogy, M. M. Gawad, K. F. A. and Shalaby, E. A. 2011. Influence of compost, amino and humic acids on the growth, yield and chemical parameters of strawberries. *Journal of Medicinal Plants research.* 5(11): 2304-2309.

Tzortzakis, N., Gouma, S., Dagianta, E., Saridakis, c., Papamichalaki, M., Goumas, D., Manios, T. 2012. Use of fertigation and and municipal solid waste compost for green pepper cultivation. *The Scientific World Journal Volume.* Article Id: 973193:8.