

Evaluation of the performance of parental line and their F₁ hybrids for yield and quality traits in brinjal (*Solanum melongena* L.)

Abstract

Brinjal or eggplant (*Solanum melongena* L.) is one of the most important cultivated solanaceous vegetable, mainly grown for its edible fruits. The present investigation was carried out in order to obtain *per se* performance of parents and their cross combinations for genetic improvement in brinjal. Ten parents were crossed in diallel fashion excluding reciprocal. Half diallel set of crosses and their all possible 45 F₁'s (excluding reciprocals) in brinjal (*Solanum melongena* L.) were evaluated in Randomized Complete Block Design (RBD) with three replications for nineteen yield and yield contributing traits during *Kharif* 2020-21 (Y₁) and 2021-22 (Y₂) at the Main Experiment Station (MES) of Department of Vegetable Science, Acharya Narendra Deva University of Agriculture and Technology, Narendra Nagar, Kumarganj, Ayodhya (U.P.) India. The study revealed that highly significant differences were observed for most of the traits under study. Based on the *per se* performance, the parent P₉ (2.61 kg) registered highest fruit yield per plant followed by P₇ (2.49 kg) in both the year and pooled. The *per se* performance of crosses *viz.*, P₂ x P₉ (3.11 kg), P₄ x P₆ (2.66 kg), P₁ x P₈ (2.55 kg) and P₆ x P₇ (2.42 kg) were produced significantly highest fruit yield per plant than the general mean. These hybrids may be subjected to multi-locational and multi-seasonal trials for their release for commercial cultivation. The findings with respect to salinity tolerance must be correlated to develop the salt tolerant cultivars.

Key word: *Per se* performance, brinjal, half diallel, yield, hybrids.

Introduction

Brinjal or eggplant (*Solanum melongena* L. 2n=24) is one of the most cultivated solanaceous vegetable, which is mainly grown for its edible fruit. In India it is known by many regional name *viz.*, Baigan (Hindi), Badanekai (kannada), Vangi (Marathi), Katharikai (Tamil), Vankai (Telugu) while in worldwide it is popularly known as aubergine (France) or guinea squash. It is the most popular and major fruit vegetable crop in India and many other countries of the world. It is a perennial plant but grown as an annual crop. It is a flexible crop adapted to different agro-climatic regions and can be grown throughout the year. It is an important crop in the tropical regions of world and is being grown commonly in India, China, Turkey, Japan, Italy, Indonesia, Iraq, Syria, Spain and Phillipines.

Its immature fruits are generally used as vegetable and other culinary preparations, unripe fruit is essentially consumed as cooked vegetable in various forms and the dried shoots are used as firewood in rural areas. Brinjal is consumed by many ways like salad, bhaji, stuffed brinjal, bhārtha, pickles etc., has made the brinjal more popular vegetables in India. Its fruits are widely consumed in various culinary preparations and are rich source of protective nutrients (**Hedges and Lister, 2007**) [7].

Globally, India ranked second in vegetable production next to China and contributed 10.80 M ha and 196.26 MT to global vegetable area and production, respectively. In India, brinjal occupies an area of 0.758 million hectares with 13.154 million tonnes of annual production which have the 17.5 tonnes per hectare productivity (**Anonymous, 2021**) [2]. In Uttar Pradesh brinjal is existence in cultivation on an area of 0.080 million hectare with annual production of 2.75 million tonnes with 34.40 tonnes per hectare productivity. In Uttar Pradesh, Agra, Meerut, Lucknow, Kanpur, Aligarh, Chitrakoot and Gorakhpur district share more area and production in the state (**Anonymous 2018**) [1].

The peel of brinjal have a great amount of anthocyanin. A higher anthocyanin (0.16 - 1.04 g/100 g) and glyco-alkaloid level (6.25- 20.50 mg/100 g) low in brinjal which are essential for good fruit quality. These antioxidant act as anti-ageing agent. In particular antioxidant found in brinjal helps prevent skin cancer. Brinjal is also a good source of bone building vitamin K and magnesium as well as heart healthy copper which give nourishment to your scalp and keep it healthy. White brinjal are highly beneficial for regulation of blood sugar levels in human body and also controls absorption of glucose. This makes them the best option for people suffering from diabetes.

A number of cultivars are under cultivation depending upon the yield, consumer preference about the colour, size and shape of the fruit. But it is not possible to have one common cultivar to suit different localities and local preferences. It is therefore required to improve the locally preferred cultivars for yield and adaptation or development of new hybrid combinations. Earlier, egg plant breeding was relied both on mass selection and pure line selection from land races for the development of improved varieties. Brinjal, being native to India and often cross pollinated crop, possesses considerable diversity for plant type, fruit colour, fruit shape, fruit size, yield and other quality traits (Ravali *et al.*, 2017) [12], which offers much scope for improvement through heterosis breeding.

The selection of parents is the important step in heterosis breeding which could combine well and produce desirable hybrids. Several research workers reported the importance of genetic diversity in crops Reddy *et al.*, 2017 [13], Triveni *et al.*, 2017 [17], and

Srivatsava *et al.*, 2019 [14] with this perception, the present investigation was carried out with the objective to study the *per se* performance of parents and hybrids for yield and quality traits.

Materials and Methods

The present investigation was carried out at the main experiment station, Department of Vegetable Science, Acharya Narendra Deva University of Agriculture and Technology, Narendra Nagar (Kumarganj), Ayodhya (UP), India, during *Kharif*, 2020-21 (Y₁) and 2021–22 (Y₂). The experimental farm falls under humid subtropical climate and is located between 24.47° and 26.56°N latitude and 82.12° and 83.58°E longitude at an altitude of 113 m above mean sea level. The experimental materials comprised of ten promising and diverse inbreds and varieties of brinjal selected on the basis of genetic variability from the germplasm stock maintained in the department. The selected parental lines *i.e.*; Balfahava (P₁), Punjab Sadabahar (P₂), NDB-3 (P₃), NDB-2 (P₄), NDB White-1 (P₅), Pusa Kranti (P₆), Pant Samrat (P₇), Mukta Keshi (P₈), NDB Sel-1 (P₉) and Co-2 (P₁₀) were crossed in all possible cross combinations, excluding reciprocals, following diallel mating design during the year, 2019-20 to get 45 F₁'s for the study of mean performance of parental line and their resultant F₁. Emasculation followed by covering of flower with fine cotton was done in the evening and pollination was done around 7-9 AM in the next morning.

The experiments were conducted in a Randomized Complete Block Design (RBD) with three replications to assess the performance of 45 F₁'s hybrids and their 10 parental lines of brinjal. The crop was planted in two rows spaced at 75 cm apart with a plant to plant spacing of 60 cm. The experiments were transplanted on 12 August, 2020 and 17 August, 2021 in salinity stress condition. The soil of the experimental plot was saline alkali with pH more than 8, Sodium Absorption Ratio (SAR) 35.12, Cation Exchange Capacity (CEC) 17.60 c mol (p⁺) kg⁻¹ and the Electrical Conductivity (EC) was 0.31 ds/m.

Observations were recorded for nineteen economic traits including biochemical traits, *viz.* days to 50% flowering, days to first fruit harvest, leaf length (cm), leaf width (cm), plant height (cm), number of primary branches per plant, harvest duration (days), fruit length (cm), fruit equatorial circumference (cm), number of fruits per plant, average fruit weight (kg), marketable fruit yield per plant (kg), unmarketable fruit yield per plant (kg), reducing sugars (%), non-reducing sugar (%), total sugars (%), chlorophyll content (mg/g), total phenol content and total fruit yield per plant (kg). *Per se* performance were evaluated for parents and hybrids following method suggested by **Panase and Sukatme, 1987 [10]** for analysis of variance of experimental for nineteen yield and yield contributing traits.

Result and discussion

The success of any breeding programme depends on the identification of suitable parents and selection of proper breeding methodology for the improvement of a specific trait. The selection of parents with high *per se* performance would be of merit in producing better hybrids and hence the parents selected for crossing programme were evaluated based on their *per se* performance.

Perusal of Table-1 revealed that the pooled analysis of variance divided in the source of variation into environments, replication, environments x replication, genotypes and environments x genotypes. Mean squares due to environment was found highly significant for all the traits under study except primary branches per plant, reducing sugars, non-reducing sugar and total sugars. Mean squares due to replication and environments x replication were found significant for only one traits *i.e.* primary branches per plant in replication and number of fruits per plant in environments x replication under the study. The mean squares due to genotypes and environments x genotypes were found highly significant for all the traits under study during over season (pooled). The results are in corroboration with findings of Srivatsava *et al.*, 2019 [14], Pidigam *et al.*, 2019 [11], Ravali *et al.*, 2017 [12], Reddy *et al.*, 2017 [13] and Triveni *et al.*, 2017 [17].

Among the parents highest *per se* performance (Table-2) for most desirable traits total fruit yield per plant were exhibited by parents P₉ and P₇ in both the season (Y₁, Y₂) and in over season (pooled). The above mentioned genotypes may be used as donor parents in hybridization programme for developing high yielding varieties of respective groups. Some other genotypes exhibiting very high mean performance for the characters other than total fruit yield per plant are also listed in Table-2. In this context, none of the parents found significantly early for days to 50 % flowering and days to first fruit harvest.

For quality traits, most desirable parents P₃ followed by P₁ and P₇ for reducing sugars in both the season (Y₁, Y₂) as well as in over season pooled recorded higher values over the grand mean. Maximum non-reducing sugar among the parents was observed in parent P₁₀ in Y₁, Y₂ and in over season pooled. Maximum total sugars was exhibited by parents P₃ followed by P₁ and P₉ in both the season (Y₁, Y₂) as well as in over season pooled. Parents P₄ and P₈ in Y₁ and in over season pooled while, none of the parents in Y₂ showed significant chlorophyll content among the parents over the grand mean. Parental line P₉ in Y₁ while, the parents P₄ and P₇ in Y₂ and P₄ in over season pooled were exhibited maximum total phenol

content over the grand mean. The above results are in conformity with the findings of earlier workers [Kanchana, *et al.*, 2021 [8], Timmareddygari, *et al.*, 2021 [15], Kumar *et al.*, 2013 [9].

Among the hybrids highest mean performance (Table-2) for most desirable traits total fruit yield per plant were exhibited by cross combination $P_2 \times P_9$ followed by $P_6 \times P_7$, $P_7 \times P_8$, $P_7 \times P_9$ and $P_1 \times P_7$ in Y_1 , $P_2 \times P_9$ followed by $P_6 \times P_7$, $P_7 \times P_8$, $P_7 \times P_{10}$ and $P_8 \times P_9$ in Y_2 while, crosses $P_2 \times P_9$ followed by $P_4 \times P_6$, $P_1 \times P_8$, $P_6 \times P_7$ and $P_7 \times P_{10}$ in over season pooled. Some other crosses exhibiting very high mean performance for the characters other than total fruit yield per plant are also listed in Table-2. In this context, the most desirable cross combination which produced significantly early 50 % flower than the grand mean were $P_6 \times P_7$ followed by $P_3 \times P_7$, $P_8 \times P_9$ in Y_1 , $P_6 \times P_7$ in Y_2 and $P_5 \times P_6$ in over season pooled. Cross combination $P_3 \times P_8$ followed by $P_6 \times P_7$ and $P_8 \times P_9$ in Y_1 , $P_6 \times P_7$ and $P_1 \times P_2$ in Y_2 and $P_8 \times P_9$ followed by $P_5 \times P_6$ and $P_7 \times P_9$ in over season pooled found significantly early for days to first fruit harvest. Increased yield in crosses in accordance with findings of Chaddha *et al.*, 2001 [4]; Bhatt *et al.*, 2004 [3]; Dubey *et al.*, 2014 [5].

Among the crosses (Table-2), quality traits *viz.* reducing sugars, non-reducing sugar, total sugars, chlorophyll content and total phenol content were more anticipated traits would be useful to identify the best hybrid combination for exploitation of superior hybrids. Out of forty five crosses, top three cross combination $P_2 \times P_4$ followed by $P_2 \times P_9$, $P_3 \times P_6$ in Y_1 ; $P_2 \times P_4$ followed by $P_2 \times P_9$, $P_3 \times P_6$ in Y_2 and $P_2 \times P_4$ followed by $P_2 \times P_9$ and $P_8 \times P_{10}$ in over season pooled produced significantly high reducing sugars than the general mean. The cross combination $P_4 \times P_6$ followed by $P_3 \times P_{10}$ and $P_3 \times P_9$ in Y_1 ; $P_3 \times P_9$ followed by $P_3 \times P_{10}$ and $P_4 \times P_6$ in Y_2 and $P_3 \times P_8$ followed by $P_3 \times P_{10}$ and $P_4 \times P_6$ in over season pooled exhibited significantly high non-reducing sugars than the general mean. Cross combination for total sugars were $P_3 \times P_6$ followed by $P_1 \times P_5$ and $P_2 \times P_6$ in Y_1 ; $P_3 \times P_6$ followed by $P_1 \times P_4$ and $P_3 \times P_4$ in Y_2 and $P_3 \times P_6$ followed by $P_4 \times P_4$ and $P_2 \times P_9$ in over season pooled showed significantly highest total sugars than the general mean. Crosses which produced significantly highest chlorophyll content than the general mean were $P_5 \times P_6$ followed by $P_3 \times P_7$ and $P_3 \times P_9$ in Y_1 ; $P_5 \times P_8$ followed by $P_3 \times P_9$ and $P_3 \times P_7$ in Y_2 ; $P_3 \times P_9$ followed by $P_5 \times P_6$ and $P_1 \times P_5$ in over season pooled. The crosses that produced significantly maximum total phenol content than the general mean were $P_3 \times P_8$ followed by $P_1 \times P_8$ and $P_2 \times P_7$ in Y_1 ; $P_1 \times P_7$ followed by $P_2 \times P_3$ and $P_4 \times P_8$ in Y_2 and $P_5 \times P_6$ followed by $P_5 \times P_7$ and $P_2 \times P_{10}$ in over

season pooled. The superiority of the F_1 's than parents is might be due to the additive and dominance both type of gene action and could be exploit for heterosis breeding. The superiority of *per se* performance of parents and F_1 for various traits in seasons and over season (pooled) have also been reported by earlier workers [Kanchana, *et al.* 2021 [8]; Tripathy, *et al.* 2021 [16]; Timmareddygari, *et al.* 2021 [15]; Gadhiya, *et al.* 2016 [6].

Conclusion

It may be concluded from the present study that, on the basis of *per se* performance, Parents P_9 and P_7 along with four F_1 crosses *viz.* $P_2 \times P_9$ followed by $P_4 \times P_6$, $P_1 \times P_8$ and $P_6 \times P_7$ were identified as superior crosses for total fruit yield plant. These parents (P_9 and P_7) may be used in future breeding plan likewise, the best performing F_1 ($P_2 \times P_9$, $P_4 \times P_6$, $P_1 \times P_8$ and $P_6 \times P_7$) may be recommended for multi-locational trials for commercialization of the crop.

References

1. Anonymous. Indian Horticulture Database. National Horticulture Board, Ministry of agriculture, Government of India, Gurgaon. 2018.
2. Anonymous. Horticulture Statistics Division, Department of Agriculture, Co-operation and Farmer's Welfare, Ministry of Agriculture, India. 2020-2021.
3. Bhatt P K, Zapadiya VJ, Sapovadiya MH, Chetariya CP. Estimation of heterobeltiosis and standard heterosis for fruit yield and its attributes in brinjal (*Solanum melongena* L.). Journal of Pharmacognosy and Phytochemistry. 2019; 8(4): 1384-1388.
4. Chadha S, Singh B, Kumar J. Heterosis in brinjal. Karnataka Journal of Agricultural Sciences. 2001; 14(4): 1130-1133.
5. Dubey R, Das,A, Ojha MD, Saha B, Ranjan,A, Singh PK. Heterosis and combining ability studies for yield and yield attributing traits in brinjal (*Solanum melongena* L.). The Bioscan. 2014; 9(2): 889-894.
6. Gadhiya AD, Chaudhari KN, Rai VP and Patel AP. *Per se* Performance of brinjal (*Solanum melongena* L.) for yield and its contributing traits. Advances in Life Sciences. 2016; 5(2):449-458.
7. Hedges LJ, and Lister CE. Nutritional attributes of spinach, silver beet and eggplant. Crop and Food Research Confidential, Report No. 1928. 2007.
8. Kanchana R, Vijayalatha KR, Anitha T, Sandeep G and Paramaguru P. *Per se* Performance of Hybrids and Parents for Yield and Quality in Brinjal (*Solanum melongena* L.). The Pharma Innovation Journal. 2021; 10(10): 14-21.
9. Kumar SR, Arumugam T, Anandakumar CR, Balakrishnan S, Rajavel DS. Heterosis

- expression, interrelationship, direct and indirect effects of component characters on yield in intervarietal crosses of eggplant (*Solanum melongena* L.). African Journal of Biotechnology. 2013; 12(45):6366-6375.
10. Panse VG, Sukhatme PV. Genetics of quantitative characters in relation to plant breeding. Indian Journal of Genetics and Plant Breeding. 1987; 17:318-328.
 11. Pidigam S, Suchandranath BM, Srinivas N, Narshimulu G, Srivani SA, Yadla H. Assessment of genetic diversity in yardlong bean (*Vigna unguiculata* (L.) Walp subsp. sesquipedalis Verdc.) Germplasm from India using RAPD markers. Genetic Resources and Crop Evolution. 2019; 66:1231-1242.
 12. Ravali B, Saidaiah P, Reddy RK, Shivraj N and Geetha A. Study on character association and path analysis in brinjal (*Solanum melongena* L.). Journal of Pharmacognocny and Phytochemistry. 2017; 6(6):393-397.
 13. Reddy RD, Saidaiah P, Reddy KR and Pandravada SR. Mean performance of cluster bean genotypes for yield, yield parameters and quality traits. International Journal of Current Microbiology and Applied Science. 2017; 6(9): 3685-3693
 14. Srivastava S, Saidaiah P, Shivraj N and Reddy KR. Yield and quality based phenotypic evaluation of germplasm of brinjal (*Solanum melongena* L.) under semi-arid conditions. International Journal of Current Microbiology and Applied Science. 2019; 8(7): 415-422.
 15. Timmareddygar S, Saidaiah P, Natarajan S, Geetha A and Komatireddy RR. *Per se* performance of hybrids for yield, yield attributes and quality parameters in brinjal (*Solanum melongena* L.). International Journal of Current Microbiology and Applied Sciences. 2021; 10(2): 32-45.
 16. Tripathy B, Tripathy P, Sahu GS, Dash SK, Pradhan BD, Lenka D and Das S. *Per Se* performances of brinjal (*Solanum melongena* L.) F1 crosses for fruit yield and bacterial wilt tolerance. The Pharma Innovation Journal. 2021; 10(10): 490-494.
 17. Triveni D, Saidaiah P, Reddy RK and Pandravada SR. Mean performance of the parents and hybrids for yield and yield contributing traits in tomato. International Journal of Current Microbiology and Applied Science. 2017; 6(11): 613-619.

Table-1: ANOVA (mean squares) for a set of 10 x 10 diallel crosses for different traits in brinjal during over season pooled.

Source of variation	d.f	Days to 50% flowering	Days to first fruit harvest	Leaf length	Leaf width	Plant height	Primary branches per plant	Crop duration	Fruit length	Fruit equatorial circumference	Number of fruits per plant
Environment	1	163.10**	598.05**	5.56**	15.14**	108.10**	0.50	2581.60**	1.30**	1.65**	20.68**
Replication	2	1.09	0.55	1.96	0.37	47.51	1.11**	11.28	0.62	0.75	0.80
Env. x Rep.	2	0.20	0.07	1.03	0.06	19.36	0.35	0.22	0.31	0.08	3.53**
Genotypes	54	11.41**	14.42**	6.87**	4.77**	157.02**	2.19**	26.47**	28.82**	29.24**	107.19**
Env. x geno.	54	10.52**	8.66*	7.35**	3.52**	175.21**	2.35**	19.36**	12.39**	8.01**	78.01**
Error	216	4.38	4.83	0.91	0.16	36.52	0.16	10.46	0.47	0.39	1.51

Source of variation	d.f	Average fruit weight	Marketable fruit yield per plant	Unmarketable fruit yield per plant	Reducing sugar	Non reducing sugar	Total sugar	Chlorophyll content	Total Phenol content	Total fruit yield per plant
Environment	1	57.43**	0.662**	0.597**	0.001	0.000	0.002	0.157**	0.048**	2.419**
Replication	2	31.28	0.017	0.020	0.014	0.003	0.006	0.076	0.006	0.001
Env. x Rep.	2	12.30	0.009	0.000	0.000	0.001	0.001	0.016	0.001	0.012
Genotypes	54	2923.14**	0.591**	0.077**	0.271**	0.034**	0.315**	0.088**	0.102**	0.793**
Env. X geno.	54	1052.30**	0.418**	0.067**	0.107**	0.021**	0.116**	0.061**	0.112**	0.485**
Error	216	25.52	0.008	0.002	0.003	0.001	0.003	0.005	0.002	0.010

*, ** Significant at 5 per cent and 1 per cent probability levels, respectively.

Table-2 Mean performance, general mean, range, coefficient of variation and critical difference for nineteen characters of diallel set of 45 F₁'s and their 10 parents in Y₁ (2020-21), Y₂ (2021-22) and pooled

Sr. No.	Genotypes	Days to 50% flowering			Days to first fruit harvest			Leaf length (cm)		
		Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
1	P ₁ × P ₂	46.67	45.00	48.33	67.67	64.67	68.50	13.01	13.43	12.69
2	P ₁ × P ₃	47.33	45.67	47.68	70.67	67.67	69.00	13.51	13.50	13.10
3	P ₁ × P ₄	48.00	46.00	49.50	72.67	67.67	68.17	12.84	11.61	12.29
4	P ₁ × P ₅	48.67	45.67	47.33	70.82	69.67	70.67	11.49	14.45	13.89
5	P ₁ × P ₆	51.00	49.00	48.83	71.91	70.33	70.33	13.59	13.34	13.32
6	P ₁ × P ₇	48.67	46.67	49.68	72.19	71.00	70.33	13.39	13.33	11.99
7	P ₁ × P ₈	50.67	48.67	50.00	70.67	69.33	71.33	12.99	13.07	13.02
8	P ₁ × P ₉	47.67	48.33	48.83	69.00	67.67	67.67	13.05	9.50	13.04
9	P ₁ × P ₁₀	49.67	50.33	49.67	70.33	70.33	70.67	13.28	12.91	13.37
10	P ₂ × P ₃	50.33	49.67	51.00	70.73	69.67	72.50	12.20	12.08	11.83
11	P ₂ × P ₄	49.00	47.00	46.83	72.00	69.00	68.33	9.94	9.76	11.38
12	P ₂ × P ₅	46.33	50.33	48.83	69.00	69.33	70.00	14.29	14.61	14.06
13	P ₂ × P ₆	52.33	51.33	49.67	72.01	70.33	71.50	14.36	14.09	13.47
14	P ₂ × P ₇	49.67	48.33	48.50	70.67	69.00	69.91	14.65	13.96	12.72
15	P ₂ × P ₈	49.00	49.67	52.33	72.34	68.67	70.29	11.92	12.19	12.89
16	P ₂ × P ₉	47.00	45.67	47.17	68.33	67.00	69.60	12.83	12.92	13.15
17	P ₂ × P ₁₀	47.00	48.67	49.67	71.67	69.67	70.17	11.82	11.62	12.30
18	P ₃ × P ₄	47.33	46.33	47.00	70.00	67.33	68.17	14.78	15.23	14.14
19	P ₃ × P ₅	48.67	46.33	48.00	71.63	68.33	69.33	16.20	16.63	14.95
20	P ₃ × P ₆	46.67	47.33	48.83	68.63	66.33	68.53	11.87	11.86	12.03
21	P ₃ × P ₇	45.33	48.00	48.50	67.67	66.67	69.33	12.82	13.24	11.69
22	P ₃ × P ₈	45.67	47.00	46.67	65.67	66.00	67.50	10.45	10.77	12.53
23	P ₃ × P ₉	46.67	45.00	48.67	72.33	69.00	70.50	14.65	14.57	14.45
24	P ₃ × P ₁₀	47.00	48.67	49.17	71.51	68.00	69.33	9.74	10.08	12.37
25	P ₄ × P ₅	52.33	49.67	49.33	71.00	69.67	71.00	13.78	13.41	12.66
26	P ₄ × P ₆	50.00	49.33	48.17	71.67	70.67	69.50	12.55	12.24	12.53
27	P ₄ × P ₇	50.67	48.33	47.67	73.00	69.00	70.33	12.08	12.16	11.99
28	P ₄ × P ₈	50.33	48.33	47.83	72.67	69.00	69.50	12.81	13.11	13.95
29	P ₄ × P ₉	52.67	50.00	49.33	71.93	70.33	70.98	12.49	12.33	14.26
30	P ₄ × P ₁₀	49.67	47.67	47.17	70.74	68.00	68.32	11.03	11.55	11.71
31	P ₅ × P ₆	47.67	45.67	45.50	68.05	65.33	66.50	14.56	16.46	14.64
32	P ₅ × P ₇	53.00	51.00	48.33	72.00	69.33	67.50	12.95	12.91	11.68
33	P ₅ × P ₈	47.67	45.67	46.17	67.33	68.33	70.33	16.34	14.20	14.43
34	P ₅ × P ₉	52.00	49.67	48.33	74.82	69.00	70.25	13.79	17.03	13.38
35	P ₅ × P ₁₀	52.33	49.67	51.00	71.78	71.33	71.17	17.51	13.37	13.57
36	P ₆ × P ₇	45.33	44.67	47.33	66.00	63.67	67.67	15.23	15.73	14.14
37	P ₆ × P ₈	50.00	50.00	50.33	69.00	66.33	69.67	14.07	15.53	13.81
38	P ₆ × P ₉	49.00	49.00	49.67	67.19	66.00	69.33	14.33	13.81	13.31
39	P ₆ × P ₁₀	50.33	47.67	50.17	68.31	66.67	69.30	14.58	13.50	13.00
40	P ₇ × P ₈	49.00	46.00	47.83	69.00	65.00	67.87	15.50	15.24	13.13
41	P ₇ × P ₉	49.67	46.00	46.83	68.98	65.00	66.53	14.12	13.35	13.96
42	P ₇ × P ₁₀	47.67	47.00	50.00	67.42	66.33	69.17	13.24	13.60	13.28
43	P ₈ × P ₉	45.33	46.00	46.83	66.03	65.00	66.17	14.51	14.87	15.60
44	P ₈ × P ₁₀	51.00	48.67	50.33	72.67	69.67	72.24	11.27	12.39	13.09
45	P ₉ × P ₁₀	50.00	47.67	50.00	68.65	66.67	69.39	13.66	13.25	15.38
F₁ Hybrid mean		48.93	47.90	48.61	70.19	68.05	69.43	13.34	13.31	13.20
Parents										
1	P ₁	51.67	48.67	50.17	72.33	70.00	71.17	11.95	11.61	11.78
2	P ₂	49.67	49.00	49.33	70.33	69.33	69.83	12.69	12.63	12.66
3	P ₃	53.00	50.00	51.50	68.67	67.33	68.00	12.96	13.57	13.26
4	P ₄	49.00	50.67	49.83	71.67	68.67	70.17	13.33	13.69	13.51
5	P ₅	48.67	48.00	48.33	70.33	67.00	68.67	13.30	13.97	13.64
6	P ₆	49.67	47.33	48.50	69.67	65.67	67.67	10.65	10.71	10.68
7	P ₇	51.33	49.67	50.50	73.33	69.67	71.50	12.97	13.32	13.14
8	P ₈	49.33	48.33	48.83	67.67	67.33	67.50	12.58	12.23	12.41
9	P ₉	49.00	47.00	48.00	71.00	68.00	69.50	13.82	13.61	13.72
10	P ₁₀	52.33	50.67	51.50	75.33	72.00	73.67	11.57	11.42	11.49
Parental mean		50.37	48.93	49.64	71.03	68.50	69.76	12.58	12.67	12.62
Grand mean		49.19	48.08	48.82	70.34	68.13	69.49	13.20	13.19	13.09
CV		4.16	4.33	3.60	3.43	2.93	2.67	6.83	8.17	6.27
CD 5%		3.27	3.33	2.84	3.86	3.19	2.99	1.44	1.72	1.34
Range	Lowest	45.33	44.67	45.5	65.67	63.67	66.17	9.74	9.50	10.68
	Highest	53.00	51.33	52.33	75.33	72.00	73.67	17.51	17.03	15.06

Table-2. Contd...

Sr. No.	Genotypes	Leaf width (cm)			Plant height (cm)			Primary branches /plant		
		Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
1	P ₁ × P ₂	8.46	8.42	7.87	72.63	76.27	78.72	5.13	5.30	5.61
2	P ₁ × P ₃	7.90	7.80	8.06	74.15	73.78	75.00	5.33	5.53	5.70
3	P ₁ × P ₄	9.11	7.25	7.69	72.31	76.26	68.49	6.46	6.58	6.11
4	P ₁ × P ₅	7.50	9.09	8.20	71.15	73.78	71.98	6.07	6.25	5.84
5	P ₁ × P ₆	7.75	7.71	8.14	68.31	70.26	72.06	5.92	6.01	6.17
6	P ₁ × P ₇	7.84	7.64	7.01	92.94	71.36	71.52	5.95	5.86	5.95
7	P ₁ × P ₈	7.54	8.38	7.68	69.50	68.56	86.21	6.16	7.74	6.53
8	P ₁ × P ₉	7.56	9.46	8.81	72.85	69.21	74.98	5.96	6.49	6.41
9	P ₁ × P ₁₀	7.66	7.53	8.29	70.13	90.06	71.50	6.18	5.87	6.04
10	P ₂ × P ₃	7.56	8.43	7.89	74.42	68.29	83.00	5.96	6.47	6.38
11	P ₂ × P ₄	6.03	6.65	7.56	83.77	71.62	76.56	5.92	6.29	5.71
12	P ₂ × P ₅	8.07	8.77	8.34	74.34	95.62	73.55	5.88	7.48	6.40
13	P ₂ × P ₆	9.19	9.17	9.14	71.89	80.48	83.10	6.54	6.69	6.57
14	P ₂ × P ₇	10.78	9.46	8.48	69.47	72.94	74.61	5.88	6.08	6.07
15	P ₂ × P ₈	7.54	8.20	7.98	94.17	93.88	74.92	6.18	7.87	6.90
16	P ₂ × P ₉	9.55	7.92	7.88	76.95	78.06	83.02	4.68	4.68	5.32
17	P ₂ × P ₁₀	7.62	6.84	7.19	72.11	81.53	79.00	5.00	5.18	5.68
18	P ₃ × P ₄	7.77	6.91	7.24	74.26	73.10	75.03	5.86	5.98	5.97
19	P ₃ × P ₅	9.77	9.96	8.81	68.50	68.50	69.15	6.15	6.15	6.17
20	P ₃ × P ₆	9.26	8.24	7.90	78.48	77.21	75.99	3.64	3.71	4.84
21	P ₃ × P ₇	8.28	8.35	7.19	72.16	68.17	88.77	5.86	5.41	5.67
22	P ₃ × P ₈	7.16	7.26	7.67	95.99	77.56	76.96	6.08	6.15	6.01
23	P ₃ × P ₉	11.17	10.27	9.73	84.82	93.76	81.69	4.42	4.57	5.55
24	P ₃ × P ₁₀	6.55	6.72	8.75	69.25	79.57	71.09	4.50	5.23	5.55
25	P ₄ × P ₅	7.38	8.62	8.08	70.44	91.49	84.07	6.92	6.70	6.44
26	P ₄ × P ₆	6.98	8.09	8.82	68.40	72.71	74.38	6.08	6.50	5.59
27	P ₄ × P ₇	7.91	8.81	8.22	72.84	73.97	74.30	5.38	5.57	5.28
28	P ₄ × P ₈	8.36	10.50	9.13	68.90	71.82	73.30	4.96	4.66	5.26
29	P ₄ × P ₉	10.16	11.39	10.58	72.60	76.48	68.73	6.03	6.27	6.21
30	P ₄ × P ₁₀	8.24	8.80	9.09	95.31	72.34	75.64	5.79	6.49	5.07
31	P ₅ × P ₆	7.87	9.20	8.74	66.92	68.97	80.14	6.46	4.82	5.34
32	P ₅ × P ₇	8.63	8.63	7.90	76.15	72.79	84.17	5.69	6.16	6.12
33	P ₅ × P ₈	9.03	9.13	10.15	77.05	88.12	74.20	4.88	7.46	5.94
34	P ₅ × P ₉	7.51	10.78	8.66	75.62	72.35	74.47	3.68	6.55	5.52
35	P ₅ × P ₁₀	12.76	7.66	7.52	78.26	63.57	71.14	5.63	4.59	5.75
36	P ₆ × P ₇	11.52	10.97	8.97	85.19	79.68	73.67	5.30	5.78	5.93
37	P ₆ × P ₈	11.34	8.06	7.99	72.36	71.84	82.22	3.64	4.15	4.77
38	P ₆ × P ₉	9.75	9.75	9.05	76.06	78.94	74.38	4.79	4.81	4.88
39	P ₆ × P ₁₀	8.85	9.84	10.00	79.89	91.60	81.29	4.21	3.64	4.83
40	P ₇ × P ₈	11.06	10.88	9.56	71.43	79.87	85.15	6.37	6.31	6.05
41	P ₇ × P ₉	8.90	8.81	8.34	93.08	89.98	66.34	4.88	4.85	5.66
42	P ₇ × P ₁₀	7.91	9.33	8.98	69.22	75.00	83.12	4.88	5.06	5.38
43	P ₈ × P ₉	9.31	9.38	9.20	79.28	65.76	74.02	3.71	3.83	4.35
44	P ₈ × P ₁₀	8.18	8.22	7.87	72.89	90.09	72.43	4.59	5.22	4.45
45	P ₉ × P ₁₀	8.93	8.80	10.78	70.79	70.99	72.75	3.46	3.80	4.71
F₁ Hybrid mean		8.63	8.17	8.46	75.94	76.77	76.37	5.40	5.71	5.70
Parents										
1	P ₁	7.32	7.80	7.56	81.17	76.95	79.06	5.91	6.43	6.17
2	P ₂	8.32	8.21	8.27	76.22	78.27	77.25	5.88	5.69	5.79
3	P ₃	8.13	8.36	8.24	66.71	70.05	68.38	5.65	5.71	5.68
4	P ₄	7.31	7.13	7.22	72.60	76.90	74.75	5.44	5.62	5.53
5	P ₅	8.56	8.84	8.70	75.56	75.68	75.62	6.32	6.87	6.59
6	P ₆	6.37	6.90	6.64	73.84	77.53	75.68	6.04	7.32	6.68
7	P ₇	6.99	7.28	7.13	82.36	76.81	79.58	5.31	5.06	5.19
8	P ₈	8.17	8.17	8.17	81.66	78.25	79.95	6.33	6.69	6.51
9	P ₉	9.06	8.84	8.95	71.37	74.94	73.15	6.20	6.61	6.41
10	P ₁₀	7.35	7.50	7.43	70.39	73.91	72.15	6.28	7.43	6.86
Parental mean		7.76	7.90	7.83	75.19	75.92	75.55	5.94	6.34	6.14
Grand mean		8.47	8.57	8.35	75.80	76.62	76.22	5.50	5.81	5.78
CV		4.61	5.33	4.09	8.89	7.65	7.03	8.31	5.91	5.78
CD 5%		0.62	0.73	0.57	10.79	9.38	8.68	0.73	0.55	0.53
Range	Lowest	6.03	6.65	6.64	66.71	63.57	66.34	3.46	3.64	4.35
	Highest	12.76	11.39	1.78	95.99	95.62	88.77	6.92	7.87	6.90

Table-2. Contd...

Sr. No.	Genotypes	Crop duration (Days)			Fruit length (cm)			Fruit equatorial circumference (cm)		
		Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
1	P ₁ × P ₂	194.00	198.00	196.17	15.82	16.32	13.69	12.45	11.83	12.82
2	P ₁ × P ₃	193.00	198.00	192.50	13.56	13.56	15.02	10.51	10.90	9.26
3	P ₁ × P ₄	188.67	200.33	196.83	16.58	16.77	16.92	14.71	11.47	9.26
4	P ₁ × P ₅	195.67	193.67	191.83	16.64	16.48	17.84	12.59	13.14	10.82
5	P ₁ × P ₆	189.67	193.67	193.67	15.45	15.71	13.37	13.66	13.36	12.74
6	P ₁ × P ₇	185.33	196.00	193.00	12.29	15.30	14.16	13.96	12.64	14.26
7	P ₁ × P ₈	191.33	190.00	190.33	13.35	12.93	11.90	15.20	13.55	14.67
8	P ₁ × P ₉	191.33	196.00	195.00	15.11	13.90	13.64	12.30	14.36	15.76
9	P ₁ × P ₁₀	194.67	199.67	196.33	13.09	12.99	13.72	16.12	15.69	16.72
10	P ₂ × P ₃	196.00	200.67	194.67	16.97	16.57	11.92	7.75	8.29	10.44
11	P ₂ × P ₄	192.00	197.33	195.83	18.47	18.14	16.98	12.59	12.62	12.53
12	P ₂ × P ₅	193.67	196.00	194.50	13.89	15.53	14.55	12.59	10.75	10.63
13	P ₂ × P ₆	191.67	196.67	194.83	13.26	13.95	15.26	16.44	15.93	15.32
14	P ₂ × P ₇	188.00	198.33	196.50	14.20	13.89	15.27	10.01	12.66	12.62
15	P ₂ × P ₈	190.67	195.67	194.33	17.04	16.81	16.13	14.56	14.56	14.11
16	P ₂ × P ₉	190.00	195.00	191.33	16.72	16.21	14.25	11.63	12.29	13.13
17	P ₂ × P ₁₀	192.00	197.33	194.83	12.74	12.95	13.15	12.94	13.27	14.24
18	P ₃ × P ₄	188.67	194.00	192.67	18.72	17.92	16.52	10.56	11.29	11.79
19	P ₃ × P ₅	191.33	196.33	195.50	18.29	17.64	15.37	9.39	10.05	13.09
20	P ₃ × P ₆	188.33	193.33	194.00	17.62	17.13	17.05	9.94	10.98	9.36
21	P ₃ × P ₇	189.00	194.00	193.00	14.35	13.83	16.15	11.67	11.56	12.07
22	P ₃ × P ₈	198.00	168.67	199.00	13.13	13.47	13.68	11.32	11.61	12.10
23	P ₃ × P ₉	190.67	198.67	195.17	16.70	16.10	14.68	10.96	12.29	14.37
24	P ₃ × P ₁₀	193.67	198.67	194.50	15.34	14.85	14.52	11.95	12.19	11.10
25	P ₄ × P ₅	195.00	200.00	195.33	17.29	16.55	16.80	11.35	11.20	12.88
26	P ₄ × P ₆	191.67	197.00	193.50	18.19	17.46	17.09	11.76	12.06	11.84
27	P ₄ × P ₇	196.67	200.67	197.50	17.07	16.74	14.74	11.60	11.76	12.35
28	P ₄ × P ₈	194.67	199.33	195.67	13.17	13.03	15.88	13.14	12.50	11.53
29	P ₄ × P ₉	192.33	197.00	194.17	16.04	15.29	16.79	12.55	12.23	10.81
30	P ₄ × P ₁₀	194.00	198.00	193.17	13.99	13.99	15.80	12.14	12.13	11.03
31	P ₅ × P ₆	190.67	200.33	194.67	16.37	11.17	12.76	13.98	12.91	12.29
32	P ₅ × P ₇	191.33	196.33	197.17	14.74	14.66	13.90	15.10	15.10	13.31
33	P ₅ × P ₈	196.33	195.67	193.17	11.65	16.37	16.54	13.49	13.46	12.31
34	P ₅ × P ₉	194.00	198.00	195.83	13.82	12.09	13.71	12.78	13.23	12.59
35	P ₅ × P ₁₀	193.33	199.33	197.17	12.09	13.53	15.41	13.14	13.36	12.36
36	P ₆ × P ₇	191.33	196.67	194.17	17.19	19.27	18.73	13.21	12.88	12.32
37	P ₆ × P ₈	195.33	199.33	199.83	12.63	12.93	15.00	12.46	15.96	13.78
38	P ₆ × P ₉	192.33	196.67	195.67	13.02	12.54	12.85	13.99	13.96	13.55
39	P ₆ × P ₁₀	195.67	200.33	196.33	12.89	13.30	14.67	15.99	12.34	12.44
40	P ₇ × P ₈	197.00	201.67	197.83	17.26	16.48	15.23	10.88	10.51	11.33
41	P ₇ × P ₉	192.67	194.67	192.67	12.85	12.85	14.61	12.17	16.61	15.30
42	P ₇ × P ₁₀	189.67	196.67	195.67	12.85	12.67	13.17	16.85	12.11	13.61
43	P ₈ × P ₉	197.00	201.00	198.67	10.89	11.23	11.44	13.89	14.29	13.89
44	P ₈ × P ₁₀	190.33	195.33	194.67	7.72	11.16	12.49	14.74	15.28	14.03
45	P ₉ × P ₁₀	197.33	201.33	197.33	8.92	8.44	10.26	13.88	13.36	13.25
F₁ Hybrid mean		192.58	196.70	195.03	14.67	14.68	14.74	12.78	12.77	12.66
Parents										
1	P ₁	194.33	199.33	196.83	11.05	11.29	11.17	13.82	13.88	13.85
2	P ₂	187.00	194.33	190.67	16.47	16.79	16.63	7.63	7.32	7.47
3	P ₃	193.33	198.00	195.67	17.07	17.20	17.13	7.04	7.44	7.24
4	P ₄	190.00	194.00	192.00	19.20	17.45	18.32	8.49	8.16	8.33
5	P ₅	192.33	198.00	195.17	11.03	10.90	10.96	12.13	11.63	11.88
6	P ₆	190.00	194.33	192.17	13.03	13.63	13.33	15.88	15.16	15.52
7	P ₇	190.33	195.00	192.67	10.88	11.98	11.43	15.78	15.65	15.71
8	P ₈	192.00	197.00	195.50	13.39	12.99	13.19	17.15	16.91	17.03
9	P ₉	191.00	195.33	193.17	14.44	14.32	14.38	17.74	17.99	17.87
10	P ₁₀	189.00	194.00	191.50	7.27	7.14	7.30	12.58	12.52	12.55
Parental mean		190.93	195.93	193.53	13.38	13.36	13.38	12.82	12.66	12.74
Grand mean		192.28	196.56	194.76	13.43	14.44	14.49	12.78	12.75	12.68
CV		1.58	1.70	1.33	4.06	5.24	3.79	5.52	5.22	4.36
CD 5%		4.88	5.93	4.19	0.93	1.21	0.89	1.13	1.06	0.90
Range	Lowest	185.33	168.27	190.33	7.27	7.14	7.3	7.04	7.32	7.24
	Highest	198.00	201.67	199.83	19.20	19.27	18.73	17.74	17.99	17.87

Table-2. Contd...

Sr. No.	Genotypes	Number of fruits /plant			Average fruit weight (g)			Marketable fruit yield per plant (kg)		
		Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
1	P ₁ × P ₂	16.90	16.57	16.24	108.47	108.69	115.56	1.04	1.00	1.29
2	P ₁ × P ₃	20.24	22.58	26.28	80.07	80.44	65.04	1.24	1.21	1.03
3	P ₁ × P ₄	31.20	22.92	27.88	78.12	84.71	69.86	1.48	1.33	1.18
4	P ₁ × P ₅	22.55	28.42	27.06	83.77	77.93	67.06	1.34	1.44	1.12
5	P ₁ × P ₆	23.18	23.18	23.02	78.54	78.42	77.85	1.09	1.09	1.20
6	P ₁ × P ₇	31.69	26.55	22.47	102.87	76.95	92.21	1.81	1.39	1.43
7	P ₁ × P ₈	23.88	28.60	24.27	112.46	95.78	109.72	1.92	1.82	1.87
8	P ₁ × P ₉	23.83	22.88	19.35	76.26	107.39	117.03	1.37	1.93	1.73
9	P ₁ × P ₁₀	31.36	30.93	24.17	58.46	63.07	107.32	1.13	1.19	1.59
10	P ₂ × P ₃	34.63	34.75	32.90	45.75	43.58	56.90	0.94	0.97	0.97
11	P ₂ × P ₄	26.62	25.58	21.24	66.33	65.16	86.82	1.17	1.23	1.13
12	P ₂ × P ₅	21.81	31.62	25.93	111.35	63.77	71.92	1.92	1.45	1.34
13	P ₂ × P ₆	21.83	22.77	26.99	73.69	78.61	78.36	1.20	1.22	1.35
14	P ₂ × P ₇	33.22	22.58	22.57	63.90	111.03	97.40	1.43	1.93	1.64
15	P ₂ × P ₈	21.32	21.52	22.35	79.73	80.39	79.47	1.42	1.45	1.27
16	P ₂ × P ₉	35.88	35.62	33.65	101.30	102.35	102.61	2.40	2.46	2.13
17	P ₂ × P ₁₀	27.84	29.11	26.50	60.59	60.95	86.70	1.16	1.21	1.56
18	P ₃ × P ₄	36.45	36.91	30.37	51.75	51.93	64.10	1.28	1.30	1.34
19	P ₃ × P ₅	18.29	18.26	24.81	81.49	79.53	69.00	1.15	1.18	1.16
20	P ₃ × P ₆	18.27	18.68	26.66	82.38	83.17	64.46	1.21	1.25	1.09
21	P ₃ × P ₇	31.73	26.61	26.61	55.99	63.86	65.9	1.34	1.39	1.28
22	P ₃ × P ₈	19.34	19.76	20.78	85.13	85.65	98.50	1.13	1.19	1.55
23	P ₃ × P ₉	18.26	20.15	20.99	109.15	110.52	92.11	1.44	1.48	1.34
24	P ₃ × P ₁₀	25.26	24.96	29.09	58.95	59.91	61.90	0.83	0.86	1.14
25	P ₄ × P ₅	22.21	22.48	21.90	67.47	68.44	74.08	0.92	0.93	1.18
26	P ₄ × P ₆	25.48	24.87	30.38	76.56	76.10	88.70	1.34	1.43	1.91
27	P ₄ × P ₇	19.65	20.19	24.02	75.70	76.94	68.77	1.18	1.23	1.19
28	P ₄ × P ₈	17.31	17.43	26.94	101.53	103.35	77.55	1.44	1.46	1.37
29	P ₄ × P ₉	16.46	17.67	17.98	73.41	72.20	76.85	0.81	0.84	1.00
30	P ₄ × P ₁₀	22.11	23.47	20.87	67.36	65.60	73.99	0.91	0.92	1.06
31	P ₅ × P ₆	21.92	17.21	24.47	105.34	91.13	73.56	1.66	1.24	1.29
32	P ₅ × P ₇	19.53	20.35	19.85	104.25	103.44	94.28	1.39	1.41	1.27
33	P ₅ × P ₈	16.78	26.79	22.52	91.38	105.32	107.23	1.20	1.69	1.57
34	P ₅ × P ₉	25.18	20.65	22.95	101.99	68.53	63.74	2.28	0.93	0.88
35	P ₅ × P ₁₀	19.86	25.55	23.88	70.08	102.33	84.90	0.90	2.30	1.61
36	P ₆ × P ₇	30.34	27.76	26.62	115.21	116.07	96.32	2.18	2.22	1.78
37	P ₆ × P ₈	26.23	23.78	21.71	87.10	84.97	80.34	1.72	1.21	1.20
38	P ₆ × P ₉	20.88	26.42	21.86	103.43	104.47	103.00	1.84	1.91	1.68
39	P ₆ × P ₁₀	23.80	26.68	21.57	82.49	89.03	81.22	1.17	1.74	1.28
40	P ₇ × P ₈	21.88	23.31	22.71	127.14	127.81	97.58	2.28	2.34	1.62
41	P ₇ × P ₉	21.41	19.35	20.63	135.60	84.33	94.83	2.26	0.92	1.29
42	P ₇ × P ₁₀	16.63	21.80	20.67	84.50	122.87	113.56	0.89	2.28	1.84
43	P ₈ × P ₉	24.27	24.02	20.40	116.35	114.84	103.11	2.28	2.33	1.77
44	P ₈ × P ₁₀	18.35	18.56	21.87	95.95	96.57	99.28	1.35	1.38	1.83
45	P ₉ × P ₁₀	25.52	26.36	23.11	73.54	77.12	73.60	1.28	1.34	1.12
F₁ Hybrid mean		23.81	24.14	23.97	85.84	85.89	82.67	1.42	1.44	1.38
Parents										
1	P ₁	15.92	17.59	16.75	122.43	128.97	125.70	1.58	1.64	1.61
2	P ₂	29.98	32.73	31.36	49.65	43.19	46.42	0.84	0.88	0.86
3	P ₃	32.85	33.85	33.55	55.02	47.93	51.48	1.03	1.02	1.03
4	P ₄	25.69	27.51	26.60	56.19	52.33	54.26	0.80	0.83	0.81
5	P ₅	22.86	23.44	23.15	77.27	75.42	76.35	1.32	1.30	1.31
6	P ₆	18.38	20.41	19.39	107.47	97.33	102.40	1.47	1.45	1.46
7	P ₇	19.94	21.92	20.93	123.66	123.99	123.83	1.92	1.85	1.89
8	P ₈	15.82	17.58	16.70	126.68	123.86	125.27	1.53	1.53	1.53
9	P ₉	17.41	18.74	18.08	151.57	160.72	156.15	1.99	1.89	1.94
10	P ₁₀	31.05	29.60	30.33	70.23	47.45	58.84	0.97	0.94	0.95
Parental mean		22.99	24.33	23.68	94.02	90.12	92.07	1.35	1.33	1.33
Grand mean		23.66	24.17	23.92	87.33	86.66	84.38	1.40	1.42	1.37
CV		5.25	4.79	4.30	6.02	5.98	5.28	7.40	5.32	5.79
CD 5%		1.99	1.85	1.67	8.42	8.30	7.45	0.16	0.12	0.13
Range	Lowest	15.82	16.57	16.24	45.75	43.19	46.42	0.80	0.83	0.81
	Highest	36.45	36.91	33.65	151.57	160.72	156.15	2.40	2.46	2.13

Table-2. Contd...

Sr. No.	Genotypes	Unmarketable fruit yield per plant (kg)			Reducing sugar			Non reducing sugar		
		Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
1	P ₁ × P ₂	0.79	0.81	0.60	1.22	1.24	1.28	0.53	0.53	0.51
2	P ₁ × P ₃	0.43	0.50	0.55	1.14	1.15	1.15	0.45	0.40	0.38
3	P ₁ × P ₄	0.53	0.58	0.53	0.69	1.25	1.31	0.41	0.59	.55
4	P ₁ × P ₅	0.55	0.57	0.55	1.24	0.78	0.91	0.63	0.39	0.44
5	P ₁ × P ₆	0.48	0.53	0.43	0.95	0.99	0.68	0.53	0.55	0.39
6	P ₁ × P ₇	0.87	0.60	0.51	0.94	1.09	1.09	0.58	0.62	0.57
7	P ₁ × P ₈	0.38	0.79	0.69	1.17	0.95	1.10	0.55	0.58	0.55
8	P ₁ × P ₉	0.56	0.46	0.48	1.07	1.18	1.21	0.65	0.53	0.46
9	P ₁ × P ₁₀	0.47	0.57	0.62	0.97	1.01	1.10	0.60	0.60	0.55
10	P ₂ × P ₃	0.51	0.58	0.53	1.23	1.21	1.13	0.51	0.49	0.55
11	P ₂ × P ₄	0.44	0.52	0.65	1.33	1.38	1.30	0.41	0.41	0.47
12	P ₂ × P ₅	0.26	0.56	0.49	0.98	0.69	0.91	0.63	0.52	0.49
13	P ₂ × P ₆	0.36	0.48	0.51	1.30	1.30	0.99	0.52	0.48	0.45
14	P ₂ × P ₇	0.46	0.35	0.45	0.63	1.06	1.15	0.53	0.63	0.63
15	P ₂ × P ₈	0.25	0.35	0.42	1.08	1.14	1.05	0.52	0.52	0.52
16	P ₂ × P ₉	0.98	1.09	0.98	1.32	1.35	1.14	0.46	0.46	0.52
17	P ₂ × P ₁₀	0.62	0.73	0.56	1.00	1.04	1.10	0.63	0.63	0.59
18	P ₃ × P ₄	0.66	0.74	0.65	1.21	1.23	1.15	0.60	0.60	0.62
19	P ₃ × P ₅	0.28	0.48	0.47	0.88	0.90	0.93	0.41	0.41	0.50
20	P ₃ × P ₆	0.24	0.39	0.45	1.32	1.30	1.26	0.56	0.57	0.54
21	P ₃ × P ₇	0.35	0.46	0.45	0.74	0.74	1.03	0.51	0.55	0.48
22	P ₃ × P ₈	0.52	0.61	0.44	0.99	0.97	0.97	0.49	0.49	0.56
23	P ₃ × P ₉	0.61	0.72	0.54	0.73	0.80	1.05	0.67	0.67	0.60
24	P ₃ × P ₁₀	0.72	0.80	0.63	0.69	0.75	0.69	0.67	0.64	0.58
25	P ₄ × P ₅	0.51	0.65	0.45	0.54	0.62	0.85	0.49	0.49	0.51
26	P ₄ × P ₆	0.36	0.52	0.75	0.67	0.86	1.09	0.69	0.63	0.54
27	P ₄ × P ₇	0.24	0.34	0.48	0.99	1.04	1.02	0.48	0.45	0.54
28	P ₄ × P ₈	0.32	0.40	0.53	0.78	0.80	1.00	0.50	0.50	0.55
29	P ₄ × P ₉	0.34	0.42	0.35	0.97	0.98	0.93	0.58	0.58	0.49
30	P ₄ × P ₁₀	0.52	0.62	0.43	0.91	0.90	1.11	0.44	0.41	0.49
31	P ₅ × P ₆	0.74	0.36	0.36	0.60	0.82	0.78	0.59	0.30	0.41
32	P ₅ × P ₇	0.37	0.41	0.46	0.66	0.71	0.85	0.58	0.53	0.51
33	P ₅ × P ₈	0.30	0.81	0.71	0.80	0.62	0.68	0.30	0.60	0.63
34	P ₅ × P ₉	0.28	0.48	0.60	0.60	0.64	0.66	0.57	0.59	0.63
35	P ₅ × P ₁₀	0.42	0.35	0.43	0.65	0.63	0.59	0.60	0.57	0.53
36	P ₆ × P ₇	0.90	0.92	0.64	0.85	0.83	0.75	0.47	0.47	0.58
37	P ₆ × P ₈	0.60	0.77	0.50	1.13	1.01	1.00	0.56	0.51	0.49
38	P ₆ × P ₉	0.48	0.54	0.43	0.79	0.88	0.83	0.64	0.63	0.56
39	P ₆ × P ₁₀	0.71	0.66	0.50	1.04	1.10	1.04	0.54	0.54	0.56
40	P ₇ × P ₈	0.59	0.64	0.58	1.18	1.11	1.01	0.44	0.44	0.44
41	P ₇ × P ₉	0.49	0.59	0.67	0.98	0.75	0.68	0.40	0.49	0.54
42	P ₇ × P ₁₀	0.53	0.55	0.46	0.77	0.94	0.80	0.49	0.40	0.49
43	P ₈ × P ₉	0.39	0.46	0.38	1.14	1.05	0.92	0.56	0.56	0.43
44	P ₈ × P ₁₀	0.37	0.45	0.37	1.31	1.29	0.94	0.45	0.44	0.51
45	P ₉ × P ₁₀	0.60	0.61	0.51	0.87	0.92	0.78	0.53	0.52	0.56
F₁ Hybrid mean		0.50	0.57	0.52	0.96	0.98	0.97	0.53	0.52	0.52
Parents										
1	P ₁	0.39	0.54	0.47	1.33	1.37	1.35	0.50	0.50	0.50
2	P ₂	0.59	0.63	0.61	1.16	1.21	1.18	0.36	0.36	0.36
3	P ₃	0.48	0.52	0.50	1.36	1.36	1.36	0.50	0.52	0.51
4	P ₄	0.53	0.57	0.55	1.05	1.07	1.06	0.49	0.49	0.49
5	P ₅	0.34	0.43	0.38	0.37	0.36	0.37	0.24	0.23	0.24
6	P ₆	0.42	0.46	0.44	1.08	1.04	1.06	0.52	0.51	0.51
7	P ₇	0.58	0.62	0.60	1.24	1.31	1.28	0.43	0.43	0.43
8	P ₈	0.50	0.54	0.82	1.23	1.24	1.24	0.38	0.33	0.36
9	P ₉	0.68	0.67	0.68	1.20	1.27	1.24	0.50	0.49	0.49
10	P ₁₀	0.48	0.52	0.50	1.05	1.09	1.07	0.61	0.58	0.60
Parental mean		0.50	0.55	0.55	1.10	1.13	1.12	0.45	0.51	0.44
Grand mean		0.50	0.57	0.53	0.98	1.01	1.00	0.52	0.51	0.50
CV		10.48	8.53	8.11	4.76	5.75	4.39	6.26	6.51	4.45
CD 5%		0.08	0.07	0.07	0.07	0.09	0.07	0.05	0.05	0.05
Range	Lowest	0.24	0.34	0.35	0.37	0.36	0.37	2.24	0.23	0.24
	Highest	0.98	1.09	0.98	1.36	1.38	1.36	0.69	0.67	0.63

Table-2. Contd...

Sr. No.	Genotypes	Total sugar			Chlorophyll content (mg/g)			Total Phenol content		
		Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled	Y ₁	Y ₂	Pooled
1	P ₁ × P ₂	1.75	1.77	1.80	1.38	1.47	1.35	1.38	1.19	1.06
2	P ₁ × P ₃	1.59	1.55	1.53	1.09	1.19	0.70	0.94	0.82	0.97
3	P ₁ × P ₄	1.10	1.84	1.86	1.39	1.32	1.30	1.19	1.13	1.13
4	P ₁ × P ₅	1.87	1.18	1.35	1.29	1.40	1.45	0.83	1.15	1.03
5	P ₁ × P ₆	1.48	1.54	1.08	1.13	1.17	1.17	0.99	1.08	0.95
6	P ₁ × P ₇	1.52	1.71	1.65	1.40	0.98	1.14	1.12	1.44	1.14
7	P ₁ × P ₈	1.72	1.53	1.60	1.32	1.42	1.28	1.42	1.12	1.12
8	P ₁ × P ₉	1.72	1.71	1.66	0.93	1.38	1.41	0.83	0.76	0.83
9	P ₁ × P ₁₀	1.57	1.61	1.65	1.22	1.27	1.12	0.79	1.16	1.15
10	P ₂ × P ₃	1.74	1.70	1.68	1.03	1.07	1.14	1.21	1.41	1.09
11	P ₂ × P ₄	1.74	1.79	1.77	1.03	1.09	1.24	1.12	0.94	1.16
12	P ₂ × P ₅	1.60	1.21	1.40	1.42	1.34	1.21	0.86	0.99	0.96
13	P ₂ × P ₆	1.82	1.78	1.44	1.23	1.25	1.31	0.86	0.91	1.05
14	P ₂ × P ₇	1.16	1.69	1.78	1.29	1.47	1.38	1.41	0.90	0.86
15	P ₂ × P ₈	1.60	1.66	1.57	1.22	1.26	1.20	1.32	0.84	0.92
16	P ₂ × P ₉	1.78	1.80	1.66	1.31	1.39	1.39	0.88	1.21	1.16
17	P ₂ × P ₁₀	1.63	1.67	1.70	1.05	1.11	1.21	1.08	1.12	1.27
18	P ₃ × P ₄	1.80	1.83	1.77	1.11	1.16	1.04	1.09	0.86	0.84
19	P ₃ × P ₅	1.29	1.31	1.44	1.01	1.07	1.15	0.94	0.86	0.82
20	P ₃ × P ₆	1.88	1.87	1.80	1.22	1.28	1.16	1.11	0.94	1.07
21	P ₃ × P ₇	1.25	1.29	1.52	1.48	1.48	1.25	1.35	1.09	1.11
22	P ₃ × P ₈	1.48	1.46	1.53	1.34	1.35	1.39	1.44	0.88	0.87
23	P ₃ × P ₉	1.40	1.47	1.65	1.45	1.51	1.37	0.86	1.16	1.01
24	P ₃ × P ₁₀	1.36	1.38	1.27	1.17	1.21	1.25	0.76	0.88	1.15
25	P ₄ × P ₅	1.03	1.11	1.36	1.06	1.09	1.15	1.25	0.82	1.07
26	P ₄ × P ₆	1.35	1.48	1.63	1.00	1.08	1.19	1.16	1.32	1.10
27	P ₄ × P ₇	1.47	1.48	1.55	1.22	1.28	1.17	0.94	0.87	0.97
28	P ₄ × P ₈	1.28	1.29	1.55	1.31	1.29	1.20	0.87	1.35	1.22
29	P ₄ × P ₉	1.55	1.56	1.42	1.38	1.41	1.21	1.13	1.21	1.07
30	P ₄ × P ₁₀	1.35	1.31	1.60	1.33	1.31	1.27	0.98	0.94	1.02
31	P ₅ × P ₆	1.19	1.12	1.19	1.49	1.44	1.46	0.88	1.26	1.30
32	P ₅ × P ₇	1.24	1.24	1.36	1.36	1.40	1.37	1.19	1.13	1.28
33	P ₅ × P ₈	1.10	1.22	1.31	1.41	1.54	1.49	1.21	1.01	0.94
34	P ₅ × P ₉	1.17	1.23	1.29	1.03	0.95	1.06	1.32	1.19	0.98
35	P ₅ × P ₁₀	1.25	1.20	1.12	0.91	1.09	1.07	0.82	0.94	1.10
36	P ₆ × P ₇	1.32	1.30	1.33	1.43	1.44	1.22	0.94	0.86	1.01
37	P ₆ × P ₈	1.69	1.52	1.50	1.35	1.25	1.24	1.17	1.32	1.13
38	P ₆ × P ₉	1.43	1.51	1.40	1.37	1.41	1.36	1.16	1.17	1.02
39	P ₆ × P ₁₀	1.58	1.64	1.59	1.22	1.39	1.38	0.99	0.99	1.06
40	P ₇ × P ₈	1.62	1.55	1.45	1.09	1.12	1.22	0.88	1.33	1.15
41	P ₇ × P ₉	1.37	1.24	1.22	1.03	1.21	1.35	1.17	1.19	1.04
42	P ₇ × P ₁₀	1.26	1.33	1.29	1.20	1.08	1.22	1.23	0.99	1.09
43	P ₈ × P ₉	1.71	1.61	1.36	1.42	1.42	1.41	1.34	1.17	1.19
44	P ₈ × P ₁₀	1.76	1.73	1.45	1.32	1.35	1.19	1.19	1.22	1.27
45	P ₉ × P ₁₀	1.40	1.44	1.34	1.39	1.42	1.17	0.99	0.88	0.85
F₁ Hybrid mean		1.49	1.50	1.49	1.24	1.28	1.24	1.08	1.07	1.05
Parents										
1	P ₁	1.82	1.87	1.85	1.23	1.26	1.25	0.94	0.78	0.86
2	P ₂	1.51	1.57	1.54	0.96	0.97	0.97	1.12	1.12	1.12
3	P ₃	1.88	1.88	1.88	1.28	1.31	1.30	1.13	0.83	0.98
4	P ₄	1.53	1.56	1.55	1.49	1.34	1.42	0.91	1.42	1.16
5	P ₅	0.62	0.59	0.60	1.17	1.18	1.18	0.83	0.94	0.88
6	P ₆	1.60	1.55	1.57	1.30	1.32	1.31	0.84	1.12	0.98
7	P ₇	1.67	1.74	1.71	1.15	1.18	1.17	1.12	1.38	1.25
8	P ₈	1.61	1.57	1.59	1.43	1.35	1.39	0.90	0.83	0.86
9	P ₉	1.69	1.76	1.73	0.97	1.02	1.00	1.15	1.12	1.13
10	P ₁₀	1.66	1.67	1.67	1.21	1.24	1.23	0.78	0.79	0.79
Parental mean		1.56	1.57	1.56	1.22	1.21	1.22	0.97	1.03	1.00
Grand mean		1.50	1.51	1.51	1.24	1.27	1.24	1.06	1.06	1.04
CV		3.83	4.11	3.23	5.60	5.10	4.58	4.28	4.46	3.06
CD 5%		0.09	0.09	0.08	0.11	0.10	0.09	0.07	0.07	0.05
Range	Lowest	0.62	0.59	0.60	0.91	0.95	0.7	0.76	0.76	0.79
	Highest	1.88	1.88	1.88	1.49	1.54	1.49	1.44	1.44	1.30

Table-2. Contd...

Sr. No.	Genotypes	Total fruit yield per plant (kg)		
		Y ₁	Y ₂	Pooled
1	P ₁ × P ₂	1.83	1.81	1.90
2	P ₁ × P ₃	1.67	1.71	1.57
3	P ₁ × P ₄	2.06	1.91	1.71
4	P ₁ × P ₅	1.89	2.04	1.69
5	P ₁ × P ₆	1.57	1.62	1.64
6	P ₁ × P ₇	2.68	1.99	1.95
7	P ₁ × P ₈	2.30	2.60	2.55
8	P ₁ × P ₉	1.94	2.39	2.21
9	P ₁ × P ₁₀	1.60	1.76	2.21
10	P ₂ × P ₃	1.44	1.55	1.50
11	P ₂ × P ₄	1.60	1.75	1.79
12	P ₂ × P ₅	2.18	2.00	1.84
13	P ₂ × P ₆	1.57	1.70	1.88
14	P ₂ × P ₇	1.89	2.28	2.09
15	P ₂ × P ₈	1.67	1.80	1.68
16	P ₂ × P ₉	3.38	3.54	3.11
17	P ₂ × P ₁₀	1.78	1.94	2.12
18	P ₃ × P ₄	1.94	2.04	1.99
19	P ₃ × P ₅	1.43	1.66	1.63
20	P ₃ × P ₆	1.45	1.64	1.54
21	P ₃ × P ₇	1.66	1.78	1.69
22	P ₃ × P ₈	1.64	1.80	1.99
23	P ₃ × P ₉	2.06	2.21	1.89
24	P ₃ × P ₁₀	1.55	1.66	1.77
25	P ₄ × P ₅	1.42	1.58	1.62
26	P ₄ × P ₆	1.70	1.95	2.66
27	P ₄ × P ₇	1.42	1.57	1.68
28	P ₄ × P ₈	1.76	1.86	1.90
29	P ₄ × P ₉	1.15	1.26	1.35
30	P ₄ × P ₁₀	1.42	1.54	1.50
31	P ₅ × P ₆	2.40	1.60	1.63
32	P ₅ × P ₇	1.76	1.82	1.73
33	P ₅ × P ₈	1.50	2.50	2.28
34	P ₅ × P ₉	2.56	1.41	1.48
35	P ₅ × P ₁₀	1.32	2.65	2.03
36	P ₆ × P ₇	3.08	3.14	2.42
37	P ₆ × P ₈	2.32	1.97	1.70
38	P ₆ × P ₉	2.32	2.45	2.10
39	P ₆ × P ₁₀	1.88	2.41	1.78
40	P ₇ × P ₈	2.87	2.98	2.28
41	P ₇ × P ₉	2.75	1.51	1.96
42	P ₇ × P ₁₀	1.41	2.83	2.30
43	P ₈ × P ₉	2.67	2.79	2.15
44	P ₈ × P ₁₀	1.72	1.83	2.20
45	P ₉ × P ₁₀	1.87	1.94	1.63
F₁ Hybrid mean		1.91	2.02	1.91
Parents				
1	P ₁	1.99	2.10	2.05
2	P ₂	1.43	1.51	1.47
3	P ₃	1.51	1.55	1.53
4	P ₄	1.33	1.40	1.37
5	P ₅	1.65	1.73	1.69
6	P ₆	1.90	1.91	1.91
7	P ₇	2.50	2.47	2.49
8	P ₈	2.03	2.07	2.05
9	P ₉	2.67	2.56	2.61
10	P ₁₀	1.45	1.46	1.46
Parental mean		1.84	1.87	1.86
Grand mean		1.90	1.99	1.90
CV		6.24	4.71	4.93
CD 5%		0.19	0.15	0.16
Range	Lowest	1.15	1.26	1.35
	Highest	3.38	3.54	3.11