

Original Research Article

Effect of STCR-IPNS Based Nutrient application on Soil Health, Yield, Nutrient Content and Uptake of Mustard (*Brassica juncea* L.) in eastern plain zone of Uttar Pradesh, India

Abstract

The improvement in grain yield characters was the manifestation of improved growth characters as a result of higher uptake of nutrients caused by balanced supply of nutrients in this regard soil test based nutrient management approaches aims provide a scientific basis for balanced fertilization to obtain more yield per unit of fertilizer investment. An experiment was conducted during kharif season 2017-18 in the Soil Science and Agricultural Chemistry Research Farm, SHUATS, Prayagraj. The cursory glance of data revealed that the bulk density and particle density of soil was found to be non-significant in different levels of fertilizer recommendation based on soil test values. The maximum soil pore space (60.37%) and water retaining capacity (81.25 %) was recorded in treatment T₄ [STCR + 5 t FYM]. The maximum available Nitrogen (305.82 kg ha⁻¹), available Phosphorus (26.90 kg ha⁻¹), available Potassium (205.07 kg ha⁻¹) and available Sulphur (14.23 ppm ha⁻¹) in soil was recorded in treatment T₈ [STCR + @ 50 % FYM + @ 50 % S]. The maximum seed yield of mustard (11.53 q ha⁻¹) and stover yield (16.03 q ha⁻¹) was associated with the treatment T₈ [STCR + @ 50 % FYM + @ 50 % S]. Result showed that that application of T₈ [STCR + @ 50 % FYM + @ 50 % S] significantly recorded maximum nutrient content viz. N (2.19%), P (0.23%), K (1.68%) and S (4.8%) content in grain N (1.73%), P (0.21%), K (1.47%) and S (3.9%) content in stover and maximum nutrient uptake viz. N (25.25 kg ha⁻¹), P (2.65 kg ha⁻¹), K (19.37 kg ha⁻¹) and S (55.34 kg ha⁻¹) uptake in grain is and N (27.73 kg ha⁻¹), P (3.36 kg ha⁻¹), K (23.56 kg ha⁻¹) and S (62.51 kg ha⁻¹) uptake in stover.

Key Words: Mustard, Nutrient Content, Nutrient Uptake, STCR and Yield

Introduction

Rapeseed-mustard (*Brassica campestris*) is a major oilseed crop contributing important share in oilseed production in the country. Production of rapeseed and mustard declined from 8.03 MT in 2012-13 to 6.82 MT in 2015-16 (**Anonymous 2017**).

India is amongst the largest vegetable oil economic in the world. The present average per capita consumption of oils and fats has not been more than 11g day⁻¹ as against the nutritional

standard of 30g/day for a balanced diet. Mustard is rich in minerals like calcium, manganese, copper, iron, selenium, zinc, vitamin A, B, C and proteins. 100g mustard seed contains 508 kcal energy, 28.09 Mg carbohydrates, 26.08g proteins, 36.24g total fat and 12.2g dietary fiber. (Upadhyay *et al.*, 2016)

The nutrient elements of major significance for yield and quality of yellow mustard are nitrogen, phosphorus sulphur and Zinc. Nitrogen is the most important which determines the growth of yellow mustard that increases the amount of protein, methionine, dry matter and yield. Phosphorus and potash are known to be efficiently utilized in the presence of nitrogen to promote flowering, setting of siliqua and increases the size of siliqua and yield. (Singh *et al.*, 2012)

Sulphur is considered to occupy fourth place among major plant nutrient after nitrogen, phosphorus and potassium. It increases phosphorus uptake by plant and nitrogen in protein synthesis and is indispensable for the synthesis of essential amino acid like cysteine and methionine. Besides, sulphur is also involved in various metabolic processes of plants. It is a constituent of glutathione, a compound supposed to be associated with the plant respiration and the synthesis of essential oils. Sulphur also plays a vital role in chlorophyll formation. (Yadav *et al.*, 2017)

There are several reasons behind such yield reduction including poor soil nutrient status. Soil fertility declination under continuous cropping has been witnessed which need to be restored for sustaining and increasing crop yield. Soil fertility restoration can effectively be achieved by integrated management of nutrient sources (Joshi *et al.*, 2017), but due to lack of proper knowledge of method and time of manuring and fertilizer application; the cost of cultivation increased. Soil fertility maintenance required adequate knowledge of soil nutrient status, fertilizer efficiency soil efficiency, time and methods of fertilizer application. Adoption of soil test crop response (STCR) suggested by Ramamoorthy *et al.* (1967) is efficient approach concerning all aspects of nutrient management. Supplying of plant nutrients based on STCR approach significantly improved crop yield as well as soil health (Rajput *et al.* 2016) and is very important for yield sustainability and reducing fertilizer cost (Saxena *et al.*, 2008). Implementation of inductive approach of STCR in Chhattisgarh may reduce cost of cultivation and may also encourage smart and strategic nutrient management practices.

In the targeted yield approach, it is assumed that there is linear relationship between grain yield (economic produce) and nutrient uptake by the crop. Targeted yield concept, thus strikes a balance between “Fertilizing the crop” and “Fertilizing the soil”. This approach can be used not only for individual field situations but also as a better approximation for planning

the requirement of fertilizers on area basis for a given level of crop production. Fertilizer application and the yield targets chosen can be so manipulated that both high profits from fertilizer investment and maintenance of soil fertility can be achieved (Velayutham 1979). The targeted yield approach has been used to formulate fertilizer recommendations across the country (Puri *et al.* 1993, 1994; Verma and Bhagat 1995; Verma *et al.* 2017; Singh *et al.* 2017).

Resources and Methods

Soil of Experimental Field

The soil of experimental field is sandy loam in texture, good aeration (47.53 % porosity), alkaline in reaction (pH 7.58), low in organic carbon (0.45%), low in available N (238.21 kg ha⁻¹), medium in available P (20.73 kg ha⁻¹), high in available K (127.65 kg ha⁻¹) and low in available sulphur (9.82 ppm ha⁻¹).

Experimental site

The experiment was conducted at research Farm of Soil Science and Agricultural Chemistry at Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj. The area is situated on the south of Prayagraj on the right side of the river Yamuna on the South of Rewa Road at a distance of about 6 Km from Prayagraj city. It is situated at 25°24'23" N latitude, 81°50'38" E longitude and at the altitude of 98 meter above the sea level.

Layout and Design of the Experiment

The experiment was laid out in randomized block design with three replications. The total numbers of unit plots were 27. The size of a unit plot was 2.0 m X 2.0 m. The width of the main irrigation channel is 1.0 m and the width of the sub-irrigation channel is 0.5 m.

Treatments of the Investigation:

Table 1: Treatment Combination

S. No.	Symbol	Description
1.	T ₀	[Control]
2.	T ₁	[RDF + 5 t FYM]
3.	T ₂	[STL + 5 t FYM]
4.	T ₃	[FP + 5 t FYM]
5.	T ₄	[STCR + 5 t FYM]
6.	T ₅	[RDF + 5 t FYM + 50 % S]
7.	T ₆	[STL + 5 t FYM + 50 % S]

8.	T ₇	[FP + 5 t FYM + 50 % S]
9.	T ₈	[STCR + 5 t FYM + 50 % S]

RDF- Recommended dose of fertilizers (80:40:40 kg ha⁻¹)

STL- Soil Test Levels (80:28:28 kg ha⁻¹)

FP- Farmer's Practice (50:30:30 kg ha⁻¹)

STCR- Soil Test Crop Response (40:15:15 kg ha⁻¹)

STCR approach

The following STCR equation developed for Mustard was used for achieving 25 q ha⁻¹ yield target.

With FYM

1. Nitrogen dose (kg ha⁻¹) = 12.27T - 0.56SN - 0.09FYM - N

2. Phosphorus dose (kg ha⁻¹) = 4.60T - 3.29SP - 0.06FYM - P

3. Potassium dose (kg ha⁻¹) = 4.69T - 0.24SK - 0.05FYM - K

Where, T = Yield target (q ha⁻¹), SN = Alkaline KMnO₄-N, SP = Olsen's P (kg ha⁻¹) and SK = Ammonium Acetate - K (kg ha⁻¹).

The fertilizer adjustment equations were ready for determining requirement of fertilizer. Say for 25 q ha⁻¹ the yield target of mustard with varying soil test values in table 3.5. These results were shows that the fertilizer requirement varies with the soil test values for a particular target yield. Similar result was also reported by **Mishra *et al.* (2010) and Singh *et al.* (2014).**

1. Nutrient Requirement

- | | |
|--|---|
| a. Kg N required per quintal of seed = | $\frac{\text{Total uptake of N (kg ha}^{-1}\text{)}}{\text{Seed yield (q ha}^{-1}\text{)}}$ |
| Production | |
| b. Kg P required per quintal of seed = | $\frac{\text{Total uptake of P (kg ha}^{-1}\text{)}}{\text{Seed yield (q ha}^{-1}\text{)}}$ |
| Production | |
| c. Kg K required per quintal of seed = | $\frac{\text{Total uptake of K (kg ha}^{-1}\text{)}}{\text{Seed yield (q ha}^{-1}\text{)}}$ |
| Production | |

Preparation and analysis of soil samples

Soil samples from each plot at 0-15 cm depth were collected at different stages were air-dried, grind and passed through 2 mm sieve and finally stored in polythene bags for analysis of different physico-chemical parameters and changes in available N, P, K and % Organic carbon content. The soil sample was analyzed for Bulk density, particle density, % pore space (**Muthuaval *et.al.*,1992**), soil texture (**Bouyoucos Hydrometer Method, 1952**), pH

(Jackson, 1973), Available N (Subbaih and Asija, 1956), P (Olsen *et al.*, 1954), K (Toth and Price, 1949) and S (Bardsley and Lancaster, 1960).

Plant Analysis for Content and Uptake of Nutrient

The chemical analysis of plants for the nutrient content was done when grain and straw samples were collected from each treatment at harvest to analyse nitrogen, phosphorous, potassium concentration (%) and sulphur concentration (ppm) and their uptake (kg ha⁻¹). The plant material was oven dried (70 ± 5°C for 72 hours) and ground separately and then subjected to analysis. Plant analysis for the determination of nutrient content in grain and stover were done with the standard procedures viz., nitrogen concentration in plant (both grain and stover) was determined by micro-kjeldahl's method, phosphorus by vanado-molybdo phosphoric acid yellow colour method, potassium by flame photometer and sulphur by Turbidometric Method (Jackson, 1973). The uptake of nitrogen, phosphorus, potassium and sulphur were calculate by the following formulas:

$$\text{Nutrient uptake (N, P, K kg ha}^{-1}\text{)} = \frac{\text{Nutrient content in grain and straw (\%)} \times \text{Seed and Stover Yield (kg ha}^{-1}\text{)}}{100}$$

Nutrient response ratio (kg yield kg nutrient⁻¹)

It was calculated by using following equation (Indian Society of Agronomy, New Delhi).

$$\text{NRR} = \frac{\text{Yield (kg)}}{\text{Amount of nutrient applied (kg)}}$$

Result and Discussion

1. Physical Properties of Soil

It is obvious from the data given in table-2 and depicted in fig.1 clearly shows that response Bulk density and Particle density of soil was found to be non-significant in different levels of fertilizer recommendation based on soil test values. The maximum Bulk density (1.09 Mgm⁻³) and Particle density (2.73 Mgm⁻³) of soil was recorded in treatment T₂ [STL + 5 t FYM] and minimum Bulk density (1.03 Mgm⁻³) and Particle density (2.24 Mgm⁻³) of soil was

recorded in treatment [STCR + 5 t FYM + 50 % S]. Similar results were also reported by Nagar *et al.*, (2015) and Sahu *et al.*, (2015)

The response of soil pore space and water retaining capacity (WRC) was found to be significant in different levels of fertilizer recommendation based on soil test values. The maximum soil pore space (60.37 %) and maximum WRC (81.25%) was recorded in treatment T₄ [STCR + 5 t FYM] and minimum soil pore space (52.63 %) and minimum WRC (60.00%) was recorded in treatment T₇ [FP + 5 t FYM + 50 % S]. The results of the present investigation are also in agreement with the findings of Ahmadi and David (2016) and Alam *et al.*, (2014).

Table 2: Effect of different levels of fertilizer recommendation based on soil test values on physical properties of soil after crop harvest.

Treatments	BD (Mg m ⁻³)	PD (Mg m ⁻³)	% PS	WRC (%)
T ₀	1.04	2.35	53.62	73.33
T ₁	1.07	2.45	55.45	75.00
T ₂	1.09	2.73	60.04	61.11
T ₃	1.09	2.52	56.33	69.23
T ₄	1.04	2.62	60.37	81.25
T ₅	1.05	2.36	54.58	68.75
T ₆	1.05	2.45	56.16	62.50
T ₇	1.05	2.35	52.63	60.00
T ₈	1.03	2.24	53.51	75.00
F-test	NS	NS	S	S
S. Em_±	0.025	0.171	0.169	1.38
C.D. (P= 0.05)	0.054	0.362	0.359	2.94




Fig. 1: Effect of different levels of fertilizer recommendation based on soil test values on physical properties of soil after crop harvest.

2. Chemical Properties of Soil

An appraisal of the data given in table 3 and depicted in fig. 2 clearly shows that available N, P, K and S in soil increased significantly with the increase in different levels of fertilizer recommendation based on soil test values. The maximum available N ($305.82 \text{ Kg ha}^{-1}$), available P (26.90 Kg ha^{-1}), available K ($205.07 \text{ Kg ha}^{-1}$) and available S ($14.23 \text{ ppm ha}^{-1}$) in soil was recorded in treatment T_8 [STCR + 5 t FYM + 50 % S] and the minimum available N ($289.13 \text{ Kg ha}^{-1}$), available P (19.10 Kg ha^{-1}), available K ($183.97 \text{ Kg ha}^{-1}$) and available S ($10.43 \text{ ppm ha}^{-1}$) in soil was recorded in treatment T_0 [control]. The consequences of the current investigation are additionally in concurrence with the investigation of **Upadhyay et al., (2015)**, **Rajput et al., (2016)** and **P. Dey (2016)**

Table 3: Effect of different levels of fertilizer recommendation based on soil test values on Chemical Properties of soil after crop harvest.

Treatments	N (kg ha^{-1})	P (kg ha^{-1})	K (kg ha^{-1})	S (ppm ha^{-1})
T_0	289.13	19.10	183.97	10.43
T_1	299.46	23.95	196.63	11.40
T_2	302.51	25.03	202.50	13.40
T_3	297.46	22.90	192.43	10.73
T_4	300.52	24.11	197.07	12.30
T_5	303.86	25.75	203.47	13.47
T_6	298.83	23.93	193.07	10.97
T_7	301.16	24.70	198.20	12.87
T_8	305.82	26.90	205.07	14.23

F-test	S	S	S	S
S. Em _±	0.980	0.311	0.430	0.060
C.D. (P= 0.05)	2.078	0.661	0.912	0.127




Fig. 2: Effect of different levels of fertilizer recommendation based on soil test values on Chemical properties of soil after crop harvest.

3. Seed and Stover Yield (q ha⁻¹)

It is visualized from the data given in table-4 and depicted in fig. 3 that Seed Yield and of Mustard was found to be increased significantly but stover yield of mustard was found to be non-significant with the increase in different levels of fertilizer recommendation based on soil test values. The maximum Seed Yield (11.53 q ha⁻¹) and maximum Stover Yield (16.03 q ha⁻¹) was recorded as in T₈ [STCR + 5 t FYM + 50 % S] and the minimum Seed Yield (9.33 q ha⁻¹) and Stover yield (13.77 ha⁻¹) was recorded as in T₀ [control]. Comparative findings were detailed by Kumar *et al.*, (2016) and Pal and Pathak (2016).

Table 4: Effect of different levels of fertilizer recommendation based on soil test values on Seed Yield (q ha⁻¹)

Treatment	Seed Yield (q ha ⁻¹)	Stover Yield (q ha ⁻¹)
T ₀	9.33	13.77
T ₁	9.53	15.13
T ₂	9.83	15.33
T ₃	9.47	14.90
T ₄	11.13	15.53

T₅	9.67	15.20
T₆	9.50	15.10
T₇	9.70	15.23
T₈	11.53	16.03
F-test	S	NS
S. Em₊	0.222	0.395
C.D. (P= 0.05)	0.472	0.836




Fig. 3: Effect of different levels of fertilizer recommendation based on soil test values on yield of mustard.

4. Plant Nutrient Concentration

A critical perusal of the data given in table-5 and depicted in fig. 4 clearly shows the nutrient concentration (N, P, K and S) in Stover and grain increased significantly with the increase in different levels of fertilizer recommendation based on soil test values. The maximum N (1.73% in Stover and 2.19 % in grain), P (0.21% in Stover and 0.23 % in grain), K (1.47% in Stover and 1.68 % in grain) and S (3.9 % in Stover and 4.8 % in grain) in treatment T₈ [STCR + 5 t FYM + 50 % S] which was significantly higher than any other treatment combination and the minimum N (1.05% in Stover and 1.18 % in grain), P (0.12% in Stover and 0.14 % in grain), K (1.08% in Stover and 1.19 % in grain) and S (2.8 % in Stover and 4.0 % in grain) was recorded in treatment T₀ [control]. The results of the present investigation are also in agreement with the findings of **Bharose *et al.*, (2011)** and **Chaurasia *et al.*, (2009)**

Table 5: Effect of different levels of fertilizer recommendation based on soil test values on Nutrient Concentration in Stover and Grain.

Treatments	N (%)		P (%)		K (%)		S (%)	
	Stover	Grain	Stover	Grain	Stover	Grain	Stover	Grain
T ₀	1.05	1.18	0.12	0.14	1.08	1.19	2.8	4.0
T ₁	1.17	1.44	0.14	0.16	1.11	1.21	3.1	4.2
T ₂	1.40	1.59	0.15	0.18	1.15	1.30	3.0	4.1
T ₃	1.14	1.70	0.14	0.17	1.17	1.27	3.4	4.3
T ₄	1.61	1.99	0.17	0.21	1.39	1.57	3.7	4.6
T ₅	1.50	1.82	0.16	0.18	1.32	1.37	3.5	4.4
T ₆	1.53	1.77	0.15	0.15	1.21	1.44	3.2	4.3
T ₇	1.57	1.82	0.16	0.19	1.26	1.51	3.6	4.6
T ₈	1.73	2.19	0.21	0.23	1.47	1.68	3.9	4.8
F-test	S	S	S	S	S	S	S	S
S. Em _±	0.08	0.07	0.01	0.01	0.06	0.07	0.14	0.15
C.D. (P= 0.05)	0.25	0.21	0.03	0.03	0.19	0.20	3.15	3.19




Fig. 4: Effect of different levels of fertilizer recommendation based on soil test values on nutrient content in grain and stover.

5. Nutrient uptake by Grain and Stover of Mustard

At a glance over the data given in the table-6 and depicted in fig. 5 clearly shows the nutrient uptake (N, P, K and S) in Stover and grain increased significantly with the increase in different levels of fertilizer recommendation based on soil test values. The maximum uptake of N (27.73 kg ha^{-1} in Stover and 25.25 kg ha^{-1} in grain), P (3.36 kg ha^{-1}

¹ in Stover and 2.65 kg ha⁻¹ in grain), K (23.56 kg ha⁻¹ in Stover and 19.37 kg ha⁻¹ in grain) and S (62.51 kg ha⁻¹ in Stover and 55.34 kg ha⁻¹ in grain) in treatment T₈ [STCR + 5 t FYM + 50 % S] which was significantly higher than any other treatment combination and the minimum uptake N (14.45 kg ha⁻¹ in Stover and 11.00 kg ha⁻¹ in grain), P (1.65 kg ha⁻¹ in Stover and 1.30 kg ha⁻¹ in grain), K (14.87 kg ha⁻¹ in Stover and 11.10 kg ha⁻¹ in grain) and S (38.55 kg ha⁻¹ in Stover and 37.32 kg ha⁻¹ in grain) was recorded in treatment T₀ [control]. The consequences of the current investigation are additionally in concurrence with the investigation of **Raghvendra *et al.*, (2017) and Dhruw *et al.*, (2018)**

Table 6: Effect of different levels of fertilizer recommendation based on soil test values on Nutrient Uptake in Stover and Grain.

Treatments	N (kg ha ⁻¹)		P (kg ha ⁻¹)		K (kg ha ⁻¹)		S (kg ha ⁻¹)	
	Stover	Grain	Stover	Grain	Stover	Grain	Stover	Grain
T₀	14.45	11.00	1.65	1.30	14.87	11.10	38.55	37.32
T₁	17.70	13.72	2.11	1.52	16.79	11.53	46.90	40.02
T₂	21.46	15.62	2.29	1.76	17.62	12.77	45.99	40.30
T₃	16.98	16.09	2.08	1.60	17.43	12.02	50.66	40.72
T₄	25.00	22.14	2.64	2.33	21.58	17.47	57.46	51.19
T₅	22.80	17.55	2.43	1.74	20.06	13.24	53.20	42.54
T₆	23.10	16.81	2.26	1.42	18.27	13.68	48.32	40.85
T₇	23.91	17.65	2.43	1.84	19.18	14.64	54.82	44.62
T₈	27.73	25.25	3.36	2.65	23.56	19.37	62.51	55.34
F-test	S	S	S	S	S	S	S	S
S. Em_±	0.05	0.10	0.04	0.07	0.36	0.29	1.01	0.17
C.D. (P= 0.05)	0.16	0.32	0.14	0.23	1.09	0.89	3.06	0.54




Fig. 5: Effect of different levels of fertilizer recommendation based on soil test values on nutrient uptake by grain and stover.

6. Nutrient Response Ratio (kg kg⁻¹)

It is apparent from the data given in table-7 and depicted in fig. 6 clearly shows that the nutrient response ratio (kg kg⁻¹) was found to be increased significantly with the increase in different levels of fertilizer recommendation based on soil test values. The maximum total nutrient response ratio was recorded as 436.36 kg kg⁻¹ in T₈ [STCR + 5 t FYM + @ 50 % S] followed by T₄ [STCR + 5 t FYM] with the total NRR value 244.37 kg ka⁻¹ and the minimum total nutrient response ratio was recorded as 146.22 kg kg⁻¹ in T₃ [FP + 5 t FYM].

Table 7: Effect of different levels of fertilizer recommendation based on soil test values on Nutrient Response Ratio (kg kg⁻¹)

Treatment	Nitrogen Response Ratio (kg kg ⁻¹)	Phosphorus Response Ratio (kg kg ⁻¹)	Potassium Response Ratio (kg kg ⁻¹)	Total Nutrient Response Ratio (kg kg ⁻¹)
T ₀	-	-	-	-
T ₁	30.83	61.65	61.65	154.13
T ₂	31.45	89.85	89.85	211.15
T ₃	48.74	48.74	48.74	146.22
T ₄	66.65	88.86	88.86	244.37
T ₅	31.08	62.17	62.17	155.42
T ₆	30.82	88.07	88.07	206.96
T ₇	49.86	83.10	83.10	216.06
T ₈	68.90	183.73	183.73	436.36
F-test	S			
S. Em±	16.62			




Fig. 6: Effect of different levels of fertilizer recommendation based on soil test values on Nutrient Response Ratio (kg kg⁻¹)

Conclusion

On the basis of results emanated from present investigation, it could be concluded that STCR based integrated nutrient management not only gave higher crop yield but also provide highest nutrient content and uptake in grain and stover of mustard which is subjected to nutrient enrichment in mustard seed. Our results also highlight that STCR-IPNM based nutrient application is effective tool of sustaining soil health. Therefore, STCR-IPNM based nutrient management can be recommended as an effective tool for balanced fertilization.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

References

Anonymous, (2017). Agricultural Statistics at a glance. Directorate of Economics & Statistics, Ministry of Agriculture, Govt. of India.

Ahmadi, S. A. and David, A. A. (2016), Effect of nitrogen and zinc on Physico-chemical properties of soil of Allahabad Uttar Pradesh India. *International Journal of Multidisciplinary Research and Development*. Volume (3); Issue 5; Page No. 288-290 ISSN: 2349-4182.

Alam, Md. S., Mishra, A. K., Singh, K., Singh, S. K. and David, A. A. (2014). Response of sulphur and FYM on soil physico-Chemical Properties and Growth, Yield and Quality of

Mustard (*Brassica Nigra L.*) *Journal of Agricultural Physics*, Vol. (14), No. 2, ISSN 0973-032X, pp. 156-160.

Bardsley, C.E. and Lancaster, J. D. (1960). Determination of reserve sulphur and soluble sulphates in soil. *Soil Sci. Soc. Amer. Proc.* 24 265- 268.

Bharose, R., Chandra, S., Thomas, T. and Dhan, D. (2011). Effect of different levels of phosphorus and sulphur on yield and availability of NPK protein and oil content in Toria (*Brassica sp.*) VAR. P.T.–303. *ARPN Journal of Agricultural and Biological Science*. Vol (6) No 2.

Bouyoucos, G.J. (1927). The hydrometer as a new method for the mechanical analysis of soil. *Soil Sci.* 23343-353.

Chaurasia, A., Singh, S. B., & Namdeo, K. N. (2009). Integrated nutrient management in relation to nutrient contents and uptake of Ethiopian mustard (*Brassica carinata*). *Research on Crops*, 10(2), 246-249.

Dey, P. (2016). Soil health management. *Soil Health*, 79.

Dhruw, T. K., Sharma, G. K., & Joshi, S. K. (2018). Effect of soil test based nutrient management on crop yield, nutrient requirement and relationship between nutrient uptake and yield of mustard (*Brassica campestris*) in Alfisol. *IJCS*, 6(6), 2022-2024.

Jackson, M. L. (1973). Soil Chemical Analysis, *Prentice Hall of India Private Limited, New Delhi*.

Joshi, S. K., Nag, G. P., Singh, D. P., Sahu, Y. K., & Kumawat, N. (2017). Long-term effect of nutrient management on active organic pools: A review. *International Journal of Chemical Studies*, 5(4), 576-579.

Kumar, V., Nath, P., Kumar, R., Kumar, V., Verma, J. K. and Naresh, R. K. (2016). Interactive effect of sulphur and nitrogen on growth, yield and quality of Indian Mustard (*Brassica juncea L.*). *I.J.S.N.*, Vol.7 (1): 57-61 ISSN 2229 – 6441.

Muthuval P., Udaysoorian, C., Natesan, R., and Ramaswami, P. P. (1992). Introduction to Soil analysis, Tamil Nadu Agricultural University, Coimbatore-641002.

Nagar, K. C., Meena, H. and Swaroop, N. (2015). Effect of different levels of inorganic fertilizer and sulphur on physico-chemical properties of soil, and yield of mustard (*Brassica juncea L.*). *An Asian Journal of Soil Science*. Vol. (10), ISSN–0976–7231.

Olsen, S.R., Cole, C.V., Watnahe, F.S. and Dean, L. A. (1954). Estimation of available phosphorous in soil by extraction with sodium bicarbonate U.S. Dept. Agr. *Cric.* 939.

Pal, R. L. and Pathak, J. (2016). Effect of integrated nutrient management on yield and economics of mustard. *Indian Journal of Science and Nature*. Vol. (7) ISSN 2229 – 6441.

- Puri, G. and Jaipurkar, S.A. (1993)** Predication and optimization of soil test based fertilizer recommendation for safflower production in Vertisols of central India. In Third International Safflower Conference, held on June 14-18 at Institute of Botany at Beijing, China, pp. 45-51.
- Puri, G. and Jaipurkar, S.A. (1994)** Evaluating the effects of soil fertility variables on the yield of mustard (*Brassica compestris*) on swell shrink soil (Vertisol) of central India accepted for presentation. In 15th International Congress of Soil Science, pp. 45-49.
- Raghavendra, Rao K.N., Wani S.P., Ravi M.V., Veeresh H., Channabasavanna A.S. And Swamy M. (2017).** Effect of soil test based nutrient management approaches on grain yield and nutrient uptake of Dry DSR-mustard cropping system. *Agriculture Update* Volume 12 | TECHSEAR-5 | 2017 | 1286-1290
- Rajput, P. S., Srivastava, S., Sharma, B. L., Sachidanand, B., Dey, P., Aher, S. B., & Yashona, D. S. (2016).** Effect of soil-test-based long-term fertilization on soil health and performance of rice crop in Vertisols of central India. *International Journal of Agriculture, Environment and Biotechnology*, 9(5), 801-806.
- Ramamoorthy, B., Narasimhan, R. L., & Dinesh, R. S. (1967).** Fertilizer recommendations based on fertilizer application for specific yield of Sonara-64. *Indian Farming*, 17(443), 51.
- Sahu, Y. K, David, A. A, Upadhyay, Y, Dhruw, S. S. & Sidar, R. S (2015).** Influence of organic manure and various level of NPK on soil physico-chemical properties of Mustard (*Brassica juncea* L.) cv. Euro Shakti. *International Journal of Agricultural Science and Research (IJASR)* vol (5), ISSN (P): 2250-0057.
- Saxena, A. K., Singh, S., Srivastava, A., & Gautam, P. (2008).** Yield target approach under integrated nutrient management for assessing fertilizer requirements of onion in Mollisols of Uttarakhand. *Indian Journal of Horticulture*, 65(3), 302-306.
- Singh, J., Singh, N.S.H. and Bhadauria, H.S. (2012)** Nitrogen and sulphur requirement of mustard under different crop sequences. *Ann. Pl. Soil Res.* 14(2): 113-115
- Singh, M., & Dixit, S. P. (2017).** Yield, profitability and nutrient uptake of wheat under soil test crop response based fertilizer application with different levels of lime in an acid Alfisol. *Journal of Pharmacognosy and Phytochemistry*, 6(6), 1985-1988.
- Singh, M. and Kumar, M. (2014).** Effect of nitrogen and sulphur levels on seed yield and some other characters in mustard [*Brassica juncea* (L.) Czern and Coss]. *International Journal of Agricultural Sciences*. Volume (10), 449-452.
- Subbiah, B. V. and Asija, C. L. (1956).** A rapid procedure for the estimation of available nitrogen in soil, *Current Sci.* 25. 259-260.

Toth, S. J. and Prince, A. L. (1949). Estimation of cation exchange capacity and exchangeable Ca K and Na Content of Soil by Flame photometer technique. *Soil Sci.* 67 439-445.

Upadhyay, Y., Swaroop, N., Sahu, Y. K., Dhruw, S. S. and Verma, P. D. (2016). Interaction effects of different doses of sulphur and zinc with NPK on physico-chemical properties of soil in yellow mustard (*Brassica campestris* L.) Cv. Krishna Super Goldi. *International Journal of Agricultural Science and Research (IJASR)*, vol. (6), ISSN (P): 2250-0057.

Velayutham, M. (1979) Fertilizer recommendation based on targeted yield concept problem and prospects. *Fertiliser News* 24, 12-17.

Verma, T.S. and Bhagat, R.H. (1995) Nitrogen use efficiency as affected by time variant nitrogen application to wetland rice. *Oryza* 32, 276-279

Verma, M., Singh, Y. V., Dey, P., & Babu, A. (2017). Soil test based fertilizer recommendation for mustard (*Brassica juncea* L.) in Eastern Plain Zone of Uttar Pradesh, India. *International Journal of Current Microbiology and Applied Science*, 6(2), 155-161.

Yadav, R., Singh, P. K., Singh, R. K., Tiwari, P. and Singh, S. N. (2017). Impact of Sulphur Nutrition on Promising Mustard Cultivars in Eastern Uttar Pradesh. *Int. J. Pure App. Biosci.* (5) ISSN: 2320 – 7051 p: 389-394.