

# PERFORMANCE OF GLUCONATE AND LACTATE BASED FORMULATIONS ON PLANT GROWTH AND YIELD ATTRIBUTES IN MAIZE (*Zea mays* L.)

**Aim/Objectives:** To check the efficacy of gluconate and lactate based formulations on plant growth and yield characteristics of maize

**Place and Duration of Study:** School of Agricultural Sciences, Malla Reddy University, Hyderabad, spring 2021

**Methodology:** The experiment was laid-out in Randomized Block Design (RBD), with Maize hybrid (DHM-117) spacing adopted (60 x 20 cm) with 4 replications. The treatments viz; T<sub>1</sub>: Control (No fertilizer), T<sub>2</sub>:100 % RDF, T<sub>3</sub>: 75 % RDF + 25 % Sahasra Organic Formulations (SOF), T<sub>4</sub>:50 % RDF + 50 % SOF, T<sub>5</sub>: 25 % RDF + 75 % SOF, T<sub>6</sub>: 100 % SOF. The RDF (N:72-80; P:24; K:20 kg/acre).

**Results:** The yield attributes were recorded intermittently and maximum grain yield (7,649.8 kg ha<sup>-1</sup>) and Stover yield (8,859.87 kg ha<sup>-1</sup>) were recorded with T<sub>2</sub> (100% RDF), followed by T<sub>3</sub> (75 % RDF + 25 % Sahasra Organic Formulations). Similarly gross returns (₹ 123606.5 ha<sup>-1</sup>), net returns (₹ 87116.49 ha<sup>-1</sup>) and B: C (3.4) ratio was also recorded highest in T<sub>2</sub> followed by T<sub>3</sub> (75 % RDF + 25 % SOF), while the lowest were recorded with T<sub>1</sub> (control).

**Conclusion:** Adoption of gluconate and lactate based formulations enhanced soil fertility, soil organic carbon and yield in Maize crop. The study concluded that the integrated approach *i.e.*, T<sub>3</sub> (75 % RDF + 25 % SOF) has realized the highest net returns (₹ 82466.8 ha<sup>-1</sup>) and B: C ratio (3.4) in comparison to the 100 % RDF treatment thus reducing the cost of fertilizers and benefiting the net returns to the farmers.

---

**Keywords:** Gluconate, Lactate, Innovative Technologies, Soil Fertility, Net Returns.

## 1 Introduction

Maize is grown almost all over the world in different agro-climatic conditions and different seasons. It is the third most important food crop next to rice and wheat. It is also known as the miracle crop or queen of cereals due to its high productivity potential among the cereal crops of the Graminaceae family. In India, maize is not only grown for food and fodder but

also several industrial uses and acquired a dominant role in the farming sector and macro-economy of the Asian region (Mauria *et al.*, 1998). Furthermore, one of the most critical elements influencing maize crop development and yield productivity is fertilizer management (Ghosh *et al.*, 2020). The current NPK fertilizer consumption ratio is 10: 2.9 : 1 against the optimal ratio of 4: 2: 1 (Walia and Kler, 2010). Maize area and production have steadily increased in India during the past two decades and it is cultivated in an area of 9.2 M ha with an average production of 28.7 Mt and productivity of 3115 kg ha<sup>-1</sup>. This non-judicious nutrient application by farmers causes multi-nutrient deficiencies. In this context, the smart use of integrated nutrient management and use of organic formulations is one of the greatest options for ensuring long-term crop productivity, while maintaining soil fertility in maize and other cereal-based cropping systems. This ultimately improves crop yield (Garima and Pant, 2018). Cultivation of maize in the spring season is a common practice in Peninsular India (Telangana, Andhra Pradesh, Karnataka and Tamil Nadu) as well as in North Eastern plains, where the winter remains frost free and the temperature does not fall below 12°C (Nirupama Singh, 2012).

### **1.1 Gluconic acid based formulations:**

Among various organic acids, gluconic acid seems to be the major mechanism of phosphate solubilization by gram negative bacteria (Goldstein *et al.*, 1993). Gluconic acid is produced by the oxidative metabolism of glucose dehydrogenase enzyme through microbial fermentation with different types of *Aspergillus niger* strains (Kim *et al.*, 1997).

### **1.2 Lactic acid based formulations:**

Different strains of *Lactobacillus* are used for Lactic acid production. Lactic acid bacteria have also been used for the treatment of animal manures, farm yard manure and sewage for odor abatement and as an inoculant to accelerate the composting of organic wastes (Okada, 1998).

### **1.3 Amino acids based formulations:**

The chelating ability of amino acids has been used in fertilizers for agriculture to facilitate the delivery of minerals to plants to correct mineral deficiencies, such as iron chlorosis and other nutrient deficiencies. These fertilizers are also used to prevent deficiencies from occurring and improve the overall health of the plants. Amino acids can be extracted from marine algae like seaweeds (*Ascophyllum nodosum*), plants like soybean, maize gluten, protein cake from vegetable oil industries and animal proteins (Ashmead, 1986).

### **1.4 Protein Lacto gluconate Nutrients:**

Several organic carbon (OC) rich formulations tailored with amino acids, gluconic and lactic acids blended with elemental Nitrogen (N), phosphorous (P), potassium (K), sulfur (S), calcium (Ca), magnesium (Mg), boron (B), copper (Cu), iron (Fe), molybdenum (Mo),

manganese (Mn) etc., were produced from research & development based biotech industries. These formulations were proved through bio-efficacy studies by several national and international agriculture universities and research laboratories on various crop systems and environmental conditions (Elisha *et al.*, 2014).

## 2. Material and Methods:

### 2.1 Preamble

The experiment will be conducted to study the efficacy of integrated usage of the recommended dose of fertilizers (N:72-80; P:24 K:20 kg/acre) and Lactate, Gluconate based organic formulations on Maize. The study will focus on growth and development in terms of plant growth and yield attributes.

### 2.2 Location

The experimental site Malla Reddy University is situated in Southern Telangana Agro Climatic Zone at longitude - 78° 46' 22.69" E (78.772971), latitude - 17° 21' 9.86" N (17.352743) with an altitude of 547 meters above Mean Sea Level (Fig. 1).




Fig. 1. Current location map of the field experiment

### 2.3 Crop and Variety

Maize variety, DHM-117 a short duration high yielding genotype suitable for *kharif* and *rabi* seasons was used for the field experiment.

## 2.4 Experimental details:

**Table 1. Experimental details of Maize crop**

S.No.	Particulars	Details
1.	Year of Experiment	2021-22
2.	Crop	Maize
3.	No. of Treatments	6
4.	No. of Replications	4
5.	Spacing	60×20 cm
6.	Experimental Design	Randomized Block Design
7.	Season	<i>spring -2021</i>
8.	Variety	DHM-117
9.	Location	Malla Reddy University, Telangana

### **Treatments** Details:

**T<sub>1</sub>**: Control (No fertilizer)

**T<sub>2</sub>**: 100 % RDF

**T<sub>3</sub>**: 75 % RDF + 25 % Sahasra Organic Formulations

**T<sub>4</sub>**: 50 % RDF + 50 % Sahasra Organic Formulations

**T<sub>5</sub>**: 25 % RDF + 75 % Sahasra Organic Formulations

**T<sub>6</sub>**: 100 % Sahasra Organic Formulations

Recommended Dose of Fertilizer (kg/acre) N: 72-80; P: 24; K: 20

## 2.5 Sahasra Organic Products:

Sahasra organic products i.e., Organic manure, Sage, Nutri and Poshak were formulated and supplied by Sahasra crop science, Hyderabad for testing the efficacy of the products for improving the soil health and yield.

**Table 2. Details of Sahasra Organic Formulations**

S.No.	Crop	Product	Dosage	Time of Application	Spraying Intervals
1.	Maize	Organic Manure	50 kg/acre	Basal Dose	Single Dose

			(Before sowing)	
Sage	3 ml/lit		20, 40, 60 DAS	Three Sprays
Poshak	3 ml/lit		30 & 40 DAS	Two Sprays
Nutri	2 ml/lit		20 & 40 DAS	Two Sprays

\*Coragen 10 DAS (for fall army worm)

### Spraying Schedule:

20, 40 & 60 Days After Sowing

### Note:

**Organic Manure** (50 kg/acre as Basal Application); **Sage** (3 ml/lit at 20-40-60 DAS); **Nutri** (2 ml/lit at 20 & 40 DAS); **Poshak** (3 ml/lit at 30 & 40 DAS); If micronutrient deficiency observed on leaves additional spray can be given.

Crop raised as per recommended package of practices of the University. All the recommended agronomic practices are uniformly followed for all treatments viz; hoeing, intercultural operations, weedicides application and irrigations. The soil physico chemical characteristics, microbial characteristics, plant growth parameters, yield attributes and post harvest parameters (Elisha *et al.*, 2014). For data recording five sample plants and cobs were identified randomly collected and stacked separately for sun drying and various observations for treatment evaluation were recorded as per the experimental design. The data were statistically analyzed by standard tools for interpretation of the results. Data recorded from the experimental identified plants in each sub-plot were randomly selected and taken the average weight of all the cobs weight per plant<sup>-1</sup>, no.of cobs per plant, cob length (cm), cob girth (cm), No. of kernels cob<sup>-1</sup>, grain (kg ha<sup>-1</sup>) and stover yield (kg ha<sup>-1</sup>).

### 2.6 Grain yield

The grain yield was noted by weighing the grains shelled from the cobs obtained from the central four rows of each sub-plot and converting it into kg ha<sup>-1</sup>.

### 2.7 Biological yield

It was recorded by weighing the sun dried plants along with ears obtained from central four rows of each sub-plot. The biological yield thus obtained in each sub-plot was converted into kg ha<sup>-1</sup>.

### 2.8 Harvest Index (%)

$$HI (\%) = \text{Grain yield} / \text{Biological yield} \times 100 \text{ (Reddy, 2004)}$$

### 2.9 Statistical analysis

The data obtained from the field experiment was subjected to statistical analysis. Wherever the treatments difference were significant, Critical differences were worked out at a 0.05 per cent probability level and the values were furnished (Jan *et al.*, 2009). The treatment differences that were not significant at five per cent were denoted as “NS”. A brief interpretation of the results was given at the end of each parameter.


### 3. Results and Discussion

#### 3.1 Plant Height (cm)

It is evident from (Table 3 and Fig. 2) that the increased plant height among all the treatments in the 6<sup>th</sup> leaf stage was observed in T<sub>2</sub> (100 % RDF) 45.8 cm, T<sub>3</sub> (75 % RDF + 25 % Sahasra Organic Formulations) 172.2 cm on silking stage, T<sub>2</sub> (100 % RDF) 108.4 cm on dough stage and in T<sub>3</sub> (75 % RDF + 25 % Sahasra Organic Formulations) 185.6 cm at maturity stage.

**Table 3. Plant height of maize crop under different stages of crop growth**

Plant height (cm)	6 <sup>th</sup> Leaf	Silk stage	Dough stage	Maturity
T <sub>1</sub>	34.1	130.8	82.4	142.6
T <sub>2</sub>	45.8	171.1	108.4	178.1
T <sub>3</sub>	42.6	172.2	107.4	185.6
T <sub>4</sub>	40.3	160.1	100.2	165.4
T <sub>5</sub>	38.1	143.9	91.0	158.1
T <sub>6</sub>	37.7	138.9	88.3	149.8
<b>C.D.</b>	<b>1.755</b>	<b>6.850</b>	<b>5.286</b>	<b>17.476</b>
<b>SE(m)</b>	0.577	2.252	1.738	5.475
<b>SE(d)</b>	0.816	3.185	2.458	7.743
<b>C.V.</b>	<b>2.904</b>	<b>2.947</b>	<b>2.171</b>	<b>5.675</b>



**Fig. 2. Plant height of maize crop under different stages of crop growth**

#### 3.4 Number of cobs per plant

It is clear from (Table 4.) that the cob-bearing capacity is one of the most important crops yield components. More or less cobs, it is the genetic character of any cultivar but some improvement can be expected owing to agronomic manipulations. The number of cobs per plant was not affected significantly by different treatment combinations. The highest number of cobs per plant was observed with the recommended dose of fertilizer treatment. (T<sub>2</sub>).

### 3.5 Cob length (cm)

As mentioned in (Table 4) that the highest cob length (20.7 cm) was recorded with T<sub>3</sub> (75 % RDF + 25 % SOF) followed by T<sub>2</sub> (100 % RDF) *i.e* 19.8 cm, while the least cob length has been recorded in T<sub>1</sub> control (16.5 cm) treatment. An increase in cob length might be due to adequate and uniform availability of nutrients during the crop growth which resulted in enhancement of the yield attributes Pathak *et al.* (2005) also observed similar results.

### 3.6 Cob girth (cm)

It is evident from (Table 4) that the maximum cob girth (14.5 cm) was recorded with T<sub>2</sub> (100 % RDF) followed by T<sub>3</sub> (75 % RDF + 25 % SOF) *i.e* 14.3 cm, while the least cob girth (11.7 cm) was recorded with T<sub>1</sub> (control)

### 3.7 Cob weight (g)

The highest cob weight (180.6 g) was recorded with T<sub>3</sub> (75 % RDF + 25 % SOF) followed by T<sub>2</sub> (100 % RDF) (175.9), while the lowest cob weight (128.3 g) was noted with T<sub>1</sub> (control).

### 3.8 Number of kernels cob<sup>-1</sup>

The highest number of kernels per cob (554.6) was recorded with T<sub>3</sub> (75 % RDF + 25 % SOF) followed by T<sub>2</sub> (100 % RDF) (550.7), while the lowest number of kernels per cob (457.4) was recorded with T<sub>1</sub> (control)

### 3.9 Grain yield (kg ha<sup>-1</sup>)

Significantly higher grain yield (7649.8 kg ha<sup>-1</sup>) was recorded with T<sub>2</sub> (100% RDF) which was statistically on par with T<sub>3</sub> (75 % RDF + 25 % SOF) (*i.e* 7241.3 kg ha<sup>-1</sup>), followed by T<sub>4</sub> which recorded 7,019.9 kg ha<sup>-1</sup> grain yield closely followed by T<sub>5</sub> and T<sub>6</sub> (6,763.6 and 6,179.1 kg ha<sup>-1</sup> respectively), while significantly lower grain yield (4808.2 kg ha<sup>-1</sup>) was recorded with T<sub>1</sub> (control). The maximum grain yield might be attributed due to the combined

effect of fertilizer along with organic formulations which escalated the availability of nutrients and transport of major nutrients like N, P and K which also resulted in increased plant height, no. of leaves and leaf area which in turn lead to increased production and translocation of photosynthates and yield attributes like cob length (cm), cob girth (cm), number of rows cob<sup>-1</sup>, cob weight (g), grain weight (g). The results are in a similar trend to the results of an increase in stover yield due to fertigation of microbes as reported by Abdelhamid *et al.* (2011), Baral and Adhikari (2013) and Meena *et al.* (2013). Sonawane *et al.* (2009) also reported better grain and dry fodder yield with combined application of nutrients due to improvement in yield parameters of maize crops. Application of organic manures because of combination with inorganic fertilizers not only enhanced the nutrient supply for higher grain yield but also alters the soil physical and chemical properties which favor better crop growth and yield (Prasad *et al.*, 2011).

### 3.10 Stover yield (kg ha<sup>-1</sup>)

Significantly higher stover yield (8859.8 kg ha<sup>-1</sup>) was recorded with T<sub>2</sub> (100% RDF) which was statistically on par with T<sub>3</sub> (75 % RDF + 25 % SOF) which is (8850.6 kg ha<sup>-1</sup>), followed by T<sub>4</sub>, T<sub>5</sub>, T<sub>6</sub> respectively (*i.e.* 8394.9, 8252.7, 8049.3 kg ha<sup>-1</sup>), while the lowest stover yield (7835.6 kg ha<sup>-1</sup>) was recorded with T<sub>1</sub> (control). The increase in stover yield might be due to the combined effect of fertilizer along with organic formulations which enhanced the availability of nutrients like N, P and K, which increased plant biometric observations such as plant height, the number of leaves and leaf area which in turn lead to higher production and translocation of photosynthates and more dry matter production plant<sup>-1</sup>. The results are in a similar trend were the increase of stover yield due to fertigation of microbes as reported by Abdelhamid *et al.* (2011) through fertigation, by seed inoculation in maize by Baral and Adhikari (2013) and Meena *et al.* (2013). Manjhi *et al.* (2014) reported the maximum grain and stover yield under integrated treatment. Karki *et al.* (2005) also recorded similar results in respect of grain and stover yield of maize.

### 3.11 Test weight (g)

The highest test weight (31.8 g) was observed with T<sub>3</sub> (75 % RDF + 25 % SOF) which was on par with T<sub>2</sub> (100% RDF) (*i.e.* 31.5 g) followed by T<sub>4</sub>, T<sub>5</sub> and T<sub>6</sub> respectively (*i.e.* 30.4, 30.4, 30.2 g), while the lowest test weight (29.1 g) was recorded with T<sub>1</sub> (control). Due to the application of organic formulations plants received maximum nutrients throughout their

growth period and nourished sufficiently, which resulted in a maximum 100 kernel weight. Similar findings of more grain weight with imposing integrated treatment in Maize crop was reported by Cheema *et al.* (2010).


### 3.12 Harvest index (%)

A significantly more harvest index (46.34) was noted with T<sub>2</sub> (100% RDF) followed by T<sub>3</sub> (75 % RDF + 25 % SOF which is (45.54), while the lowest harvest index (38.03) was observed with T<sub>1</sub> (control).

**Table 4. Yield attributes of maize (*Zea mays* L.) as influenced by different doses of fertilizers and Sahasra Organic formulations.**

Treatments	No.of Cobs/plant	Cob length (cm)	Cob girth (cm)	Cob weight (g)	No. of Kernels/Cob	Grain yield (kg ha <sup>-1</sup> )	Stover yield (kg ha <sup>-1</sup> )	Test weight (g)	Harvest Index (%)
T <sub>1</sub>	9.6	16.5	11.7	128.3	457.4	4,808.2	7,835.60	29.1	38.03
T <sub>2</sub>	11.9	19.8	14.5	175.9	550.7	7,649.8	8,859.87	31.5	46.34
T <sub>3</sub>	11.8	20.7	14.3	180.6	554.6	7,241.3	8,850.63	31.8	45.00
T <sub>4</sub>	10.4	18.8	13.9	167.6	526.3	7,019.9	8,394.90	30.4	45.54
T <sub>5</sub>	10.2	18.0	13.2	157.3	509.1	6,763.6	8,252.73	30.4	45.04
T <sub>6</sub>	10.4	16.9	12.1	142.2	494.2	6,179.1	8,049.30	30.2	43.43
SE.m±	0.388	0.305	0.186	2.457	7.546	136.68	31.179	0.395	1.172
CD (P=0.05)	1.238	0.975	0.593	7.842	24.085	415.75	99.515	1.260	3.741

T<sub>1</sub>: Control (No fertilizer), T<sub>2</sub>: 100 % RDF, T<sub>3</sub>: 75 % RDF + 25 % Sahasra Organic Formulations, T<sub>4</sub>: 50 % RDF + 50 % Sahasra Organic Formulations, T<sub>5</sub>: 25 % RDF + 75 % Sahasra Organic Formulations, T<sub>6</sub>: 100 % Sahasra Organic Formulations. Recommended Dose of Fertilizer (kg/acre) N: 72-80; P: 24 K: 20



**Fig. 3. Grain yield of maize as influenced by different doses of fertilizers and Sahasra Organic Formulations.**

**Table 5. Economics of maize (*Zea mays* L.) as influenced by different doses of fertilizers and Sahasra organic formulations**

Treatment	Gross returns	Net returns	B:C
T <sub>1</sub>	79958.975	51308.98	2.8
T <sub>2</sub>	123606.491	87116.49	3.4
T <sub>3</sub>	117469.759	82466.76	3.4
T <sub>4</sub>	113693.399	79477.40	3.3
T <sub>5</sub>	109706.733	76627.73	3.3
T <sub>6</sub>	100736.175	68811.18	3.2

- Grain cost Rs 15 /kg ; Stover cost = Rs 1/kg

From this study, application of integrated application of (75 % RDF + 25 % Sahasra Organic Formulations) was found to be on par with the 100% recommended dose of fertilizer was recorded with the highest grain yield (7649.8 kg ha<sup>-1</sup>) and stover yield (8859.8 kg ha<sup>-1</sup>), gross returns (₹ 123606.5 ha<sup>-1</sup>), net returns (₹ 87116.49 ha<sup>-1</sup>) with B: C ratio 3.4. and economical with the highest net returns (₹ 82466.8 ha<sup>-1</sup>) and B: C ratio (3.4) in both the treatments as mentioned in Table 5.

#### 4. Conclusion

From this study, it can be concluded that the application of recommended dosages of chemical fertilizers in combination with Sahasra organic nutrients (75 % RDF + 25 % Sahasra Organic Formulations) recorded higher yields. In addition to this, the application of Sahasra organic nutrients enhances soil health by reducing the cost of cultivation to the farmers in respect of total nitrogen, phosphorus and potassium content in the soil.

## 5. Acknowledgments

The author acknowledge the Department of Agronomy, School of Agricultural Sciences, Malla Reddy University for extending the facilities to conduct this field experiment smoothly. I would like to extend my gratitude to the Research and Development team Sahasra Crop Science for sharing their pearls of wisdom and for financial support during the course of this research.

## 6. Ethical Approval

The authors declare that all the experiments were conducted according to the current laws of the country in which they were performed.

## 7. Conflict of Interest

The Author (s) declare (s) that there is no conflict of interest.

## 8. References:

Abdelhamid, M.T., Selim, E.M and El-Ghamry, A.M. 2011. Integrated effects of bio and mineral fertilizers and humic substances on growth, yield and nutrient contents of fertigated cowpea (*Vigna unguiculata* L.) grown on sandy soils. *Journal of Agronomy*. 10 (1): 34-39.

Ashmead, H. D. 1986. Foliar Feeding of Plants with Amino Acid Chelates. Park Ridge, New Jersey USA: Noyes Publications: S662.5-F65.

Baral, B.R and Adhikari, P. 2013. Effect of Azotobacter on growth and yield of maize. *SAARC Journal of Agriculture*. 11 (2): 141-147.

Cheema, M.A., Farhad, W., Saleem, M.F., Khan, H.Z., Munir, M.A., Wahid, A., Rasul, F. and Hammad, H.M. 2010. Nitrogen management strategies for sustainable maize production. *Crop Environment*. 1: 49-52.

Elisha, P., Rajesh Kumar Sharma and Varma, V. 2014. Studies on organic acid based biotech nutrients to enhance soil organic carbon (SOC) and grain yield in Maize (*Zea mays*) crop. *International Journal of Science Engineering and Advance Technology*. 2 (10): 1-10.

- Garima and Pant, K.S. 2018. Effect of integrated nutrient management and tree spacing on production potential of maize (*Zea mays*) under poplar-based agroforestry system. *International Journal of Current Microbiology and Applied Sciences*. 6: 2692-2697.
- Ghosh, D., Brahmachari, K., Brestic, M., Ondrisik, P., Hossain, A and Skalicky, M. 2020. Integrated weed and nutrient management improve yield, nutrient uptake and economics of maize in the rice maize cropping system of eastern India. *Agronomy*. 10: 1906-1915.
- Goldstein, A.H., Kogers, R.D and Mead, G. 1993. Mining by the microbe. *Nature Biotechnology*. 11: 1250-1254.
- Jan, M. T., P. Shah, P. A. Hoolinton, M. J. Khan and Q. Sohail. 2009. Agriculture research: Design and Analysis, A monograph. Dept. of Agronomy, KP Agric. Uni. Peshawar, Pakistan.
- Karki, T.B., Kumar, A and Gautam, R.C. 2005. Influence of integrated nutrient management on growth, yield, content and uptake of nutrient and soil fertility status in maize (*Zea mays*). *Indian Journal of Agricultural Sciences*. 75: 682-685.
- Kim, K.Y., Jordan, D and McDonald, G.A. 1997. Solubilization of hydroxyapatite by *Enterobacter agglomerans*, phosphate solubilizing bacteria and microbial activity in soil: Effect of carbon sources. *Soil Biology Biochemistry*. 30: 995-1003.
- Manjhi, R.P., Yadav, M.S and Thakur, R. 2014. Effect of integrated nutrient management on crop productivity and changes in soil fertility in maize-wheat cropping sequence. *Indian Journal of Agronomy*. 59: 371-376.
- Mauria S., Gupta N.P., Zaidi, P.H and Singh, N.N. 1998. Maize research in India-Progress and future challenges. *Indian Farming*. 48 (1): 37-41.
- Meena, M.D., Tiwari, D.D., Chaudhari, S.K., Biswas, D.R., Narjary, B., Meena, A.L and Meena, R.B. 2013. Effect of biofertilizer and nutrient levels on yield and nutrient uptake by maize (*Zea mays* L.). *Annals of Agri-Bio Research*. 18 (2): 176-181.
- Nirupma Singh, R.Ambika Rajendran, Meena Shekhar, S.L. Jat, Ramesh Kumar & R.Sai Kumar. Rabi Maize Opportunities Challenges, Directorate of Maize Research, Pusa Campus, New Delhi -110 012, Technical Bulletin, No. 9: 32-42.
- Okada, S. 1998. Lactic acid Bacteria for developing a clean environment. *Biseibutsu* (in Japanese). 4 (1): 56-70.
- Pathak, S.K., Singh, S.B., Jha, R.N and Sharma, R.P. 2005. Effect of nutrient management on nutrient uptake and changes in soil fertility in maize (*Zea mays*)-wheat (*Triticum aestivum*) cropping system. *Indian Journal of Agronomy*. 50: 269-273.
- Prasad, J., Karmakar, S., Kumar, R and Mishra, B. 2011. Influence of integrated nutrient management on yield and soil properties in maize-wheat cropping system in an Alfisol of Jharkhand. *Journal of Indian Society of Soil Science*. 58: 200-204.

Reddy, S. R. 2004. Principles of crop production. 2<sup>nd</sup> Ed. Kalyani Publishers, New Delhi, India. pp 46.

Sonawane, D.A., Goethe, R.M., Pawar, V.Y., Jadhav, A.G and Wadile, S.C. 2009. Performance of maize (*Zea mays* L.) under integrated nutrient management. *Bioinfolet*. 6: 270-280.

Walia, S.S and Kler, D.S. 2010. Effect of organic and inorganic sources of nutrition on growth, macro and micronutrient uptake in maize under maize–wheat sequence. *Indian Journal of Ecology*. 37: 27-29.