

Response of plant growth regulators on growth , yield and quality of okra cv. Kashi Kranti

ABSTRACT

The present work was conducted to evaluate the response of okra cv. Kashi Kranti to plant growth regulators and micronutrients at the vegetable seed production area, Bihar Agricultural University, Sabour, Bhagalpur, Bihar, India during Kharif season. The experiment was laid out in RBD (Factorial) and replicated thrice. The experiment was framed with two levels of PGRs, GA₃ (100 ppm, 150 ppm) and NAA (100 ppm and 150 ppm), sprayed twice, first at twenty days after sowing and second at forty days after sowing. The maximum number of leaves (18.13), number of branches (3.48), plant height (104.17 cm), number of fruit per plant (23.39), fruit fresh weight (17.95g) and fruit yield (130.88q/ha), 1000 seed weight (78.49g) chlorophyll-a (1.84), chlorophyll-b (0.81), zinc content (30.81 ppm), boron (22.55 ppm), copper (18.24 ppm) and germination (85.48%) was observed on application of GA₃-100ppm, which was found to be significantly superior in most of the cases. Maximum net return highest benefit cost ratio (2.67) was also obtained on application of GA₃100 ppm

Key words: Plant growth regulators, okra

INTRODUCTION

Okra, *Abelmoschus esculentus* (L.) Moench, is grown throughout the tropical and warm temperate regions of the world for its fibrous pods eaten as a vegetable. Plant growth regulators play very important role in crop production. These have significant role in crop growth along the increase in longitudinal area, increase number of branches, early flower initiation, fruit set, fruit quality and subsequently contributes towards higher production when applied at various concentration. Although plant growth regulators have great potential for growth improvement but their application has to be planned sensibly in terms of optimal dose, stage of application, crop specificity and seasons. Even the same growth regulator at different doses can bring about different results. These are considered as new

generations agrochemicals after fertilizers, insecticides and herbicides. Plant physiologists have recognized five well defined groups of plant growth substances viz., auxins, GA's, cytokinins, inhibitors (abscisic acid) and senescent hormone ethylene. Therefore, the use of plant growth regulators has led to intensive scientific activity for their commercial exploitation. Keeping these points in view, the present study was undertaken to ascertain the most suitable concentration of GA₃ and NAA on growth yield and quality of okra cv. Kashi Kranti.

MATERIALS AND METHODS

The investigation was conducted at Bihar Agriculture College Sabour, Bhagalpur, Bihar, India during Kharif season. The treatments included two plant growth regulators at two different concentrations (GA₃ 100ppm, 150 ppm and NAA 100ppm, 150 ppm). The trial was laid out in factorial Randomized Block Design having three replications. The plants were sown at a distance of 50 cm either way. Recommended package of practices were followed to raise the crop. Spraying of GA₃ and NAA was done twice, first at twenty days after sowing and second at forty days after sowing. Observations were recorded on the traits namely, number of leaves per plant, number of branches per plant, plant height (cm), number of days setting first flowering, number of fruits per plant, fruit fresh weight (g), average fruit dry weight (g), average fruit diameter (cm), fruit yield (q/ha), 1000 seed weight(g), seed yield (g/plant), chlorophyll-a(mg/plant), chlorophyll-b(mg/plant), zinc content(%), boron content(%), copper content(%) and B:C ratio was worked out. The statistical analysis of the data was carried out by the method of analysis of variance as suggested by Panse and Sukhatme (1967). Comparison of the genotypes was made with the help of critical differences (C.D.).

RESULTS AND DISCUSSION

Application of plant growth regulators significantly increased growth, yield and quality parameters. Increase in number of leaves (18.13) was noticed with GA₃ 100 ppm which was significantly superior to all the treatments followed by GA₃150 ppm. The maximum number of branches (3.48) and plant height (104.17) were observed with application of GA₃ 100 ppm. The growth regulator GA₃ is recognized as a growth promoter which stimulates the rapid cell elongation in the meristematic zone of vegetative plant organs. Similar findings were reported by

Chandiniraj et al. (2016), Bhagure and Tambe (2011), Patil and Patel (2010), Mukesh et al. (2009) and Naga et al (2010) in Okra. The plant receiving growth promoter GA₃100 ppm took minimum days (41.62) to first flowering which was significantly superior to all other treatments. The growth regulators might have influenced the physiological regulation of flower formation in the plants, possibly influencing the timing of anthesis mechanism (Das and Prusty, 1972). The maximum number of fruits per plant (23.39) and fruit fresh weight (17.95) and thousand seed weight (78.49) was associated with foliar application of GA₃-100ppm and was found to be significantly superior to other treatments. However, minimum was observed in the absence of PGRs in each case. The highest gross income, net income with the maximum benefit: cost ratio of (3.14) was obtained by foliar application of GA₃-100 ppm. (Table-1).The result gained on economic aspect is also in conformity with the result reported earlier by Kumar et al. (2005). Comparisons between two growth regulators revealed that GA₃ treatments had more positive influence on growth and yield parameters than NAA. It was obvious from the results that GA₃-100 ppm recorded best performance towards growth and yield attributes among all other treatments. It was also noticed that GA₃ treatments have more pronounced effects on the data recorded in comparison with respective NAA treatments. It might be due to that GA₃ could be involved in many aspects of plant growth and development, such as cell enlargement, internodes elongation, stimulated RNA and protein synthesis and thereby leading to enhanced growth and development, flowering and fruiting (Yamaguchi et al., 2000). The positive influence of plant growth regulators on growth and yield of okra was in agreement of Kokare et al., 2006 and Nawalkar et al., 2007. The results of increased seed yield by GA₃ were in accordance of Singh et al., 2012 and Sanganagoud et al., 2014. Marie et al., 2007 has also reported an increase in number of seeds per fruit under GA₃ treatment.

As for the quality characters, maximum chlorophyll-a (1.84) was associated with foliar application of GA₃-150 ppm and which was at par with other levels of different PGRs under study. However, it was significantly superior over the control. These results supported by Kumar et al. (2009) and Yadav et al. (2003). Application of plant growth regulators significantly enhanced zinc content in fruit of okra over the other treatments. The foliar application of GA₃100 ppm registered the highest zinc content giving (30.81) in fruit. The increase in zinc content might be due to the fact that GA₃ must have accelerated the physiological process like synthesis of carbohydrate resulting in increase in zinc content. Boron (22.55) and copper (18.24) content showed significant variation on application of different levels of PGRs being maximum with

foliar application of GA₃100 ppm which in turn was found to be significantly superior to other treatments. The foliar application of GA₃-100 ppm significantly induced highest germination percentage (85.48) which was significantly superior to all other levels (Table-2). This might be due to the fact that GA₃ must have induced production of reducing sugar from the starch as a result of increase in water soluble amylolytic activity of endosperm and enhanced some metabolic activities in embryo seeds which have caused quicker germination and early establishment in plant. These results are supported by Bhagure and Tamba (2013) and Singh (1998). Maity et al. (2016) have reported that gibberellins (GA₃) and auxins (IBA) have significant effect in increasing growth and development and yield of okra with increasing concentration of both. However, GA₃ is found to be a superior growth regulator as compared to respective doses auxins.

Conclusion

Thus it is concluded that gibberellic acid and NAA have significant effect in increasing growth and development and yield of okra with increasing concentration of both. However, GA₃ is found to be a superior growth regulator as compared to respective doses of NAA. GA₃-100 was responsible for overall in growth, yield and quality of okra. The highest gross income, net income with the maximum benefit: cost ratio of was also obtained by foliar application of GA₃-100 ppm.

REFERENCES

- Bhagure YL, and Tambe TB (2011) Effect of seed soaking and foliar sprays of plant growth regulators on germination, growth and yield of okra [*Abelmoschus esculentus* (L.) Moench] var. Parbhani Kranti. The Asian Journal of Horticulture, 8(2): 399-402.
- Bhagure, Y. L. and Tambe, T. B. (2013). Effect of seed soaking and foliar sprays of plant growth regulators on germination, growth and yield of okra [*Abelmoschus esculentus* (L.) Moench] var. Parbhani Kranti. Asian J. Hort. 8 : 399-40
- Chandiniraj A, Holebasappa K, Hore JK, and Chattopadyay N (2016) Growth and yield of chilli (*Capsicum annuum* L.) as influenced by different growth regulators. The BIOSCAN, The Journal an International Quarterly Journal of life sciences, 11(1): 385-388.
- Das, R. C. and Prusty, S. S. (1972). Growth regulators effect on seed treated brinjal plants (*Solanum melongena* L.) with relation to vegetative development. Indian J. Hort., 29: 334-335

- Kokare RT, Bhalerao RK, Prabu T, Chavan SK, Bansode AB, and Kachare GS (2006) Effect of plant growth regulators on growth, yield and quality of okra (*Abelmoschus esculentus* L.). *Agricultural Science Digest* 26(3): 178-181.
- Kumar, S., Chankhar, S.K. and Rana, M. K. (2009). Response of okra to zinc and boron micronutrients. *Vegetable Science*, 36: 327- 331.
- Kumar M, and Sen NL (2005) Effect of zinc, boron and gibberellic acid on yield of okra (*Abelmoschus esculentus* (L.) Moench). *Indian Journal of Horticulture* 62 (3): 308- 309.
- Kumar, S., Chankhar, S.K. and Rana, M. K. (2009). Response of okra to zinc and boron micronutrients. *Vegetable Science*, 36: 327- 331.
- Maity, U., Dutta, P. and Layek, B. (2016). Effect of Plant Growth Regulators on Growth, Yield and Quality of Okra [*Abelmoschus esculentus* (L.) Moench] *Journal of Agroecology and Natural Resource Management* p-ISSN: 2394-0786, e-ISSN: 2394-0794, Volume 3, Issue 3; July-September 2016 pp. 251-253 © Krishi Sanskriti Publications <http://www.krishisanskriti.org/Publication.htm>
- Marie AI, Ihsan A, and Salith SH (2007) Effects of sowing date, topping and some growth regulators on growth, pod and seeds yield of okra [*Abelmoschus esculentus* (L.) Moench]. *Afr Crop Sci Conf Proc* 8:473-478.
- Mukesh LC, Tirakannanavar S, Ahmed S, Munikrishnappa PM, and Mastiholi AB (2009) Effect of growth regulators and method of application on growth and fruit yield in paprika chilli (*Capsicum annuum* L.) cv. Kt-PI-19. *Seed Research*, 37(1-2): 14-19.
- Naga, S. K., Swain, S. K., Verma, S. and Raju, B. (2010). Effect of foliar application of micronutrients on growth parameters in tomato (*Lycopersicon esculentum* Mill.) *Discourse journal of Agriculture and food science*, 1(10):146-151.
- Nawalkar LR, Khiratkar SD, Badge SA, Chopde NK, and Dadgal SS (2007) Effect of biofertilizers and growth regulator with reduced doses of NPK on growth and yield of okra. *Journal of Soils and Crops* 17(1): 145-149.
- Panse VG, and Sukhatme PV. *Statistical methods for agricultural workers*, ICAR, New Delhi. 1967, 152-161.
- Patil DR, and Patel MN (2010) Effect of seed treatment with GA₃ and NAA on growth and yield of okra (*Abelmoschus esculentus* L. Moench). cv. GO-2, *Asian Journal of Horticulture*, 5 (2): 269-272.
- Sanganagoud PR, Chaitanya HS, and Nagesh L (2014) Effect of plant growth regulators and fruit picking on growth characters of okra [*Abelmoschus esculentus* (L.) Moench] at coastal Karnataka. *Environ Ecol* 32(3): 896-900.

- Singh J, Singh BK, Singh AK, Manju Vani V, and Singh SP (2012) Performance of plant growth regulators on yield and yield attributing characters of okra [*Abelmoschus esculentus* (L.) Moench]. *Environ Ecol* 30(4): 1333-1335.
- Singh, P. V. and Kumar, J. (1998). Effect of gibberllic acid as a pre-sowing seed treatment and different level of nitrogen on germination, growth, flowering and yield of okra. *Indian Journal Agricultural Research*, 32(1):31-36.
- Yadav, P.V.S., Tikko, A. and Sharma, N. K. (2003). Effect of zinc and boron application on growth, flowering and fruiting of tomato (*Lycopersicon esculentum* Mill.). *Haryana Journal of Horticulture Science*, 39(1-2):104-107
- Yamaguchi S, and Kamiya Y (2000) Gibberellin biosynthesis: its regulation by endogenous and environmental signals. *Plant Cell Physiology* 41(3): 251-257.

UNDER PEER REVIEW

Table-1: Effect of plant growth regulators on growth and yield characters of okra cv. Kashi Kranti.

Treatments	No. of leaves/Pl. plant	No. of branches per plant	Pant height at maturity	Days to first Flowering	No. of fruits per plant	Fruit fresh weight	Average fruit diameter	Fruit yield (q/ha)	1000 seed (g)	Seed yield (g/plant)	B:C
Control	15.5	2.69	97.78	43.57	20.23	15.51	1.45	120.81	74.8	62.21	2.47
NAA-100ppm	16.32	3	99.75	43.14	22.31	17.36	1.52	126.68	76.15	64.67	2.57
NAA-150ppm	16.43	2.89	100.23	43.67	22.1	17.29	1.54	123.61	76.35	64.69	2.5
GA3-100ppm	18.13	3.48	104.17	41.62	23.39	17.95	1.66	130.88	78.49	67.33	2.67
GA3-150ppm	17.3	3.08	103	43.38	21.14	17.37	1.54	123.85	77.26	63.48	2.39
CD	0.71	0.08	3.03	1.35	0.63	0.44	0.045	3.04	0.99	1.95	
CV	10.49	6.35	6.9	7.21	6.6	5.94	6.79	6.68	2.97	6.95	

Table-2: Effect of plant growth regulators on quality characters of okra cv. Kashi Kranti.

parameters	Chlorophyll a	Chlorophyll b	Zinc (ppm)	Boron (ppm)	Copper (ppm)	Germination%
Control	1.28	0.77	24.28	15.28	15.1	83.14
NAA-100ppm	1.69	0.78	29.14	21.72	16.05	83.67
NAA-150ppm	1.57	0.79	28.52	20.3	15.67	84.38
GA3-100ppm	1.81	0.81	30.81	22.55	18.24	85.48
GA3-150ppm	1.84	0.8	27.1	21.2	16.05	84.57
CD	0.33	0.01	1.31	0.58	0.79	1.17
CV	8.87	9.41	10.83	6.66	11.22	4.04

UNDER PEER REVIEW