

DEVELOPMENT OF FRESH VALUE ADDED GUAVA (*Psidium guajava* L.) AND PAPAYA (*Carica papaya* L.) NECTAR AND STANDARDIZATION OF ITS RECIPE

ABSTRACT:

A lab experiment was conducted during winter season of 2021 at Post Harvest technology Laboratory department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology & Sciences, Prayagraj, and Uttar Pradesh, India to determine the “**Development of fresh value added guava (*Psidium guajava* L.) and papaya (*Carica papaya* L.) nectar and standardization of its recipe**”. The experimental material consisted of ten treatments with combination of different recipes including two standard recipes without blending of guava and papaya with varying levels of TSS for nectar. The treatment combinations were 10 and replicated thrice under Completely Randomized Design.

The fruits of guava and papaya were analysed for physic-chemical characteristics. The blended nectar prepared from different recipes and blending ratio were also analysed periodically at 4 days interval for their various chemical constituents. The blended nectar organoleptically evaluated by adopting 9 point Hedonic rating scale and observations were recorded for their chemical changes during storage for 12 days. A critical analysis of physical composition of guava and papaya fruits revealed that average fruit weight was recorded 163.65 g and 1610 g, pulp weight 158.20 g and 1430 g, seed weight 5.45 g and 10.50 g, weight of non-edible waste 7.13 g and 82 g and pulp seed ratio 29.02 and 136.19, respectively. Among the chemical composition, the TSS was recorded 12.31% and 8.36%, acidity 0.43% and 0.38%, ascorbic acid 243mg/100 ml and 66.62 mg/100 ml, pH 4.62 and 4.20, respectively. Among various recipe tried in this investigation, the nectar prepared from the treatment T4 (60% Guava: 40% Papaya: 18% TSS: 0.3% acidity) recorded highest TSS, Ascorbic acid, organoleptic score with respect to aroma, taste and overall acceptability. Whereas, the nectar containing the recipe T6 (45% Guava: 55% Papaya) recorded highest pH. During storage of nectar the acidity, TSS, showed an increasing trend with increasing period of storage (0 to 12 days) under ambient condition. While, there was a decreasing trend of ascorbic acid, pH and organoleptic score during storage period upto 12 days of storage under ambient condition.

Key words: *Guava-Papaya nectar, citric acid, no preservatives, recipe.*

INTRODUCTION

Fruits and vegetables contain many vitamins and minerals that are good for your health. These include vitamins A (beta-carotene), C and E, magnesium, zinc, phosphorous and folic acid. Fruit juice is a popular choice of beverage for both adults and children because of its attractive taste and color, and

the fact it is associated with many health benefits. India is a country, well-known for its tradition and culture. Syrup or Sharbet are offered to guests and is an important homemade soft drink. Similarly, fruit juice and beverages also hold an important position due to their richness in essential minerals, vitamins and other nutritive constituents. Guava (*Psidium guajava* L.) is one of the most important subtropical fruit crops. It belongs to family Myrtaceae. Guava is a native of tropical America perhaps from Mexico and Peru. It is widely distributed all over the equatorial regions of the tropical and subtropical climate. Guava can be grown easily in region with tropical and subtropical climate. Papaya (*Carica papaya* L.) is one of the tropical fruit crops. It belongs to family Caricaceae. The papaya is the plant *Carica papaya*, one of the 22 accepted species in the genus *Carica* of the family Caricaceae. Its origin is in the tropics of the Americas, perhaps from Central America and southern Mexico. Papayas grow in tropical climates and are also known as papaws or pawpaw. Their sweet taste, vibrant colour, and the wide variety of health benefits they provide make them a popular fruit.

The various products such as jam, jelly, candy, nectar, puree, concentrate slab, toffee, tutti-fruity, freeze dried chunk, dried rolls, dried slices and pickles can be prepared from papaya processing. Nectar juice is also called nectar drink, or pulp juice. It refers to the juice containing fruit pulps and flesh. Nectar juice is made of original fruit pulps or concentrated pulps, added with sugar and acid agent. The acidity can be increased after the storage in nectar, which is similar found in Bal. *et.al* (2014). It contains a large quantity of fruit pulps, fruit fibers and pectin, which can be drunk together with the juice. As the nectar juice reserve some solid content in fruit peels and pulps, it presents more of the fruit original taste, color and nutrition. To be general, the production of nectar juice includes: raw material cleaning, cutting, destoning, heating, pulping, mixing with sweet water and additives, colloid milling, degassing, filling, sealing, sterilization, finished products. Determine the ascorbic acid by 2, 6 – dichlorophenolindophenols dye method Ranganna (1997).

MATERIALS AND METHODS


Raw Material


Healthy and ripen guava and papaya, free from disease, pest and insects were randomly selected and purchased from fruit mandi in Prayagraj (U.P). Other ingredients like sugar, citric acid were brought from local shops in prayagraj.

Preparation of blended nectar of guava and papaya

Ripen fruits were selected for the preparation of blended nectar from guava and papaya. The fruits were washed with clean water to remove dirt and dust particles. The guava fruits were cut into small pieces with the help of knife, after that little amount of water was added and crushed it in mixer-grinder, while the mature and ripened papaya fruit were peeled from peeler and then cut into two equal halves and then seeds were removed. The papaya pulp was obtained by crushing in grinder. Straining was done by with the help of sieve. After extraction of pulp, 20 per cent for nectar was taken as per required ratio for blending of guava and papaya. Sugar syrup was prepared by heating required amount of sugar and water to adjust the total soluble solids as per requirement in different recipe for nectar. The acidity was maintained to 0.3 per cent by the addition of required amount of citric acid in the sugar syrup. The blended guava and papaya pulp was mixed with sugar syrup to obtain final product. The prepared nectar was again filtered by sieving through a muslin cloth to obtain a product of uniform consistency. The product was poured into hot, sterilized bottles of 200 ml capacity and packed air-tight. The filled bottles were pasteurized in boiling water till the temperature of product reaches 100°C. It took about 15 minutes to attain required temperature. The bottles of nectar was kept at ambient condition for further studies up to 12 days.

FLOW CHART OF GUAVA-PAPAYA NECTAR





TREATMENT DETAILS

Treatment	Treatment Combinations
T ₀	Guava Juice (100%) + Papaya juice (0%) + Sugar (110g) + citric acid (2.6g)
T ₁	Guava Juice (25%) + Papaya juice (75%) + Sugar (115g) + citric acid (2.6g)
T ₂	Guava Juice (50%) + Papaya juice (50%) + Sugar (105g) + citric acid (2.6g)
T ₃	Guava Juice (75%) + Papaya juice (25%) + Sugar (105g) + citric acid (2.6g)
T ₄	Guava Juice (60%) + Papaya juice (40%) + Sugar (105g) + citric acid (2.6g)
T ₅	Guava Juice (70%) + Papaya juice (30%) + Sugar (105g) + citric acid (2.6g)
T ₆	Guava Juice (45%) + Papaya juice (55%) + Sugar (105g) + citric acid (2.6g)
T ₇	Guava Juice (40%) + Papaya juice (60%) + Sugar (105g) + citric acid (2.6g)
T ₈	Guava Juice (35%) + Papaya juice (65%) + Sugar (105g) + citric acid (2.6g)
T ₉	Guava Juice (0%) + Papaya juice (100%) + Sugar (90g) + citric acid (2.6g)

Physico-chemical and organoleptic quality of guava-papaya nectar.

Titration acidity (%)

Titration acidity was determined by titrating a known quantity of sample (10ml) of homogenized sample was taken and made up to 100 ml volume in a volumetric flask. The contents were filtered through Whatman No. 1 filter paper. An aliquot of 10 ml was taken for titration against 0.1N NaOH using phenolphthalein as an indicator. The turn of aliquot to light

pink colour which persists for 15 seconds was considered as an end point. The titratable acidity was estimated in terms percent citric acid (Ranganna, 1986).

pH

The sample was taken into the beaker. Then the pH was noted with the help of an electronic pH meter.

Total soluble solids (%)

The percentage of total soluble solids was determined by using hand refractometer (Erma, Japan) by placing a drop of the filtered juice on the prism of the refractometer and observing the coincidence of shadow of the sample with the reading on the scale and expressed as °Brix.

Ascorbic acid (mg/100g)

Determination of ascorbic acid was done by 2, 6 – dichlorophenolindophenols dye method as described by Ranganna (1997). A known quantity of sample was blended with 3 per cent metaphosphoric acid (HPO₃) to make the final volume of 100 ml and then filtered. A known quantity of aliquot was titrated against 0.025 per cent 2, 6 - dichlorophenol indophenols dye to a pink colour end point. The ascorbic acid content of the sample was calculated taking into consideration the dye factor and expressed as mg Ascorbic acid per 100g.

Organoleptic evaluation

To assess the consumer preference, sensory evaluation of the experimental sample was conducted at different intervals by panel of 7-8 judges. The sample were evaluated for color and appearance, taste, aroma and overall acceptability.

RESULTS AND DISCUSSION

Titration Acidity (%)

The increase in acidity of nectar during storage might be due to formation of organic acids by ascorbic acid degradation as well as progressive decrease in the pectin content. It is also due to formation of acids from sugar. the significant highest titratable acidity (0.59%) was noticed under the treatment T₉ (0% Guava juice: 100% Papaya juice). While, minimum (0.35%) acidity was observed with T₄ (60% Guava juice: 40% Papaya juice). The results are also in conformity with the findings of Bal *et al.* (2014), who reported a significant increase in acidity of guava nectar during storage.

pH

The increased acidity and TSS under all the recipe treatments during storage had a corresponding decrease in pH. During the storage, maximum pH value was observed (4.81) with the T₆ (45% Guava juice: 55% Papaya juice). While, minimum pH was observed (4.07) with the treatment T₉ (0% Guava juice: 100% Papaya juice). The present findings are in agreement with Sasikumar (2015), who observed significant decrease in pH during storage of functional beverages from Aloe-vera juice blended with bael fruit juice.

Total Soluble Solids

This gradual increase in total soluble solids during the storage period (12 days) might be due to conversion of polysaccharides into sugars during hydrolysis process. The total soluble solids content was found to be maximum (18.45%) with the treatment T₄ (60% Guava juice: 40% Papaya juice). While, minimum TSS content was recorded (15.73%) with the treatment T₇ (40% Guava juice: 60% Papaya juice). Similar findings were also reported by Deka *et al.* (2004), who observed that total soluble solids showed an increasing trend throughout the storage period.

Ascorbic Acid

The reduction in ascorbic acid during the storage period due to oxidation by trapped oxygen in packaging container, which results in formation of dehydroascorbic acid. The maximum ascorbic acid was recorded (33.70mg/100ml) with the treatment T₄ (60% Guava juice: 40% Papaya juice). While the minimum ascorbic acid (31.24mg/100ml) was recorded under treatment T₀ (100%Guava juice: 0% Papaya juice). Similar results were also noted by Mall and Tondon (2007), in guava-aonla blended beverage.

Organoleptic evaluation

In the organoleptic evaluation such as color and appearance, taste, aroma and overall acceptability. Sensory scores for treatment T₄ (60% Guava juice: 40% Papaya juice) were found to be highest in all parameters of organoleptic attributes. The highest overall acceptability score (7.03) indicated that it was well received by the judges.

Shelf life of guava-papaya nectar

From the shelf life point of view during the storage period, the highest was recorded in T₄ (60% Guava juice: 40% Papaya juice) 11.23. While minimum was recorded in T₉ (0% guava juice: 100% papaya juice) 8.20.

Table 1: Changes in acidity, pH, and TSS in the guava-papaya nectar during storage.

Treatments	Acidity (%)				pH				TSS (°Bx)			
	Storage period (days)				Storage period (days)				Storage period (days)			
	0	4	8	12	0	4	8	12	0	4	8	12
T ₀	0.33	0.34	0.39	0.43	5.03	4.92	4.69	4.66	17.37	17.39	17.42	17.46
T ₁	0.34	0.35	0.41	0.42	4.65	4.59	4.57	4.54	18.26	18.29	18.31	18.33
T ₂	0.33	0.36	0.40	0.44	4.81	4.71	4.62	4.59	17.67	17.71	17.73	17.76
T ₃	0.37	0.38	0.42	0.46	4.76	4.72	4.66	4.63	16.22	16.26	16.28	16.31
T ₄	0.28	0.31	0.33	0.35	4.87	4.77	4.57	4.53	18.39	18.41	18.43	18.45
T ₅	0.32	0.33	0.38	0.41	5.11	4.85	4.72	4.69	17.02	17.05	17.07	17.10
T ₆	0.31	0.32	0.35	0.38	5.18	5.11	4.84	4.81	18.29	18.31	18.33	18.35
T ₇	0.35	0.36	0.40	0.47	4.79	4.77	4.27	4.24	15.65	15.67	15.70	15.73
T ₈	0.36	0.37	0.41	0.49	5.08	5.07	4.55	4.52	16.37	16.40	16.44	16.48
T ₉	0.35	0.36	0.43	0.51	5.3	4.32	4.11	4.07	16.79	16.84	16.88	16.90
F-test	NS	NS	S	S	S	S	S	S	NS	S	S	S
S.Ed(±)	0.006	0.01	0.18	0.13	0.1	0.11	0.09	0.1	0.28	0.28	0.28	0.27
CD @ 5%	NS	NS	0.43	0.32	0.25	0.27	0.23	0.24	NS	0.68	0.67	0.64

Table 2: Changes in ascorbic acid, colour and appearance and taste in the aonla candy during storage.

Treatments	Ascorbic acid (mg/100g)				Colour and appearance				Taste			
	Storage period (days)				Storage period (days)				Storage period (days)			
	0	4	8	12	0	4	8	12	0	4	8	12
T₀	31.33	31.30	31.27	31.24	4.61	4.52	4.14	4.05	5.61	5.22	5.12	4.91
T₁	31.71	31.69	31.67	31.64	5.48	5.38	5.08	4.42	5.82	5.62	5.22	5.11
T₂	32.50	32.48	32.46	32.44	5.43	5.33	5.13	4.66	5.76	5.56	5.16	5.08
T₃	32.82	32.80	32.78	32.75	5.90	5.80	5.13	5.03	7.08	7.01	6.79	6.66
T₄	33.77	33.75	33.73	33.70	7.78	7.68	7.34	7.11	7.14	7.11	7.01	6.95
T₅	31.71	31.69	31.66	31.63	7.12	7.02	6.69	6.35	5.79	5.45	5.28	5.04
T₆	32.64	32.62	32.60	32.57	6.11	6.09	6.08	6.07	7.11	7.08	6.86	6.73
T₇	32.58	32.57	32.55	32.53	6.22	6.12	6.04	5.97	6.12	6.08	5.95	5.75
T₈	32.66	32.64	32.60	32.57	5.52	5.42	5.42	5.29	5.52	5.39	5.22	5.12
T₉	32.60	32.58	32.56	32.53	9.08	9.01	8.81	8.48	5.43	5.36	5.26	5.19
F-test	S	S	S	S	S	S	S	S	S	S	S	S
S.Ed(±)	0.23	0.24	0.24	0.25	0.30	0.30	0.29	0.37	0.43	0.22	0.14	0.14
CD @ 5%	0.57	0.57	0.58	0.61	0.71	0.71	0.71	0.89	1.02	0.52	0.34	0.33

Table3: Changes in aroma, overall acceptability and shelf life in the guava-papaya during storage.

Treatments	Aroma				Overall acceptability				Shelf life
	Storage period (days)				Storage period (days)				Storage period (days)
	0	4	8	12	0	4	8	12	12
T₀	6.21	6.17	6.00	5.91	6.07	6.01	5.95	5.75	8.21
T₁	5.27	5.23	5.12	5.08	5.08	5.01	4.91	4.78	9.43
T₂	5.19	5.15	5.10	5.03	5.04	4.99	4.99	4.75	8.50
T₃	5.20	5.17	4.99	4.65	5.15	5.08	5.08	4.86	9.50
T₄	7.17	7.15	7.13	7.11	7.03	6.98	6.98	6.77	11.23
T₅	7.10	7.08	7.03	6.90	6.20	6.16	6.07	6.00	10.43
T₆	6.62	6.58	6.49	6.26	6.23	6.19	6.10	6.02	10.90
T₇	5.56	5.51	5.48	5.36	5.06	4.97	4.88	4.82	9.70
T₈	5.16	5.12	4.98	4.81	5.01	4.96	4.87	4.80	8.79
T₉	5.25	5.23	5.13	4.87	4.99	4.93	4.84	4.79	8.20
F-test	S	S	S	S	S	S	S	S	S
S.Ed(±)	0.12	0.12	0.20	0.18	0.20	0.17	0.17	0.13	0.30
CD @ 5%	0.29	0.30	0.48	0.42	0.48	0.41	0.40	0.36	0.71

CONCLUSION

The nectar prepared from the recipe T₄ (60% Guava : 40% Papaya juice) contained most acceptable ascorbic acid with 33.70mg/100g, pH with 4.53, TSS with 18.45 °Brix, acidity 0.35% and also in organoleptic score as compared to other recipes during storage. The study concludes that 60:40 ratio for blending of guava and papaya was most acceptable for preparation of nectar with TSS, acidity, pH, ascorbic acid.

REFERENCES

- Kaur, N., Kumar, A., Monga, P.K. and Arora, P.K. (2011). Biochemical studies in fruits of guava cultivars, *The Asian Journal of Horticulture* **6**(1): 122- 123.
- Kocher, G.S. and Pooja. (2011). Status of wine production from guava (*Psidium guajava* L.): A traditional fruit of India, *African Journal of Food Science* Vol **5**(16), pp. 851-860.
- Othman, O.C. (2009). *Physical and chemical composition of storage-ripen papaya (Carica papaya L.) Fruits of eastern tanzania* Vol **35**.
- Das, S.C. and Dinesh, M.R. (2014). Evaluation of certain papaya varieties and hybrids for physico-chemical characteristics. *The Asian Journal of Horticulture*, **9**(1): 237-239.
- Bal, L.M., Ahmad, T., Senapati, A.K., Pandit, P.S., (2014) Evaluation of Quality Attributes During Storage of Guava Nectar Cv. Lalit from Different Pulp and TSS Ratio. *The International Open Access Journal of Food Processing & Technology* **5**: 329.
- Jumde, A.D., Shukla, R.N. and Gousoddin. (2015). Development and chemical analysis of watermelon blends with beetroot juice during storage. *International journal of Science, Engineering and Technology* **3**(4).
- Kapoor, S., (2016) Antioxidant components and physico-chemical characteristics of jamun powder supplemented pear juice *Journal of Food Science and Technology -Mysore-* **53**(5).
- Rani, T. Baby and Babu, J. Dilip (2015). Acceptability and storage studies of guava - Aloe nectar blends. *The Asian Journal of Horticulture*, **10**(1): 80-85.
- R, S. (2015) Studies on Effect of Processing Quality and Storage Stability of Functional Beverages Prepared from Aloe vera, Blended with Bael Fruit. *International Journal of Food Quality and Safety*, Volume **1**.

Shaheel, SK., Swami, D. V., Prasanna Kumar, B. and Uma Krishna, V., (2015). Effect of blending of karonda (*Carissa carandas* L.) juice with guava, papaya and pineapple juices on its quality and organoleptic evaluation. *Plant Archives* Vol. **15**.

Thirukkumar, S. and Vennila, P., (2019). Processing of blended beverages and its storage stability. *Trends & Prospects in Processing of Horticultural Crops*.

[Kumar Meena](#), N., (2021). Utilization of custard apple pulp for preparation of blended nectar. *Indian Journal of Horticulture* **78**(2).

Babajide, J.M., Olaluwoye, A.A., Taofik Shittu, T.A., Adebisi M.A., (2013). Physicochemical Properties and Phytochemical Components of Spiced Cucumber-Pineapple Fruit Drink. [Nigerian Food Journal](#) Volume **31**.

Malav, M., Gupta, R. and Nagar, T. (2014). Studies on bio-chemical composition of orange based blended ready-to-serve (RTS) beverages. *Bioscience Biotechnology Research Communication* **7**(1).

Yadav, A., Samsher, Singh, J., Kumar, S.C.V. and Goyall, S.K. (2012). Studies on Development and Various Properties of Banana RTS Beverage. *Annals of Horticulture* **5**(2).

Jumde, D.A., Shukla, R.N., Gousoddin. (2015). Development and chemical analysis of watermelon blends with beetroot juice during storage. *International Journal of Science, Engineering and Technology*. Volume **3**.

Rani, T., Baby and Babu, J., Dilip (2015). Acceptability and storage studies of guava - Aloe nectar blends. *The Asian Journal of Horticulture*. **10**(1).

Sindumathi, G., Premalatha, M.R. and Kavitha, V. (2017). Studies on Therapeutic Value of Naturally Flavored Papaya-Mango Blended Ready-To-Serve (RTS) Beverage. *International Journal of Current Microbiology and Applied Sciences*. Volume **6**.

