

Geographical Distribution of Soil Transmitted Helminths and *Plasmodium falciparum* co-infections among school children in Bugesera District, Rwanda

ABSTRACT

Aims: Soil Transmitted Helminths and *Plasmodium* infections are ubiquitous within the tropical and subtropical regions. However, the extent and consequences of Helminthic and *Plasmodium* infections and co-infections geographical distributions are not fully understood. This study aimed determining the geographical distribution of these parasites.

Study design: Cross Sectional Survey

Place and Duration of Study: The survey was conducted in Bugesera District, Eastern Rwanda, between May and December 2020.

Methodology: The survey was conducted among children between ages 5 and 18 years, across 21 randomly selected primary schools. Stool samples were collected and screened for soil transmitted helminths using Kato-Katz, while finger-prick blood samples were examined under the microscope to determine *Plasmodium* infection.

Results: *P. falciparum* was common throughout the study area, with highest prevalence in provinces of Nyiragiseke (22.78%) and Shami (18.99%). The geographical distribution of STH was variably dominated by *A. lumbricoides*. The co-infection exhibited geographical variation consistent with the *A. lumbricoides* prevalence.

Conclusion: The un-proportional distribution of and higher prevalence of *P. falciparum* in the provinces of Nyiragiseke (22.78%) and Shami (18.99%) calls for the review of the control methods, strategies and efforts to address the unique conditions and activities in each province. The variably dominant *A. lumbricoides*-*Plasmodium* co-infection calls for targeted control strategies and efforts for these parasites since this association has been attributed to severe malaria.

Keywords: Geographical Distribution, *Plasmodium falciparum*, Soil transmitted helminths and Co - infections

1. INTRODUCTION

It is established that soil-transmitted helminthiasis (STH) and malaria are ubiquitous and endemic in the tropical regions but exaggerated in Africa – South of the Sahara and South East Asia (1-3). The World Health Organization (WHO) - African Region accounted for about 94 -95% of malaria cases in 2019 - 2020(4, 5). *Plasmodium falciparum* remains the foremost rife *Plasmodium* within the WHO African Region, causing 99.7% of the cases, and 50% within the South East Asia Region. Globally, children are the foremost vulnerable, accounting for 67% of all malaria mortality(6). Forty percent of the global burden of NTDs is in Africa, where these diseases cause high morbidity - especially in school children and loss of person years of their caretakers(7). The African Region has the second highest number of children infested with soil-transmitted helminths of all the WHO regions(3).

Accurate descriptions of the geographical distribution of the said parasitic infections is key to the control methods(8). Nevertheless, it is important to note that climatic and socioeconomic factors influence the distribution of single and multiple co- infections (9-13). STHs and *Plasmodium falciparum* coinfection - ubiquity has a long history, though, with discrepant reports. In addition, these coinfections are not uncommon in school children in the East Africa region as it were for the other endemic regions and are consistent with variable consequences(14-29). The situation is not any different in Rwanda where helminthiasis and malaria remain prevalent in many provinces, in spite of the vigorous and tremendous national control programs(30-33). Recent reports from the studies in Rwanda show that STH and *Plasmodium* infections are heterogeneously distributed across the country with evidence of signatures of spatial clustering with different risk factors including geographical location(33-37). We now report the geographical distribution of STHs and *Plasmodium falciparum* infections in Busegera District of Rwanda.

42 2. MATERIAL AND METHODS

44 2.1 Study setting

45 The study was conducted in Bugesera District, Rwanda. Bugesera District is one of the seven Districts that constitute the Eastern Province in of Rwanda. It is ranging between 30° 05' Eastern longitude, and 2° 09' Southern Latitude, and covers the surface of 1337 km². The District is characterized with a mixture of plateaus with an altitude varying between 1,100 m and 1,780m and undulating hills dominated by varying heights. Bugesera climate is dry with temperature varying between 20^o and 30°C. The district has two dry seasons and two rainy seasons. The hydrographical network is mainly characterized by 3 rivers, namely; Akanyaru, Akagera and Nyabarongo. Besides these rivers, there are 9 lakes, though, with little effect on rainfall, but important for fishing, tourism, transportation, power generation and agricultural irrigation among others.

57 2.2 Survey procedures and sample collection

63 Parents/guardians of the children were invited to attend sensitization meetings. The study procedures were explained in an exceedingly simpler language they felt most comfortable with. Written consent was obtained from all parents/guardians who were willing to have their children participate in the study. Finger prick blood was collected from every child using a capillary tube. Thick and thin blood smears were prepared for the diagnosis of *Plasmodium* parasites. *Plasmodium*-positive slides were re-checked by a senior laboratory technician for quality control and assurance.

2.3 Statistical analysis

The data were entered in EPI INFO 7, and statistical analysis was done using SPSS and EXCEL. Chi-square test including odd ratios at 95% CI and One-way ANOVA was used to test for differences in proportions and means, respectively. Values were considered statistically significant when P -values are <0.05 .

71 3. RESULTS

72 3.1 Geographical distribution of single infection and coinfection


P. falciparum was common throughout the study area, with highest prevalence in provinces of Nyiragiseke (22.78%) and Shami (18.99%), **Fig1**. The geographical distribution of STH was dominated by *A. lumbricoides*, **Fig2**. The STH – *Plasmodium* co-infection exhibited geographical variation and was higher in provinces with high *A. lumbricoides* prevalence, **Fig3**.

79 3.2 Single infections


Overall, the children were variably infected with any soil transmitted helminth species. The most prevalent species was *A. lumbricoides* (4.43%), $P < 0.001$; followed by *T. trichiura* (0.76%) and *Ancylostoma duodenale* (0.16 %). The prevalence of *P. falciparum* was 3.15%. These data are as shown in **Table1**.

84 3.3 Co-infections

85 Overall, the prevalence of STH - *Plasmodium* coinfection was 36.15% and, *A. lumbricoides*- *Plasmodium* co-infection was the only significant co-infection $P < 0.05$, **Table2**.



92 Fig 2. Geographical distribution of helminths



94 Fig 3. Geographical distribution of coinfection

Table 1: Prevalence of Helminth and *Plasmodium* single infections

Characteristic	Overall (n = 2,507)	Boys (n = 1,200), (47.9%)	Girls (n = 1,307), (52.1%)	p-value
Prevalence of helminth infection				
<i>A. lumbricoides</i> (% , 95 % CI)	4.43 (12.28-29.65)	4.50 (9.89-35.26)	4.36 (7.17-31.71)	<0.001
<i>T. trichiura</i> (% , 95 % CI)	0.76 (2.62-7.27)	0.58 (0.41-8.16)	0.92 (1.97-8.70)	0.05
<i>A. duodenale</i> (% , 95 % CI)	0.16 (0.70-3.30)	0.17 (0.50-8.85)	0.15 (0.61-7.85)	0.04
Prevalence of <i>Plasmodium</i> infection				
<i>P. falciparum</i> (% , 95 % CI)	3.15 (144.57-828.39)	3.33 (156.72-971.08)	2.98 (194.96-414.07)	

Table 2: Overall Prevalence of Co-infections

Characteristic	Overall (n = 2,507)	Boys (n = 1,200), (47.9%)	Girls (n = 1,307), (52.1%)	p-value
Coinfection				
<i>T. trichiura</i> - <i>A. lumbricoides</i> (% , 95 % CI)	3.08 (4.34-17.16)	0	5.97 (4.34-17.16)	0.432
<i>A. lumbricoides</i> - <i>P. falciparum</i> (% , 95 % CI)	36.15 (9.90-11.89)	30.16 (8.08-11.08)	41.79 (10.50-13.07)	<0.05

4. DISCUSSION

Reports from the studies in Rwanda and elsewhere show that STH and *Plasmodium* infections are heterogeneously distributed across the country with evidence of signatures of spatial clustering with different risk factors including geographical location(19, 33-37). Our finding of the distribution of *P. falciparum* is actually in phase with the previous reports. However, the prevalence in provinces of Nyiragiseke (22.78%) and Shami (18.99%) is far much higher than the overall prevalence of the district in the same study population(33). The finding calls for the review of the control methods, strategies and efforts to address the unique conditions and activities in each province.

The geographical distribution of single STH was variably dominated by *A. lumbricoides*, and the STH – *Plasmodium* co-infection followed the same pattern since there was only one significant *A. lumbricoides*- *Plasmodium* co-infection, $P < 0.05$. This association has been consistently reported from the studies in the same district in Rwanda and elsewhere(14, 30, 33, 38, 39). Most importantly and interestingly, this particular co-infection has been attributed to severe malaria(22, 30, 40). There is, therefore, need for targeted control strategies and efforts for these parasites.

5. CONCLUSION

152 The un-proportional distribution of and higher prevalence of *P. falciparum* in the provinces of Nyiragiseke (22.78%) and Shami (18.99%) calls for the review of the control methods, strategies and efforts to address the unique conditions and activities in each province. The variably dominant *A. lumbricoides*- *Plasmodium* co-infection calls for targeted control strategies and efforts for these parasites since this association has been attributed to severe malaria.

CONSENT

All participants and parents/guardians gave consent to participate.

154 ETHICAL APPROVAL

155 The approval was provided by the University of Rwanda IRB (No. 380/CMHS).

156

REFERENCES

1. Hay SI, Omumbo JA, Craig MH, Snow RW. Earth observation, geographic information systems and *Plasmodium falciparum* malaria in sub-Saharan Africa. *Adv Parasitol.* 2000;47:173-215.
2. Brooker S, Michael E. The potential of geographical information systems and remote sensing in the epidemiology and control of human helminth infections. *Adv Parasitol.* 2000;47:245-88.
3. <Fourth WHO Report on NTDs 2017.pdf>.
4. <World Malaria Report 2020.pdf>.
5. <Malaria Report 2021.pdf>.
6. <WHO Malaria Report 2019.pdf>.
7. <WHO AFRO CDS Annual Report 2016 web version_1.pdf>.
8. Brooker S, Clements AC, Bundy DA. Global epidemiology, ecology and control of soil-transmitted helminth infections. *Adv Parasitol.* 2006;62:221-61.
9. Bødker R, Akida J, Shayo D, Kisinza W, Msangeni HA, Pedersen EM, et al. Relationship between altitude and intensity of malaria transmission in the Usambara Mountains, Tanzania. *J Med Entomol.* 2003;40(5):706-17.
10. Baragatti M, Fournet F, Henry MC, Assi S, Ouedraogo H, Rogier C, et al. Social and environmental malaria risk factors in urban areas of Ouagadougou, Burkina Faso. *Malar J.* 2009;8:13.
11. Righetti AA, Glinz D, Adiossan LG, Koua AY, Niamké S, Hurrell RF, et al. Interactions and potential implications of *Plasmodium falciparum*-hookworm coinfection in different age groups in south-central Côte d'Ivoire. *PLoS Negl Trop Dis.* 2012;6(11):e1889.
12. Yapi RB, Hurlimann E, HOUNGBEDJI CA, NDRI PB, SILUE KD, SORO G, et al. Infection and co-infection with helminths and *Plasmodium* among school children in Cote d'Ivoire: results from a National Cross-Sectional Survey. *PLoS Negl Trop Dis.* 2014;8(6):e2913.
13. Omondi CJ, Onguru D, Kamau L, Nanyingi M, Ong'amo G, Estambale B. Perennial transmission of malaria in the low altitude areas of Baringo County, Kenya. *Malar J.* 2017;16(1):257.
14. Le Hesran JY, Akiana J, Ndiaye el HM, Dia M, Senghor P, Konate L. Severe malaria attack is associated with high prevalence of *Ascaris lumbricoides* infection among children in rural Senegal. *Trans R Soc Trop Med Hyg.* 2004;98(7):397-9.
15. Brooker S, Clements AC, Hotez PJ, Hay SI, Tatem AJ, Bundy DA, et al. The co-distribution of *Plasmodium falciparum* and hookworm among African schoolchildren. *Malar J.* 2006;5:99.
16. Brooker S, Akhwale W, Pullan R, Estambale B, Clarke SE, Snow RW, et al. Epidemiology of plasmodium-helminth co-infection in Africa: populations at risk, potential impact on anemia, and prospects for combining control. *Am J Trop Med Hyg.* 2007;77(6 Suppl):88-98.
17. Degarege A, Animut A, Legesse M, Erko B. Malaria severity status in patients with soil-transmitted helminth infections. *Acta Trop.* 2009;112(1):8-11.
18. Mazigo HD, Kidenya BR, Ambrose EE, Zinga M, Waihenya R. Association of intestinal helminths and *P. falciparum* infections in co-infected school children in northwest Tanzania. *Tanzan J Health Res.* 2010;12(4):299-301.
19. Pullan RL, Kabatereine NB, Bukirwa H, Staedke SG, Brooker S. Heterogeneities and consequences of *Plasmodium* species and hookworm coinfection: a population based study in Uganda. *J Infect Dis.* 2011;203(3):406-17.
20. Brooker SJ, Pullan RL, Gitonga CW, Ashton RA, Kolaczinski JH, Kabatereine NB, et al. *Plasmodium*-helminth coinfection and its sources of heterogeneity across East Africa. *J Infect Dis.* 2012;205(5):841-52.
21. Degarege A, Legesse M, Medhin G, Animut A, Erko B. Malaria and related outcomes in patients with intestinal helminths: a cross-sectional study. *BMC Infect Dis.* 2012;12:291.
22. Mulu A, Legesse M, Erko B, Belyhun Y, Nugussie D, Shimelis T, et al. Epidemiological and clinical correlates of malaria-helminth co-infections in Southern Ethiopia. *Malar J.* 2013;12:227.
23. Legason ID, Atiku A, Ssenyonga R, Olupot-Olupot P, Barugahare JB. Prevalence of Anaemia and Associated Risk Factors among Children in North-western Uganda: A Cross Sectional Study. *BMC Hematol.* 2017;17:10.
24. Kepha S, Nuwaha F, Nikolay B, Gichuki P, Edwards T, Allen E, et al. Epidemiology of coinfection with soil transmitted helminths and *Plasmodium falciparum* among school children in Bumula District in western Kenya. *Parasit Vectors.* 2015;8:314.
25. Adu-Gyasi D, Asante KP, Frempong MT, Gyasi DK, Iddrisu LF, Ankrah L, et al. Epidemiology of soil transmitted Helminth infections in the middle-belt of Ghana, Africa. *Parasite Epidemiol Control.* 2018;3(3):e00071.
26. Babamale OA, Ugbomoiko US, Heukelbach J. High prevalence of *Plasmodium falciparum* and soil-transmitted helminth co-infections in a periurban community in Kwara State, Nigeria. *J Infect Public Health.* 2018;11(1):48-53.
27. Kabatende J, Mugisha M, Ntirenganya L, Barry A, Ruberanziza E, Mbonigaba JB, et al. Prevalence, Intensity, and Correlates of Soil-Transmitted Helminth Infections among School Children after a Decade of Preventive Chemotherapy in Western Rwanda. *Pathogens.* 2020;9(12).
28. Oboth P, Gavamukulya Y, Barugahare BJ. Prevalence and clinical outcomes of *Plasmodium falciparum* and intestinal parasitic infections among children in Kiryandongo refugee camp, mid-Western Uganda: a cross sectional study. *BMC Infect Dis.* 2019;19(1):295.
29. Easton AV, Raciny-Aleman M, Liu V, Ruan E, Marier C, Heguy A, et al. Immune Response and Microbiota Profiles during Coinfection with *Plasmodium vivax* and Soil-Transmitted Helminths. *mBio.* 2020;11(5).

30. Marcelline U, Noella U, Tharcisse M, Corine K, Josephat M, Banson B. The Impact of Malaria and Gastrointestinal Helminthiasis Co-infection on Anaemia and Severe Malaria among Children in Bugesera District, Rwanda. *International Journal of TROPICAL DISEASE & Health*. 2016;13(4):1-7.
 31. Rujeni N, Morona D, Ruberanziza E, Mazigo HD. Schistosomiasis and soil-transmitted helminthiasis in Rwanda: an update on their epidemiology and control. *Infect Dis Poverty*. 2017;6(1):8.
 32. Geus D, Sift KC, Habarugira F, Mugisha JC, Mukampunga C, Ndoli J, et al. Co-infections with *Plasmodium*, *Ascaris* and *Giardia* among Rwandan schoolchildren. *Trop Med Int Health*. 2019;24(4):409-20.
 33. John Banson B, Sharma A, Joseph M, David T, D. Humphrey M, Marcelline U. Soil Transmitted Helminths and *Plasmodium falciparum* Co-infections among School Children in Bugesera District, Rwanda: Implications for National Control Programs. *International Journal of TROPICAL DISEASE & Health*. 2021:30-6.
 34. Ruberanziza E, Owada K, Clark NJ, Umulisa I, Ortu G, Lancaster W, et al. Mapping Soil-Transmitted Helminth Parasite Infection in Rwanda: Estimating Endemicity and Identifying At-Risk Populations. *Trop Med Infect Dis*. 2019;4(2).
 35. Rudasingwa G, Cho SI. Determinants of the persistence of malaria in Rwanda. *Malar J*. 2020;19(1):36.
 36. Habyarimana F, Ramroop S. Prevalence and Risk Factors Associated with Malaria among Children Aged Six Months to 14 Years Old in Rwanda: Evidence from 2017 Rwanda Malaria Indicator Survey. *Int J Environ Res Public Health*. 2020;17(21).
 37. Nzabakiriraho JD, Gayawan E. Geostatistical modeling of malaria prevalence among under-five children in Rwanda. *BMC Public Health*. 2021;21(1):369.
 38. Brutus L, Watier L, Briand V, Hanitrasoamampionona V, Razanatsoarilala H, Cot M. Parasitic co-infections: does *Ascaris lumbricoides* protect against *Plasmodium falciparum* infection? *Am J Trop Med Hyg*. 2006;75(2):194-8.
 39. Brutus L, Watier L, Hanitrasoamampionona V, Razanatsoarilala H, Cot M. Confirmation of the protective effect of *Ascaris lumbricoides* on *Plasmodium falciparum* infection: results of a randomized trial in Madagascar. *Am J Trop Med Hyg*. 2007;77(6):1091-5.
 40. Degarege A, Anmut A, Legesse M, Erko B. Malaria and helminth co-infections in outpatients of Alaba Kulito Health Center, southern Ethiopia: a cross sectional study. *BMC Res Notes*. 2010;3:143.
-