

Original Research Article

Assessment of Bacterial contamination in drinking water sources in Khartoum/ Sudan

ABSTRACT

Aim of the Study: To assess the bacterial contamination in drinking water sources in Khartoum/ Sudan

Place of Study: Central Veterinary Research Laboratory/ Bacteriology Department.

Study Design: One hundred water samples were collected from the three localities of Khartoum state (Khartoum= 33, Omdurman= 34, Khartoum north [Bahri] =25) and 8 from different companies of water supply.

Methodology: Fifty four Samples were collected from surface water and (38) from ground water [well]. These samples transported to bacteriology lab for microbiological analyses using filtration method and new technique Colilert and Pseudalert kits which used for the first time in Sudan.

Results: Filtration method revealed different bacterial species, they were: *Escherichia coli*, *Klebsiella pneumonia*, *Enterobacter aerogene* *Enterobacter sakazaki*, *Enterobacter cloacae*, *Serratia marinoruba*, *Proteus mirabilis*, *Salmonella* spp. *Raoultella terrigena* *planticola*, *Orchobacter anthrobi*, *Cronobater* spp., *Aeromonas salmonicida*, *Aeromonas hydrophilia*, *Pantoea agglomerans*, *Vibrio parahaemolyticus*. Coliform bacteria, *Escherichia coli* and *Pseudomonas* app were detected and most probable numbers (MPN) were counted using the previous kits according to manufacture instructions.

Conclusion: The water must be tested before using and quality control technique must be achievable to ensure continuously supply of pure drinking water.

Key words: *Drinking water, E. coli, Pseudomonas spp., Coliform, Colilert, Pseudalert, MPN*

1. INTRODUCTION

Water is essential to life. An adequate, safe and accessible supply must be available to all. Improving access to safe drinking-water can result in significant benefits to health. Every effort should be made to achieve a drinking water quality as safe as possible. In general terms, the greatest microbial risks are associated with ingestion of water that is contaminated with human or animal feces. Wastewater discharges in fresh waters and costal seawaters are the major source of fecal microorganisms, including pathogens [1-4].

Bacteria of greatest concern in drinking water are those that originate from the gut of warm-blooded animals. Sources include wildlife, pets, and contamination problems arise from improperly designed, failing, or overloaded waste water treatment systems, including septic systems from private homes, and leaking sanitary sewer pipes. Human sources are a particular concern as they include bacteria of human origin and may include human pathogens. Floodwater commonly contains high levels of bacteria from numerous sources.

Surface waters typically contain bacteria. Groundwater should be free of those bacteria that arise from animal sources. Proper well design and construction should protect drinking water sources from contamination by surface waters, livestock on farms or in feedlots.

A total coliform bacterium is a group of different kinds of bacteria commonly found in the environment, including soil, vegetation and untreated surface water. Total coliform bacteria are generally not harmful. Fecal coliform bacteria are a subgroup of the total coliform group.

The most important species of the group include *Escherichia coli*, *Klebsiella spp* and *Enterobacter spp*. At the same time, non-coliform bacteria were also found in polluted water such as *Sterptococcus*, *Proteus* and *Pseudomonas* species [5-7].

In this study The Colilert test was used which is an advance test because it uses proprietary Defined Substrate Technology (DST) to simultaneously detect total coliforms and *E. coli* two nutrient-indicators, ONPG and MUG, are the major sources of carbon in Colilert and can be metabolized by the coliform enzyme β -galactosidase and the *E. coli* enzyme β -glucuronidase, respectively. The Pseudalert test was used to detect *Pseudomonas aeruginosa* in water sample. The test is based on a bacterial enzyme detection technology that signals the presence of *Pseudomonas aeruginosa* through the hydrolysis of substrate in the Pseudalert reagent. *Pseudomonas* cells rapidly grow and reproduce using the rich supply of amino acids, vitamins and other nutrients present in the Pseudalert reagent [8].

This approach is different from traditional media, which provide a nutrient-rich environment that supports the growth of both target organisms and non-targets. When non targets grow and mimic target organisms, false positives occur. Growth of non-targets can also suppress target organisms and give false negatives in traditional media. To suppress non targets, traditional media often include high levels of salts, detergents, or other selective agents that may inadvertently suppress target organisms and give further false negatives.

2. MATERIALS AND METHODS

This study was designed to evaluate drinking water hygiene, with special reference to the incidence of bacterial water-borne disease in some parts of Sudan using new technique (Colilert and Pseudalert test) for the first time in Sudan.

The Colilert test simultaneously detects or quantifies both total coliforms and *E coli*, with results in 24 hours. As coliforms grow in the Colilert Test, they use β -galactosidase to metabolize ONPG and change it from colorless to yellow. *E. coli* use β -glucuronidase to metabolize MUG and create fluorescence. Since most non-coliforms do not have these enzymes, they are unable to grow and interfere. The few non-coliforms that do have these enzymes are selectively suppressed

by the Colilert Test's specifically formulated matrix. Pseudalert test also detects and quantifies *Pseudomonas aeruginosa*, actively growing strains have an enzyme that cleave substrate in the reagent to produce blue fluorescence under ultraviolet light.

2.1 Samples collection:

One hundred water samples from three localities in Khartoum state (Khartoum, Khartoum north (Bahri), Omdurman) were collected aseptically in sterile plastic containers then labeled and placed in an ice-boxes and transported immediately to the Central Veterinary Research Laboratory, department of Bacteriology for microbiological analysis. Each sample consist of 300 ml distributed in 3 container (each one 100 ml). Thirty six samples were taken from well (ground water), 56 samples directly from pipe (surface water) and 8 purified drinking water samples from different companies. These samples were classified as followed:

The surface water samples were: 11 samples from Khartoum, 19 from Omdurman and 26 from Khartoum north (Bahri).

The ground water samples were: 14 samples from Khartoum, 15 from Omdurman and 7 samples from Khartoum north (Bahri).

The purified samples were: 8 samples from different companies

2.2 Methodology:

2.2.1 Filtration Method:

One hundred ml of each sample was filtered through filter papers (4.5 µm in diameter) using filtration system, then the membrane was put on sterile Petri dishes containing absorbable paper impregnated with 2.5 ml sterile Endo broth media then incubated for 24h at 37°C. Well isolated colonies from filter paper as shown in (fig 1) were sub cultured on nutrient agar plates for purification then identified using conventional methods and API kits.

2.2.2 Analysis of water sample using Pseudalert and Colilert (Kits):

One hundred ml of water was used for Pseudalert and another 100 ml for Colilert.

The reagent was added to sample then poured into Quanti-Tray (counts from 1–200) or Quanti-Tray/2000 (counts from 1–2,419), Sealed and incubated at 37°C for 24 hours. After that results were recorded according to manufacturer instructors (IDEXX) and the Most Probable Number (MPN) was calculated as followed: Yellow wells = total coliforms, Yellow/fluorescent wells = *E. coli*, then positive wells were Count and refer to MPN table.

Under ultraviolet light, Blue fluorescence indicates the presence of *Pseudomonas aeruginosa*.

3. RESULTS

3.1 Out of 56 surface water samples were revealed the following bacterial species: *Escherichia coli*, *Pseudomonas* spp, *Aeromonas salmonicida*, *Aeromonas hydrophila caviae/sobria*, *Enterobacter sakazaki*, *Enterobacter aerogene*, *Enterobacter cloacae*, *Proteus mirabilis* and

other *proteus* spp, *Klebsiella pneumoniae*, *Pantoea agglomerans*, *pantoea* spp. *Raoultella terrigena*, *Staphylococcus lentus*, *Vibrio paraheamolyticus*, *Salmonella clori-arizona* and other *Salmonella* spp. Only 15 samples showed no bacterial growth.

3.2 Out of 36 ground water samples, 30 samples revealed the following bacterial species: *Escherichia coli*, *Pseudomonas* spp, *Enterobacter sakazaki*, *Orchobacter anthropi*, *Enterobacter cloaca*, *Cronobacter* spp., *Proteus* spp, *Klebsiella pneumoniae*, and other *Klebsiella* spp., *Aeromonas salmonicida*, *Raoultella terrigena/planticola*, *Serratia marinoruba*, *Orchobacter anthropi*, *Enterobacter sakazaki* and *Enterobacter cloaca*. Only 6 samples showed no bacterial growth.

3.3 Bacteria detected from surface and ground water using IDEXX were coliforms, *E. coli* and *Pseudomonas* spp. The most probable number (MPN) range from 3 to > 2419.6. The results obtained as followed:

Table 1: Bacterial contamination from surface water

Area	No of samples	Coliform	<i>Escherichia coli</i>	<i>Pseudomonas</i> spp.
Khartoum	11	1 -ve 10 +ve	4 -ve 7 +ve	1 -ve 10 +ve
Omdurman	19	4 -ve 15 +ve	6 -ve 13 +ve	8 -ve 11 +ve
Khartoum north (Bahri)	26	10 -ve 16 +ve	14 -ve 12 +ve	13 -ve 13 +ve
Total	56	15 -ve 41 +ve	24 -ve 32 +ve	22 -ve 34 +ve

Table 2: Bacterial contamination from ground water

Area	No of samples	Coliform	<i>Escherichia coli</i>	<i>Pseudomonas</i> spp.
Khartoum	14	5 -ve 9 +ve	10 -ve 4+ ve	10 - ve 4 +ve
Omdurman	15	0 -ve 15 +ve	7 - ve 8 + ve	9 - ve 6 +ve

Khartoum north (Bahri)	7	1 -ve 6 +ve	6 -ve 1 +ve	3 -ve 4 +ve
Total	36	6 -ve 30 +ve	23 -ve 13 +ve	22 -ve 14 +ve

Table 3: Number of positive and negative sample

Source of sample	Total number	Positive	Negative
Surface water	56	41(73.2%)	15 (26.8%)
Ground water	36	30 (83.3%)	6 (16.7%)
Purified water	8	0 (0%)	8 (100%)
Total	100	71%	29%




Fig 1: shows bacterial growth using filtration method



Fig 2: Negative and positive results of bacterial growth after exposing colilert and Pseudalert test to UV

3.4 Purified water samples were obtained from 8 companies named as follow: (Safia, Anhar, Crystal, Forat, Yes, Miso, Zulal and Care), all these samples were negative for bacterial growth when tested by filtration method and Colilert, Pseudalert test.

4. DISCUSSION

The need to increase attention on the safety of drinking water, rather than just the quantity has been acknowledged globally through the commitment to the Sustainable Development Goals (SDGs). SDG 6.1 focuses on achieving universal and equitable access to safe and affordable drinking water for all, and SDG 6.2 focuses on improving water quality by reducing pollution. The need for increased attention on drinking water safety (DWS) in Sudan has also been clearly illustrated by the AWD outbreak of 2016/17, which highlighted gaps and challenges being faced in ensuring DWS [9].

Total coliform bacteria are a group of different kinds of bacteria commonly found in the environment, including soil, vegetation and untreated surface water. Fecal coliform bacteria are a subgroup of the total coliform group. In this study, the percentage of coliform in surface (73.2%) (as shown in table 1) and ground water (83.3%) (as shown in table 2), were very high, these could be to the different types of bacteria that existed in great quantities in the intestines and feces of humans and animals. The presence of fecal coliform bacteria in drinking water is a strong indication of recent sewage or animal waste contamination, which should be interpreted as

an indication that there is a greater risk that pathogens are present (table 3). Microbes in these wastes may cause short-term effects, such as diarrhea, cramps, nausea, headaches or other symptoms, as well as potentially pose long-term health effects. They may pose a special health risk for animal, infants, young children, some of the elderly and people with severely compromised immune systems. Septic tank widely used for wastewater storage and treatment may contaminate groundwater supplies.

Many farmers use cellars, tanks or landfills to store manure, water leaching from these storage sites may also contaminate groundwater, especially during periods of rainfall. The application of animal manure to agricultural lands as fertilizer is common practice throughout the world. The bacteria present in the manure may leach into the groundwater by João [10].

An important source of contamination of surface and ground waters is runoff water from agricultural and pasture lands, and urban areas. In a study reported by Doran and Linn [108], runoff from a cow-calf pasture in eastern Nebraska was monitored during a three-year period. Rainfall runoff from the grazed area contained 5 to 10 times more fecal coliforms than runoff from the fenced; un grazed area [11].

This study had verified the use of IDEXX Colilert (18-hour & 24-hour) as an acceptable alternative to other test methods for the recovery of *E. coli* from drinking water, source water, and wastewater [12-18], the test has many good features, such as ease of use, simplifies training, Unit-dosed packaging eliminates media preparation, no repeat testing due to clogged filters or heterotrophic interference. Also it is rapid in detecting coliforms and *E. coli* simultaneously in 24 hours or less, no confirmations needed, no glassware cleaning or colony counting. It is so accurate, it can identify *E. coli* specifically, eliminate unnecessary public notification due to non-target organisms, suppresses up to 2 million heterotrophs per 100 ml. Eliminates the subjective interpretation found in traditional methods. It can also detect a single viable coliform or *E. coli* per sample. It is economical and flexible.

Other type of bacteria that not belong to coliform were also isolated during this study, this is a serious issue since they could harbor pathogenic and resistance genes difficult to treat. Neither *Pseudomonas* spp nor *Aeromonas* spp are indices of fecal pollution, but may be useful in assessing regrowth in distribution systems [19].

Vibrio parahaemolyticus was reported in this study from surface water, as mention by João, it is a well-documented causal agent of acute food-borne gastroenteritis, particularly in Japan and South East Asia. Cases are associated with the consumption of raw or undercooked shellfish such as oysters, shrimp, crabs, and lobster [10].

Also *Salmonella* spp were isolated from surface water, this is in agreement with Arvanitidou and Le who mentioned that, the principal habitat of *Salmonella* is the intestinal tract of humans and animals. Salmonellae are constantly found in environmental samples, because they are excreted by humans, pets, farm animals, and wild life. Municipal sewage, agriculture pollution, and storm water runoff are the main sources of these pathogens in natural waters. Salmonellae do not seem to multiply significantly in the natural environment, but they can survive several weeks in water and in soil if conditions of temperature, humidity, and pH are favorable [20,21].

E. coli was isolated in this study (30 from surface water and 15 from ground water), its strains isolated from intestinal diseases have been grouped into at least six different main groups, based on epidemiological evidence, phenotypic traits, clinical features of the disease and specific virulence factors. From these, enterotoxigenic (ETEC, namely O148), enterohemorrhagic (EHEC, namely O157) and enteroinvasive serotypes (EIEC, namely O124) are of outstanding importance and can be transmitted through contaminated water [22,23].

In recent years, *Aeromonas hydrophila* has gained public health recognition as an opportunistic pathogen. It has been implicated as a potential agent of gastroenteritis, septicemia, meningitis, and wound infections. It can play a significant role in intestinal disorders in children under five years old, the elderly, and immunosuppressed people [24].

There were no bacteria reported in purified water bottles, this may be because of the continuous disinfection of the water supply using many options like chlorination, ultraviolet radiation, distillation, and ozone treatment.

Enterobacter, *Klebsiella*, *Proteus*, *Serratia*, *Salmonella*, *Pseudomonas* were isolated in this study, this is in agreement with Ayman, (2006) from Eastern and Southern Sudan [25].

5. CONCLUSION

The water hygiene situation is poor in Khartoum state due to contamination with many bacterial contaminants which constitute hazard to be spread to both man and animal.

6. RECOMENDATIONS

6.1 Water system must conduct an assessment to find out how the contamination got into the water. If the assessment identifies the cause of the contamination, the water system can usually correct the problem with repairs, treatment, or improved operation and maintenance practices.

If total coliform and/or *E. coli* bacteria are detected in a private water supply, an immediate effort should be made to identify and eliminate the source of contamination. After addressing the contamination source, disinfect the entire water system, using shock chlorination. After shock chlorination, submit another water sample for testing. The water should test negative before use.

If the source of bacterial contamination or well construction errors cannot be identified and eliminated, continuous disinfection of the water supply may be necessary. Options include continuous chlorination, ultraviolet radiation, distillation, and ozone treatment.

6.2 Establishing a well-planned water pollution researches in all these localities to check the microbial quality constantly.

6.3 Further studies needed to evaluate the water microbial situation.

6.4 Safe drinking water for all is one of the major challenges of the 21st century.

6.5 Microbiological control of drinking water should be the norm everywhere.

REFERENCES

1. WHO (World Health Organization). Guidelines for Drinking-water Quality, Incorporating 1st and 2nd Addenda, Volume 1, Recommendations, 3rd ed.; WHO: Geneva, Switzerland, 2008.
2. Fenwick, A. Waterborne Diseases—Could they be Consigned to History? *Science* 2006, 313, 1077–1081.
3. George, I.; Crop, P.; Servais, P. Use of β -D-Galactosidase and β -D-Glucuronidase Activities for Quantitative Detection of Total and Faecal Coliforms in Wastewater. *Can. J. Microbiol.* 2001, 47, 670–675.
4. Grabow, W.O.K. Waterborne Diseases: Update on Water Quality Assessment and Control. *Water SA* 1996, 22, 193–202.
- 5- Alcano, E.I (1997). *Fundamental of Microbiology*. 4th edition. The Benjamin/ cumming publishing company Inc, Red wood city.
- 6- Jeffrey T.; Thomas E.; and Hancock, D.D. (2001). Cattle water troughs as Reservoirs of *Escherichia coli* 0157. *Appl. Envir. Microbiol* 67:3053-3057.
- 7- Murinda, S. E.; Nguyen, L.T.; Nam, H. M.; Almeida, R. A.; Headrick, S. J. and Oliver, S. P. (2004). Detection of sorbitol-negative and sorbitol-positive shiga toxin-producing *Escherichia coli*, *Listeria monocytogenes*, *Campylobacter jejuni*, and *Salmonella* spp. in dairy farm environmental samples. *Foodborne Pathog Dis.* 1(2):97-104.
- 8- Erick Burren. Surface Water Ambient Monitoring Program Using IDEXX For Fecal Indicator Bacteria Monitoring Updated August 2009. Citizen Monitoring Coordinator SWRCB-Clean Water Team. (213) 576-6788,
- 9- Ministry of Water Resources, Irrigation and Electricity Federal Ministry of Health Sudan Drinking, 2017. Water Safety Strategic Framework (SDWSSF)
- 10- João P. S. Cabral. (2010). Water Microbiology. Bacterial Pathogens and Water, *Int. J. Environ. Res. Public Health*, 7: 3657-3703
- 11- EPA. Source Water Protection Practices Bulletin. Managing Stormwater Runoff to Prevent Contamination of Drinking Water; Office of Water (4606); United States Environmental Protection Agency (EPA): Washington, DC, USA, 2009.

- 12- Edberg, S., M. Allen, D. Smith and N. Kriz. 1990. Enumeration of total coliforms and *Escherichia coli* from source water by the defined substrate technology. *Appl. Environ. Microbiol.* 56 (2):366-369.
- 13- Rice, E., J.A. Martin and S. Edberg. 1990. Efficacy of β - Glucuronidase assay for the identification of *Escherichia coli* by the defined-substrate technology. *Appl. Environ. Microbiol.* 56:1203-1205.
- 14- Edberg, S.C., M.J. Allen and D.B. Smith. 1991. Defined Substrate Technology Method for Rapid and Specific Simultaneous Enumeration of Total Coliforms and *Escherichia coli* from Water: Collaborative Study. *J. Assoc. Official Anal. Chem.* 74(3):526-529.
- 15- Fricker, C.R., E.J. Fricker, T. Goodall and J. Cowburn. 1995. Quantitative Procedures for the Detection of *E. coli*, Coliforms, and Enterococci in Water Using Quanti-Tray and Enterolert. Presented at: Water Quality Technology Conference (WQTC), 1995, New Orleans, LA. Graham, C. 1999. Comparison of Coliform/*Escherichia coli* Count Methods Trial. UK Public Health Laboratory System, Water & Environmental Research Unit (Nottingham)
- 16- Barrell, R.A.E., S.R. Cole, R. Down, C. Fricker, P. Holmes, C. Jones, N. Lightfoot, D. Sartory, S. Scanlon, J.A. Taylor, H.E. Tillet and J. Watkins. 1997. Evaluation Trials for Two Media for the Simultaneous Detection and Enumeration of *Escherichia coli* and coliform organisms. In: *Methods for the Examination of Waters and Associated Materials*. UK Environment Agency.
- 17- Eckner, K. 1998. Comparison of membrane filtration and multipltube fermentation by the colilert and enterolert methods for detection of waterborne coliform bacteria, *Escherichia coli*, and enterococci used in drinking and bathing water quality monitoring in southern Sweden. *Appl. Environ. Microbiol.* 64(8):3079-3083.
- 18- Graham, C. 1999. Comparison of Coliform/*Escherichia coli* Count Methods Trial. UK Public Health Laboratory System, Water & Environmental Research Unit (Nottingham).
- 19- WHO and UNICEF (2000) Global Water Supply and Sanitation Assessment 2000 Report. WHO/UNICEF, USA.
- 20- Arvanitidou, M.; Kanellou, K.; Vagiona, D.G. Diversity of *Salmonella* spp. and Fungi in Northern Greek Rivers and their Correlation to Faecal Pollution Indicators. *Environ. Res.* 2005, 99, 278–284.

- 21- Le Minor. The genus *Salmonella*. In *The Prokaryotes: An Evolving Electronic Resource for the Microbiological Community*, electronic release 3.14, 3th ed.; Dworkin, M., Falkow, S., Rosenberg, E., Eds.; Springer-Verlag: New York, NY, USA, 2003.
22. Bettelheim, K.A. The genus *Escherichia*. In *The Prokaryotes: An Evolving Electronic Resource for the Microbiological Community*, electronic release 3.14, 3th ed.; Dworkin, M., Falkow, S., Rosenberg, E., Eds.; Springer-Verlag: New York, NY, USA, 2003.
23. Scheutz, F.; Strockbine, N.A. Genus *Escherichia*. In *Bergey's Manual of Systematic Bacteriology*, 2nd ed.; Brenner, D.J., Krieg, N.R., Staley, J.T., Eds.; Springer: New York, NY, USA, 2005; Volume 2, Part B, pp. 607–623.
- 24- Daskalov, H. The Importance of *Aeromonas hydrophila* in Food Safety. *Food Control* 2006, 17, 474–483.
- 25- Ayman Ibrahim Al-Beeli. Survey of Bacterial Load of Drinking Water In Eastern and Southern Sudan, University of Khartoum; 2006