

Tempered distribution version of the Tumarkin result

Original Research Article

ABSTRACT

We give a tempered distribution analogue of the Tumarkin result that concerns approximation of some functions by sequence of rational functions with given poles.

Keywords: Boundary values of distributions, Distributions, Tumarkin

1. INTRODUCTION

For the needs of our subsequent work we will define the Blaschke product in the upper half plane Π^+ , $\Pi^+ = \{z \in \mathbb{C} | \text{Im}z > 0\}$. Assume

$$\sum_{n=1}^{\infty} \frac{y_n}{1 + |z_n|^2} < \infty, z_n = x_n + iy_n \in \Pi^+ \quad (1.1)$$

Then the Blaschke product with zeros z_n is

$$B(z) = \left(\frac{z-i}{z+i}\right)^m \prod_{n=1}^{\infty} \frac{(|z_n^2+1|)}{z_n^2+1} \frac{z-z_n}{z-\bar{z}_n}, z \in \Pi^+ \quad (1.2),$$

Let

$$z_{k_1}, z_{k_2}, \dots, z_{k_{N_k}}, k = 1, 2, \dots, \text{Im}z \neq 1, N_k \leq \infty \quad (1.3)$$

be given complex numbers. Some of the numbers in (1.3) might be equal and also some of them might be equal to ∞ (in that case $\text{Im}z = 0$)

Let R_k be the rational function of the form

$$R_k(z) = \frac{c_0 z^p + c_1 z^{p-1} + \dots + c_p}{(z - z_{k_1})(z - z_{k_1}) \dots (z - z_{k_{n_p}})}, z \in \Pi^+, \quad (1.4)$$

whose poles are some of the number in (1.3) and c_0, c_1, \dots, c_p are arbitrary numbers (if some $z_{k_i} = \infty$, then in (1.4) we put 1 instead $z - z_{k_i}$).

All z_{k_i} , for which $\text{Im}z_{k_i} > 0$, will be denoted by z'_{k_i} , and all those z_{k_i} , for which $\text{Im}z_{k_i} < 0$ will be denoted by z''_{k_i} .

Let

$$S'_k = \sum_i \frac{\text{Im}z_{k_i}}{1 + |z'_{k_i}|^2} \quad \text{and} \quad S''_k = \sum_i \frac{(-\text{Im}z''_{k_i})}{1 + |z''_{k_i}|^2}.$$

With (1.5) we denote the following conditions

$$\overline{\lim}_{k \rightarrow \infty} S'_k < \infty, \quad \overline{\lim}_{k \rightarrow \infty} S''_k = \infty. \quad (1.5)$$

Let B_k be the Blaschke product whose zeros are the numbers, $z_{k_1}, z_{k_2}, \dots, z_{k_{N_k}}$, from the numbers (1.3), $k = 1, 2, 3, \dots$. Assume (1.5). Then $\mu(z) = \lim_{k \rightarrow \infty} \log|B_k(z)|$ is subharmonic on Π^+ and differs from $-\infty$. Let $u(z)$ be the harmonic majorant of $\mu(z)$ in Π^+ . Since $\mu(z) \leq 0$ we have that $u(z) \leq 0$. Let $\phi(z) = e^{u(z)+v(z)}$, where $v(z)$ is the harmonic conjugate of $u(z)$.

2. RESULTS

Theorem 1. [4] Assume that (1.5) holds and that ϕ is as above. For a continuous function F on R there exists a sequence $\{R_k\}$ of rational functions of the form (1.4) which converges uniformly on R to F if and only if F coincide almost everywhere on R with the boundary value of meromorphic function F on Π^+ of the form

$$F(z) = \frac{\psi(z)}{B(z)\phi(z)}, \quad z \in \Pi^+, \quad (1.6)$$

where ψ is any bounded analytic function on Π^+ .

Let σ be a nondecreasing function of bounded variation on R . By $L_p(d\sigma, R)$, $p > 0$ is denoted the set of all complex valued functions F , for which the Lebesgue-Stieltjes integral exists i.e. $\int_R |F(x)|^p d\sigma(x) < \infty$.

With (1.7) we denote the following condition:

$$\int_R \frac{\log \sigma^+(x)}{1 + x^2} > -\infty \quad (1.7)$$

Theorem 2. Assume (1.5) and (1.7). For a function $F \in L^p(d\sigma, R)$, $p > 0$ there exists a sequence $\{R_k\}$ of rational functions of the form (1.4) such that

$\lim_{k \rightarrow \infty} \int_{\mathbb{R}} |F(x) - R_k(x)|^p d\sigma(x) = 0$ if and only if F coincide almost everywhere on \mathbb{R} with the boundary value of a meromorphic function F on Π^+ of the form (1.6), where B and φ are as in theorem 1, and ψ is analytic function on Π^+ of the class N^+ .

Note. N^+ is the class of all analytic functions on N^+ which satisfy the following condition

$$\lim_{y \rightarrow 0^+} \int_{\mathbb{R}} \frac{\log^+ |f(x + iy)|}{1 + x^2} dx = \int_{\mathbb{R}} \frac{\log^+ |f(x)|}{1 + x^2} dx$$

$S = S(\mathbb{R}^n)$ denotes the space of all infinitely differentiable complex valued function φ on \mathbb{R}^n satisfying

$$\sup_{t \in \mathbb{R}^n} |t^\beta D^\alpha \varphi(t)| < \infty$$

for all n -tuple α and β of nonnegative integers. Convergence in S is defined in the following way: a sequence $\{\varphi_\lambda\}$ of functions $\varphi_\lambda \in S$ in S as $\lambda \rightarrow \lambda_0$ if and only if

$$\lim_{\lambda \rightarrow \lambda_0} \sup_{t \in \mathbb{R}^n} |t^\beta D^\alpha [\varphi_\lambda(t) - \varphi(t)]| = 0$$

for all n -tuple α and β of nonnegative integers.

Again, S' is the space of all continuous, linear functionals on S , called the space of tempered distributions.

The space S' is called the space of tempered distributions. We use the convention $\langle T, \varphi \rangle = T(\varphi)$ for the value of the functional T acting on the function φ .

Let $\varphi \in S$ and $f(x) \in L^1_{loc}(\mathbb{R}^n)$. Then the functional T_f on S defined with

$$\langle T_f, \varphi \rangle = \int_{\mathbb{R}^n} f(t)\varphi(t)dt, \varphi \in S,$$

is an element in S' and it is called the regular distribution generated by the function f .

Let f is a locally integrable function on \mathbb{R} . With T_f we denote the corresponding regular distribution defined by

$$\langle T_f, \varphi \rangle = \int_{\mathbb{R}} f(x)\varphi(x)dx, \quad \varphi \in S.$$

Theorem 3. Let $z_{k_1}, z_{k_2}, \dots, z_{k_{N_k}}, k = 1, 2, \dots, \text{Im}z \neq 1, N_k \leq \infty$, are given complex numbers which satisfy (1.5) and F be of the form (1.6) (as in Theorem 2). Let $T_{F^*}, F^* \in L^p(\mathbb{R})$, be the distribution in S' generated by the boundary value $F^*(x)$ of $F(z)$ on Π^+ .

Then there exists a sequence, $\{R_k\}$, of rational functions of the form (1.4) and, respectively, a sequence of distributions $\{T_{R_k}\}, T_{R_k} \in S'$ generated by R_k , satisfying

- (i) $T_{R_k} \rightarrow T_{F^*}, k \rightarrow \infty$ in S'
- (ii) $\overline{\lim}_{n \rightarrow \infty} \int_{-\infty}^{\infty} |R_k(x)|^p |\varphi(x)| dx < \infty$, for all $\varphi \in S$.

Proof.

(i) General idea is to prove the convergence in S' in weak sense and to use Banach Steinhaus theorem to obtain the strong convergence (note that the space S is of second category).

We start with applying Theorem 2, and obtain a sequence $\{R_k\}$, of rational functions of the form (1.4) for which the following holds

$$\lim_{k \rightarrow \infty} \int_{\mathbb{R}} |F^*(x) - R_k(x)|^p dx = 0. \quad (2.1)$$

Triangle inequality implies

$$\begin{aligned} \|R_k\|_{L^p(\sigma)} &= \left(\int_{\mathbb{R}} |R_k(x)|^p d\sigma(x) \right)^{\frac{1}{p}} = \left(\int_{\mathbb{R}} |R_k(x) - F^*(x) + F^*(x)|^p d\sigma(x) \right)^{\frac{1}{p}} \\ &\leq \left(\int_{\mathbb{R}} |R_k(x) - F^*(x)|^p d\sigma(x) \right)^{\frac{1}{p}} + \left(\int_{\mathbb{R}} |F^*(x)|^p d\sigma(x) \right)^{\frac{1}{p}} < C, \end{aligned}$$

Hence $R_k(x) \in L^p(\mathbb{R})$.

Now choose arbitrary $\varphi \in S$ and fix it. We denote with q the Hölder conjugate of p , i.e.

$$\frac{1}{p} + \frac{1}{q} = 1,$$

$$\left| \int_{\mathbb{R}} f(x)\varphi(x) dx \right| \leq \int_{\mathbb{R}} |f(x)\varphi(x)| dx \leq \left(\int_{\mathbb{R}} |f(x)|^p dx \right)^{\frac{1}{p}} \left(\int_{\mathbb{R}} |\varphi(x)|^q dx \right)^{\frac{1}{q}}$$

for arbitrary function $f \in L^p$, $\varphi \in S$.

We use the previous and obtain

$$\begin{aligned} |\langle T_{R_k}, \varphi \rangle - \langle T_{F^*}, \varphi \rangle| &= \left| \int_{-\infty}^{\infty} R_k(x) \varphi(x) dx - \int_{-\infty}^{\infty} F^*(x) \varphi(x) dx \right| \\ &= \left| \int_{-\infty}^{\infty} [R_k(x) - F^*(x)] \varphi(x) dx \right| \leq \int_{-\infty}^{\infty} |R_k(x) - F^*(x)| |\varphi(x)| dx \\ &\leq \left(\int_{\mathbb{R}} |R_k(x) - F^*(x)|^p dx \right)^{\frac{1}{p}} \left(\int_{\mathbb{R}} |\varphi(x)|^q dx \right)^{\frac{1}{q}} \end{aligned}$$

$$\leq M \left(\int_{\mathbb{R}} |R_k(x) - F^*(x)|^p dx \right)^{\frac{1}{p}} \rightarrow 0, \quad (2.1) \text{ when } k \rightarrow \infty.$$

In the previous calculations we used $M = \sup\{|\varphi(x)| \mid x \in \mathbb{R}\}$ which is obviously finite since the inclusion $S \subset L^q$ is continuous and dense for arbitrary $1 \leq q < \infty$. The discussion on the start of the proof implies the claim or $T_{R_k} \rightarrow T_{F^*}$, $k \rightarrow \infty$ in S' in the strong topology.

(ii) Let $\varphi \in S$ be arbitrary and fixed. Then Minkowski inequality implies

$$\begin{aligned} \left(\int_{\mathbb{R}} |R_k(x)|^p \varphi(x) dx \right)^{\frac{1}{p}} &\leq M^{\frac{1}{p}} \left(\int_{\mathbb{R}} |R_k(x) - F^*(x) + F^*(x)|^p dx \right)^{\frac{1}{p}} \\ &\leq M^{\frac{1}{p}} \left(\int_{\mathbb{R}} |R_k(x) - F^*(x)|^p dx \right)^{\frac{1}{p}} + M^{\frac{1}{p}} \left(\int_{\mathbb{R}} |F^*(x)|^p dx \right)^{\frac{1}{p}} = I_1 + I_2. \end{aligned}$$

The integral I_1 tends to 0 when $k \rightarrow \infty$ which implies that $\int_{\mathbb{R}} |R_k(x)|^p \varphi(x) dx \leq M^{\frac{1}{p}} \|F^*\|_p + C$, for arbitrary $k \in \mathbb{N}$.

The latter implies (ii).

3. CONCLUSION

We were able to consider (tempered) distribution variant of Tumarkin result concerning approximation with rational functions. We embed L^p functions continuously into S' in a natural way and consider analog result, but now for their representation in S' .

AUTHORS' CONTRIBUTIONS

This work was carried out in collaboration among all authors and equally contributed to this article. All authors read and approved the final manuscript.

REFERENCES

- [1] Bremermann H. Distribution, Complex Variables and Fourier Transforms, Addison-Wesley Publishing Co., Inc., Reading, Massachusetts-London; 1965.
- [2] Beltrami EJ, Wohlers MR, Distributions and the boundary values of analytic functions, Academic Press, New York; 1966.
- [3] Carmichael R, Mitrovic D, Distributions and analytic functions, Pitman Research Notes in Mathematics Series, 206, Longman Scientific & Technical, Harlow; John Wiley & Sons, Inc., New York; 1989.
- [4] Duren PL, Theory of H^p Spaces, Acad. Press, New York; 1970.
- [5] Tumarkin GT, Approximation with respect to various metrics of functions defined on the circumference by sequences of rational fractions with fixed poles, Akad. Nauk SSSR Ser. Mat. 1966; 30(4):721-766.
- [6] Jantcher L, Distributionen, Walter de Gruyter, Berlin, New York; 1971.
- [7] Manova E V, Bounded subsets of distributions in D' generated with boundary values of functions of the space H^p , $1 \leq p < \infty$, Godisen zbornik na Institutot za matematika, Annuaire, ISSN 0351-7241. 2001: 31-40.
- [8] Manova EV, Pandevski N, Nastovski L, Distribution analogue of the Tumarkin result, Bulletin T. CXXXIII del' Akademie serbedes sciences et des arts, Classe des Sciences mathematiques et naturelles Sciences mathematiques, 2006: No. 31.

- [9] Reckovski N, One proof for the analytic representation of distributions, *Matematički Bilten* 28 (LIV), Skopje, 2004:19-30.
- [10] Reckovski N, Manova EV, Bedjeti B, Iseni E, For some boundary value problems in distributions, *Journal of Advances in Mathematics*, Volume 16, 2019: 8331-8339.
- [11] Rudin W, *Real and Complex Analysis*, McGraw-Hill Inc., New York; 1966.
- [12] Rudin W, *Functional Analysis*, North-Holland publishing company, Amsterdam; 1974.