

Original Research Article

Reproductive Endocrinologic Pattern in Infertile Black Women with Polycystic Ovarian Syndrome

Running Title: Reproductive Endocrinology in PCOS Infertile Women

Abstract

Background: Polycystic ovarian syndrome (PCOS) is a very common endocrine cause of infertility affecting about 10% of women of reproductive age. We evaluated the pattern of reproductive hormones in infertile Nigerian women with PCOS.

Methods: This was a prospective cross-sectional study at the Lagos State University Teaching Hospital over a 6-month period in which One hundred and fifty infertile women, with diagnosis of PCOS using the Rotterdam's criteria, had quantitative assessment of their reproductive hormones like Luteinizing Hormone (LH), Follicle Stimulating Hormone (FSH), Progesterone, Testosterone, Prolactin and Thyroid Stimulating Hormone (TSH), with relevant socio-demographic and clinical data noted in the study proforma. Z test, chi-square and correlation tests were used as appropriate to analyse the data with $p < 0.05$ significance level.

Results: The mean age of infertile women with PCOS was 26.50 ± 4.4 years; all the women had either oligomenorrhea (62%) or secondary amenorrhea (38%), 56.5% were obese and 33.3% had hirsutism. While 37.7% of the women had abnormally elevated serum LH, 16.4% had high serum FSH, 39% had LH:FSH ratio > 2.5 , 96% had low serum progesterone, 29% had high serum prolactin, 17.4% had high serum TSH values and 92% had features of polycystic ovaries on trans-vaginal ultrasound. Correlation of age and BMI with serum FSH, LH and prolactin values were weak and insignificant.

Conclusion: PCOS is common among young Nigerian women presenting for infertility treatment; with dominant features of oligomenorrhea and polycystic ovaries. Hormonal abnormalities are common, varied and not associated with other clinical characteristics.

Funding: No funding sources

Key Words:PCOS, Polycystic Ovarian Syndrome, Infertility, Rotterdam's Criteria, Hormonal Profile.

Introduction

Infertility is the inability of a couple to achieve conception after one year of regular unprotected sexual intercourse [1]. It is a condition constituting both clinical and social challenge to about one couple in six [1]. Infertility is said to be primary (when a woman has never conceived) or secondary and secondary infertility (the incapability to conceive in a couple who have had at least one successful conception in the past) [2]. The causes of infertility could be male factor from seminal fluid abnormalities or female factor such as ovulation dysfunction and tubal pathology [2]. Polycystic ovarian syndrome (PCOS) also called polycystic ovary syndrome, is a common cause of ovulation dysfunction and thus contributes to infertility [3].

Polycystic ovary syndrome (PCOS) is a common endocrinologic disorder in women that affects reproductive function [3]. It occurs in 5-10% of women of child bearing age and its exact cause is not known [4,5]. The Rotterdam 2003 criteria require at least presence of two of the following three conditions for the diagnosis of PCOS: Oligo- or anovulation (menstrual irregularity), clinical or biochemical signs of hyperandrogenism and polycystic ovaries on ultrasound [6]. PCOS is a well-established medical condition that negatively affects reproduction, general health, sexual health and quality of life [7].

Less is known about the influence of race on the PCOS phenotype [8]. Emerging evidence suggests that ethnicity may be associated with PCOS phenotype due to varying genetic and environmental propensity to metabolic and hormonal abnormalities [9-11]. In developed countries, although much has been clarified, on the pathogenesis, diagnosis, clinical manifestations, and treatment of PCOS, [7] adequate profiling of women with PCOS in developing countries and black African populations is relatively sparse. We, therefore, assessed the pattern of reproductive hormone levels in infertile Nigerian women with PCOS.

UNDER PEER REVIEW

Materials and Methods

This was a prospective cross-sectional study of 150 women with PCOS who were being evaluated for infertility at the gynaecology clinic of the Lagos State University Teaching Hospital between January and June 2017. Consenting women aged 18-45 years with history of primary or secondary infertility were recruited while women on ovulation induction medications, with co-existing ovarian tumours and on chemo- and/or radio-therapy were excluded from the study.

Each recruited participant completed a structured administered questionnaire to obtain socio-demographic data, sexual history, obstetrics history, medical history and menstrual history. The anthropometric measurements, weight and height of each subject were taken using a calibrated weighing scale (Medfield equipment and Scientific Ltd. England) and a meter height scale (Avery Co. Ltd, England). The women had their blood samples taken at recruitment, irrespective of the day of their menstrual cycle due to the widespread menstrual irregularity in this group of women. Blood samples were used for hormonal analysis and serum LH, FSH, Prolactin, TSH and Testosterone levels determined according to standard.

The study participants were examined in a well-lit changing room for facial, chest, abdominal, arms, thighs and back hair distribution to give the Ferriman and Gallwey score for hirsutism. Diagnosis of hirsutism was made if the score was greater than 8.

Technique of Trans-vaginal Ultrasound

Each participant was asked to empty her urinary bladder and then taken into the examination room ensuring adequate privacy with a female chaperone in attendance. She was placed in a supine lithotomy position with both knees flexed on the examination couch with adequate exposure of the vulva. The examination area was sufficiently screened to ensure privacy. All scans were performed in a systemic manner using the trans-vaginal 7.5 to 12 MHz Broad

Band ultrasound curved-array transducer. A Mindray Real Time Ultrasound Scanner Model Z5 was used. Folded sheets or pads were placed under the participant's buttocks to elevate her pelvis above the examination couch to allow room for transducers manipulation. Ultrasound coupling gel was applied into a latex male condom and used to completely cover the trans-vaginal probe. A different condom was used for each woman. After parting the labia, the probe covered with condom was then gently introduced into the participant's vagina advancing slowly toward the cervix. The transducer was then moved laterally to the adnexal area to properly visualize the ovaries. During the trans-vaginal ultrasound scanning, the ovaries could be brought into view by using one hand to compress the lower abdominal wall while the other hand manipulated the transducer. The trans-vaginal ultrasound image of the ovary in longitudinal plane and was frozen for measurements. The ovaries were often identified by the presence of follicles, which appeared hypoechoic or anechoic. Once scanning was completed in the longitudinal plane, the probe was rotated 90 degrees counterclockwise so that the transducer was now in a transverse plane and frozen for measurements. After identification of the ovaries, the size of the ovary was measured in three orthogonal planes, namely; longitudinal (length) transverse (thickness) and anteroposterior (width). On measuring the dimensions, inherent volume software in the machine automatically calculated the volume. The measurements were obtained from the frozen ultrasound images with electronic calipers. The total numbers of follicles in each ovary were counted. Follicles were counted on the frozen images of two non-overlapping planes in the longitudinal section of each ovary and summed up together. Measurements of the follicles were made in their maximum internal diameters and recorded in mm; their distributions were also noted as either peripherally or randomly.

The diagnosis of polycystic ovary (PCO) morphology was made if 12 or more follicles each measuring 2-9mm in diameter were present and peripherally arranged in the ovaries (giving a

string of pearl appearance) and/or ovarian volume was increased greater than 10mls [12]. A description of the distribution of the follicles was recorded by visual assessment of location of follicles as being scattered throughout the stroma or arranged peripherally in the classically described “string of pearl” at the margin of the ovary. Stromal echogenicity was defined as normal or increased (echogenic) in comparison to the myometrium whose echogenicity is greater than normal ovarian stroma [12,13]. To reduce intra observer variability, the measurements were taken thrice and an average obtained. At the end of the procedure, the probe was withdrawn gently and slowly, patient was cleaned up with tissue paper and politely asked to stand and dress up.

The hirsutism scores using the Ferriman/ Gallwey scoring system and trans-vaginal sonographic ovarian findings (ovarian volume, follicular number, size, distribution and stromal echogenicity) for each woman were recorded in the study proforma for subsequent data analysis.

Statistical analysis

The data obtained from the questionnaires were coded, imputed into the computer and analyzed using SPSS 20.0 statistical software. Percentage and proportions were determined for categorical variables. Pearson’s Chi-square (test for association) was used to assess the significance of relationships between categorical variables. Z test and Pearson’s correlation were used to compare continuous variables. P-values less than 0.05 were considered to be statistically significant at a confidence level less than 95%. Ethical approval was obtained from the Health Ethics and Research Committee of the Lagos State University Teaching Hospital.

Results

A total of 150 women, who met the Rotterdam diagnostic criteria for PCOS, being evaluated for infertility were studied. The mean age of the women with PCOS in this study was 26.50years with the majority (76%) ranging between age 21 and 30years (Table 1). All the women had some degree of menstrual abnormalities with oligomenorrhea occurring in 62.0% and amenorrhea in 38.0% (Table 1). More than half (56.5%) of the women were obese, 33.3% had hirsutism, 29.3% had acne and 92.0% had ultrasound findings of polycystic ovaries (Table 1).

In the study population, the median values of serum LH was 14.9mIU/ml, serum FSH was 6.0mIU/ml, serum prolactin was 13.7ng/dl, serum TSH was 2.1mIU/ml and serum testosterone was 1.9ng/ml (Table 2). Majority (62.3%) of the women had high serum LH values, 83.6% had normal serum FSH levels, 39% had a high LH:FSH ratio, 96% had low serum progesterone values and 29% had elevated serum prolactin levels (Table 3).

Clinical and endocrinologic values were similar in women with PCOS who had polycystic ovaries on ultrasonography when compared with women without polycystic ovaries (Table 4). Correlation of age and body mass index (BMI) with Hormonal values of FSH, LH and prolactin were weak and insignificant (Table 5).

Table 1: Socio-demographic and clinical characteristics of the women

| Variable | Frequency (N = 150) | Percentage (%) |
|----------------------------|--------------------------------|---------------------------|
| Age (in years) | | |
| <20 | 9 | 6.0 |
| 21 – 30 | 114 | 76.0 |
| 31-40 | 27 | 18.0 |
| <i>Mean ± SD</i> | <i>26.50±4.4</i> | |
| <i>Range</i> | <i>16-40</i> | |
| Education | | |
| None | 14 | 9.3 |
| Primary | 8 | 5.3 |
| Secondary | 33 | 22.0 |
| Tertiary | 95 | 63.4 |
| Religion | | |
| Christianity | 106 | 70.7 |
| Islam | 44 | 29.3 |
| Marital Status | | |
| Single | 62 | 41.3 |
| Married | 88 | 58.7 |
| Menstrual symptoms | | |
| Oligomenorrhea | 93 | 62.0 |
| Secondary amenorrhea | 57 | 38.0 |
| Body Mass Index | | |
| Underweight | 20 | 16.4 |
| Normal | 24 | 19.7 |
| Overweight | 9 | 7.4 |
| Obese | 69 | 56.5 |
| Hirsutism | | |
| Yes | 50 | 33.3 |
| No | 100 | 66.7 |
| Acne | | |
| Yes | 44 | 29.3 |
| No | 106 | 70.7 |
| Ultrasound features | | |
| Polycystic Ovary | 138 | 92.0 |
| Non-Polycystic Ovary | 12 | 8.0 |

Table 2: Median Values of Hormonal Levels in Women with PCOS

| Hormone | Median | IQR |
|----------------------|---------------|------------|
| LH(mIU/ml) | 14.9 | 15.8 |
| FSH(mIU/ml) | 6.0 | 4.1 |
| Prolactin (ng/dl) | 13.7 | 12.0 |
| TSH (mIU/ml) | 2.1 | 2.8 |
| Testosterone (ng/ml) | 1.9 | 1.8 |

*IQR – Inter quartile range, LH - Luteinizing hormone,
FSH -Follicular stimulating hormone, TSH – Thyroid Stimulating Hormone*

UNDER PEER REVIEW

Table 3: Distribution of hormonal levels in women with PCOS

| Hormone | Serum Level | Frequency | % |
|--------------------|-----------------------------|-----------|------|
| LH | Normal = 0.5-10.9mIU/ml | 55 | 37.7 |
| | High \geq 11mIU/ml | 91 | 62.3 |
| FSH | Normal = 2.0-12mIU/ml | 122 | 83.6 |
| | High \geq 12mIU/ml | 24 | 16.4 |
| LH / FSH ratio | Normal ratio \leq 2.5 | 89 | 61.0 |
| | High ratio $>$ 2.5 | 57 | 39.0 |
| Progesterone | Normal \geq 20ng/ml | 5 | 4 |
| | Low $<$ 20ng/ml | 145 | 96 |
| Total Testosterone | Normal = 15ng/ml-70ng/ml | 120 | 95.2 |
| | Low $<$ 15ng/ml | 6 | 4.8 |
| Prolactin | Normal level \leq 19ng/dl | 103 | 71.0 |
| | High level $>$ 19.0ng/dl | 42 | 29.0 |
| TSH | Normal \leq 6.2mIU/ml | 90 | 82.6 |
| | High $>$ 6.2mIU/ml | 19 | 17.4 |

LH - Luteinizing hormone, FSH -Follicular stimulating hormone, TSH – Thyroid Stimulating Hormone

Table 4: Comparison of PCOS women with polycystic ovaries versus women without polycystic ovaries on ultrasound

| Variables | PCO | Non-PCO | z-test | p-value |
|---------------------|--------------|-------------|---------------------|---------|
| | Median ± IQR | Median± IQR | | |
| Weight | 72 ± 18 | 43.5± 54.2 | 1.673 ^z | 0.094 |
| Height | 1.61 ± 0.09 | 1.61± 13.9 | -0.364 ^z | 0.715 |
| BMI | 42.5 ± 40.5 | 5.5±43.5 | 1.623 ^z | 0.104 |
| LH | 15.0 ± 15.7 | 7.3±1.1 | 1.313 ^z | 0.189 |
| FSH | 6.0± 5.0 | 5.2±3.0 | 0.186 ^z | 0.852 |
| Prolactin | 11.8± 11.8 | 23.4±16.7 | -0.961 ^z | 0.336 |
| TSH | 2.3± 2.7 | 2.0±1.1 | 0.863 ^z | 0.388 |
| Testosterone | 1.9± 1.5 | 1.9±1.3 | -0.094 ^z | 0.924 |
| T3 | 5.3± 2.0 | 1.1± 0.0 | 1.451 ^z | 0.146 |
| T4 | 10.5± 44.5 | 15.0± 0.0 | -0.707 ^z | 0.479 |

z- test for non-parametric test applied, PCO-Polycystic ovary, LH - Luteinizing hormone, FSH -Follicular stimulating hormone, TSH – Thyroid Stimulating Hormone, T3- Triiodothronine, T4-Thyroxine.

Table 5: Correlation of age and BMI with hormonal values in women with PCOS

| | Correlation coefficient | p-value |
|--|--------------------------------|----------------|
| Age and FSH | 0.127 | 0.127 |
| Age and Prolactin | 0.102 | 0.221 |
| Age and LH | 0.090 | 0.278 |
| Age and LH/FSH ratio | 0.082 | 0.162 |
| BMI and FSH | -0.095 | 0.299 |
| BMI and Prolactin | -0.057 | 0.533 |
| BMI and LH | 0.005 | 0.954 |
| BMI and LH/FSH ratio | 0.007 | 0.345 |
| <i>LH - Luteinizing hormone, FSH -Follicular stimulating hormone, TSH – Thyroid Stimulating Hormone, BMI-Body Mass Index</i> | | |

Discussion

The mean age of the women with PCOS in this study was 26.5years with about three-fourth of them (76%) being within the age range 21 to 30years. This is similar to the mean age of 27years reported in women with PCOS by Igwegbe et al [5] in a study in Nnewi, south-eastern Nigeria but lower than the mean age of 30years and 31.5years, in women with PCOS, observed by Ugwu et al [14] in Enugu and Omokanye et al [15] in Ilorin, Nigeria respectively. PCOS is a condition in women of reproductive age group and this explains all the mean ages in these studies being within this category.

None of the women had normal menstrual cycle length, with oligomenorrhea occurring more frequently (62%) than amenorrhea which was present in 38% of the women (Table 1). Of these women with PCOS in our study, we observed that more than half (56.5%) of the women were obese, 33.3% had hirsutism, 29.3% had acne and 92% had ultrasound findings of polycystic ovaries. The abnormal menstrual cycle length is very likely due to anovulation, a common finding in women with PCOS. Oriji et al [16] in Port Harcourt Nigeria noted that majority of women (65.5%) with PCOS in their study had normal cycle lengths, 27.5% had oligomenorrhea and 41.3% had acne. March et al in an Australian study also found that a lower proportion (23.8%) of women with PCOS were oligomenorrhic. This variation with our findings may be due to the small number of women with PCOS in the study by Oriji et al [16] and phenotypic differences with women in the Australian study [17].

Oriji et al however found that 92% of infertile women with PCOS had at least one polycystic ovary; this finding is in keeping with observations in our study [16]. Reports, by Ugwu et al [14] in south east Nigeria, of oligomenorrhea in 72.6%, obesity in 51.6% and a LH/FSH ratio > 2 in 45.2% in women with PCOS are similar to findings of our study. We noted that 39.0% of women with PCOS had a LH/FSH ratio of > 2.5 . The finding of high FSH in 16.4% of women in this study, among women with a mean age of 26years, is not typical of PCOS. This

may be because the collection of blood samples was done at recruitment, irrespective of the day of their menstrual cycle, as most of the women were either oligomenorrheic or amenorrheic. Basal FSH is best determined on the 3rd day of a woman's menstrual cycle.

We observed elevated TSH in 17.4% of women with PCOS. Although the relationship between PCOS and hypothyroidism is complex and not yet fully understood there are reports that hypothyroidism can lead to polycystic morphology of the ovaries [18]. Though this morphology can vary with the severity and duration of hypothyroidism, there is currently no evidence to suggest that hypothyroidism can lead to PCOS [18].

Median values of weight, height, body mass index (BMI) and serum LH, FSH, Prolactin, TSH, Testosterone, T3 and T4 were similar when women with PCOS with ultrasound features of polycystic ovaries were compared with women without ultrasound features of polycystic ovaries, but met the diagnostic criteria for PCOS. We also noted an unusual finding of normal total testosterone values in our study population as none of the women had elevated total testosterone >70ng/ml. These findings probably suggest that the presence of the polycystic ovarian morphology typical of PCOS is not a mandatory pre-requisite for anovulation or infertility in women with PCOS. Neither age nor BMI correlated with serum values of LH, FSH and prolactin in women with PCOS.

Conclusion

In conclusion, PCOS is common among young Nigerian women presenting for infertility treatment; with dominant features being oligomenorrhea and polycystic ovaries. Reproductive hormonal abnormalities are common, varied and not associated with other clinical characteristics in these women.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly used products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

References

1. Kamel, R.M. Management of the infertile couple: an evidence-based protocol. *Reprod Biol Endocrinol.* 2010; 8:21. <https://doi.org/10.1186/1477-7827-8-21>
2. Benksim A, Elkhoudri N, Addi RA, Baali A, Cherkaoui M. Difference between Primary and Secondary Infertility in Morocco: Frequencies and Associated Factors. *Int J Fertil Steril.* 2018;12(2):142-146. doi:10.22074/ijfs.2018.5188
3. Couto Alves, A., Valcarcel, B., Mäkinen, VP. et al. Metabolic profiling of polycystic ovary syndrome reveals interactions with abdominal obesity. *Int J Obes.* 2017; 41:1331–1340. <https://doi.org/10.1038/ijo.2017.126>
4. March WA, Moore VM, Wilson K.J. The prevalence of polycystic ovary syndrome in a community sample assessed under contrasting diagnostic criteria. *Hum. J. Reprod.* 2010;25 (2):554-6
5. Igwegbe AO, Eleje GU, Enechukwu C I. Polycystic ovary syndrome: A Review of management outcome in a low resource setting. *J Women's Health.* 2013; 2:3

6. Rotterdam ESHRE/ASRM-Sponsored PCOS Consensus Workshop Group. Revised 2003 syndrome. *Fertil Steril*. 2004;81(1):19–25.
7. Aversa A, La Vignera S, Rago R, Gambineri A, Nappi RE, Calogero AE and Ferlin A. Fundamental Concepts and Novel Aspects of Polycystic Ovarian Syndrome: Expert Consensus Resolutions. *Front. Endocrinol*. 2020; 11:516. doi: 10.3389/fendo.2020.00516
8. Ladson G, Dodson WC, Sweet SD, et al. Racial influence on the polycystic ovary syndrome phenotype: a black and white case-control study. *FertilSteril*. 2011;96(1):224-229.e2. doi:10.1016/j.fertnstert.2011.05.002
9. Louwers, Y., Lao, O. and Kayser, M. Inferred Genetic Ancestry versus Reported Ethnicity in Polycystic Ovary Syndrome (PCOS). *Human Reproduction*. 2013;28:349-349.
10. Sirota, I., Stein, D.E., Vega, M. and Keltz, M.D. Increased Insulin-Resistance and Beta-Cell Function in Polycystic Ovary Syndrome Women—Does Ethnicity Play a Role? *Reproductive Science*. 2013;20:180a-181a.
11. Akpata, C., Uadia, P. and Okonofua, F. Prevalence of Polycystic Ovary Syndrome in Nigerian Women with Infertility: A Prospective Study of the Three Assessment Criteria. *Open Journal of Obstetrics and Gynecology*. 2018;8:1109-1120. doi: 10.4236/ojog.2018.812112.
12. Hart T, Hickey M, Frank S. Definition, Prevalence and symptoms of polycystic ovary and polycystic ovary syndrome. *Best Prat Res Clin ObstetGynaecol*. 2004; 18 (5):671-783.
13. Wolfgang Dahnert, *Radiology Review Manual*. 6th Edition. Lippincott Williams and Wilkins. 2007: 1068.

14. Ugwu GO, Iyoke CA, Onah HE, Mba SG. Prevalence, presentation and management of polycystic ovary syndrome in Enugu, South East Nigeria. *Niger J Med* 2013;22:313-6
15. Omokanye L, Ibiwoye-Jaiyeola O, Olatinwo AWO, Abdul I, Durowade K, Biliaminu S. Polycystic ovarian syndrome: Analysis of management outcomes among infertile women at a public health institution in nigeria. *The Nigerian Journal of General Practice*. 2015;13(2):44-8.
16. Vaduneme Kingsley Oriji, Chidi Onwuegbulam. Prevalence of Polycystic Ovary Syndrome (PCOS) among Infertile Women Attending Fertility Clinic at a University Teaching Hospital in Nigeria. *J Gynecol Women's Health*. 2019: 15(5): 555922. DOI: 10.19080/JGWH.2019.15.555922
17. March, W.A., Moore, V.M., Willson, K.J., Phillips, D.I., Norman, R.J. and Davies, M.J. The Prevalence of Polycystic Ovary Syndrome in a Community Sample Assessed under Contrasting Diagnostic Criteria. *Human Reproduction*. 2010; 25:544-551. <https://doi.org/10.1093/humrep/dep399>.
18. Singla R, Gupta Y, Khemani M, Aggarwal S. Thyroid disorders and polycystic ovary syndrome: An emerging relationship. *Indian J Endocrinol Metab*. 2015;19(1):25-29. doi:10.4103/2230-8210.146860.