

Cytokines (IL-17, IL-23 and IL-33) in adult patients with Systemic lupus erythematosus in Trinidad and Tobago.

Abstract:

Systemic lupus erythematosus (SLE) is the most common autoimmune disease. It is characterized by the presence of hundreds of autoantibodies against many organs and tissues, including the presence of a large number of autoantibodies, which are specific to self-antigens mainly of nuclear origin such as Smith antigen, double-stranded DNA (dsDNA), anti-Sjögren's syndrome-related antigen A and B (SSA/Ro and SSB/La, respectively) and ribonucleoproteins, which are the hallmarks of the disease. Type I and II interferons, interleukin-6 (IL-6), IL-1, tumor necrosis factor-alpha (TNF- α), and immunomodulatory cytokines such as IL-10 and TGF- β are essential players in SLE. Additionally, T-cell-derived cytokines such as IL-17, IL-21, and IL-2 are dysregulated in SLE. In this study among cohorts of 60 individuals attending the hospital clinics in Trinidad and Tobago, blood samples were analyzed and the levels of the essential cytokines were measured using SLE Disease Activity Index (SLEDAI) 2000 score. The results confirmed that serum IL-17 and IL-23 levels were positively correlated with the SLE Disease Activity Index (SLEDAI) 2000 score in these patients. These findings have diagnostic and therapeutic implications. However, more work must be done targeting other cytokines relevant to autoimmunity and SLE in particular. Interleukin-33 is not an SLE marker, as has been noted in other populations.

Introduction

Studies have provided new information on interleukins 17, 23, and 33 and systemic lupus erythematosus (SLE). In their study, Wong et al., 2008 showed that the production of IL-17 was uncharacteristically high in SLE patients, and it showed a positive correlation between IL-17 and the SLEDAI score [1]. Fabien et al., in 2013, showed that IL-17 serum levels were higher in patients with lupus than controls. However, these interleukin levels did not correlate with the SLE Disease Activity Index (SLEDAI) 2000 score [2]. A previous study also found that the association between IL-23 and the IL-17 axis is vital for the inflammatory response in SLE [1]. In a study conducted by Mok et al. (2010), their investigations showed no correlation was found between serum IL-17 and 23 in SLE patients with SLE. This study also showed higher serum IL-17 and IL-23 levels, and the lack of correlation between these cytokines suggests independent regulatory mechanisms [3]. A study in 2011 by Yang et al. found that IL-33 does not play a part in the illness but is needed in the acute phase and affects red blood cells and platelets [4]. The rationale for our study was the measurements of IL-17, IL-23, and IL-33 in the serum of patients suffering from systemic lupus erythematosus in Trinidad and Tobago; studying and determining these cytokines will

be a tremendous diagnostic means and in further management of SLE patients in the country and the region.

Materials and Methods

Population, study design, and setting.

All the experiments were conducted in accordance with the Declaration of Helsinki (1964). This study describes experimental work that carries a risk of harm to human subjects, and it was conducted with the human subjects' understanding and consent, as well as a statement that the Ethical Committee of the University of the West Indies has approved the experiments.

A previous research project with fewer subjects was conducted [5]. This prospective, cross-sectional, observational study was performed. Thirty (30) patients attending the rheumatoid clinic at one of the major regional hospitals in Trinidad and Tobago were recruited. There was no interventional measure. The patients were assessed and treated after the blood was taken and were clinically assessed. However, the study sought to observe and delineate the presence of IL-17, 23, and 33 among patients attending a rheumatoid outpatient clinic. Using the Revised American College of Rheumatology criteria, lupus diagnosis was confirmed, and the disease was clinically assessed using the SLE Disease Activity Index (SLEDAI) 2000 score depicted in Table 1 [6]. A score of ≥ 3 or four was considered an active disease, and a score of < 3 was considered an inactive disease. Thirty (30) patients who were non-SLE subjects were also chosen as normal controls for the study.

The characteristics of the study population included all women aged > 18 years with an active SLE diagnosis. Female patients were enrolled in the study because their prevalence was higher in women than in men. One male patient attending the Rheumatology Clinic for SLE was also recruited in the study, as all available men were included. The target population of this study was 60, of which 30 patients with active SLE disease were included.

Inclusion and Exclusion Criteria.

Only adult patients with active SLE disease and all ethnic groups who gave permission to participate in the project were included.

None of the patients with SLE who refused to participate were included. SLE occurs mostly in women above childbearing age. Although there may be children who could suffer from the disease, they did not fall into this age category as adult females of childbearing age.

Laboratory workup as blood cell count and differential, blood chemistry, C3 and C4 levels, antinuclear antibodies, kidney function test as the presence of proteinuria and haematuria, and other blood routine was done by standard laboratory procedure in the Clinical Laboratory Department at Eric Williams Medical Science complex and were collected from the patient records,

Table 1: Describe SLEDAI 2000 (SLEDAI- 2K), a modification of SLEDAI to reflect persistent, active disease in those descriptors that had previously only considered new or recurrent occurrences; and as a measure of global disease activity in the clinic [6].

Study No.: _____ Patient Name: _____ Visit Date: _____

(Enter weight in SLEDAI Score column if descriptor is present at the time of the visit or in the preceding 10 days.)

Weight	SLEDAI SCORE	Descriptor	Definition
8	_____	Seizure	Recent onset, exclude metabolic, infectious or drug causes.
8	_____	Psychosis	Altered ability to function in normal activity due to severe disturbance in the perception of reality. Include hallucinations, incoherence, marked loose associations, impoverished thought content, marked illogical thinking, bizarre, disorganized, or catatonic behavior. Exclude uremia and drug causes.
8	_____	Organic brain syndrome	Altered mental function with impaired orientation, memory, or other intellectual function, with rapid onset and fluctuating clinical features, inability to sustain attention to environment, plus at least 2 of the following: perceptual disturbance, incoherent speech, insomnia or daytime drowsiness, or increased or decreased psychomotor activity. Exclude metabolic, infectious, or drug causes.
8	_____	Visual disturbance	Retinal changes of SLE. Include cytoid bodies, retinal hemorrhages, serous exudate or hemorrhages in the choroid, or optic neuritis. Exclude hypertension, infection, or drug causes.
8	_____	Cranial nerve disorder	New onset of sensory or motor neuropathy involving cranial nerves.
8	_____	Lupus headache	Severe, persistent headache; may be migrainous, but must be nonresponsive to narcotic analgesia.
8	_____	CVA	New onset of cerebrovascular accident(s). Exclude arteriosclerosis.
8	_____	Vasculitis	Ulceration, gangrene, tender finger nodules, periungual infarction, splinter hemorrhages, or biopsy or angiogram proof of vasculitis.
4	_____	Arthritis	≥ 2 joints with pain and signs of inflammation (i.e., tenderness, swelling or effusion).
4	_____	Myositis	Proximal muscle aching/weakness, associated with elevated creatine phosphokinase/aldolase or electromyogram changes or a biopsy showing myositis.
4	_____	Urinary casts	Heme-granular or red blood cell casts.
4	_____	Hematuria	>5 red blood cells/high power field. Exclude stone, infection or other cause.
4	_____	Proteinuria	>0.5 gram/24 hours
4	_____	Pyuria	>5 white blood cells/high power field. Exclude infection.
2	_____	Rash	Inflammatory type rash.
2	_____	Alopecia	Abnormal, patchy or diffuse loss of hair.
2	_____	Mucosal ulcers	Oral or nasal ulcerations.
2	_____	Pleurisy	Pleuritic chest pain with pleural rub or effusion, or pleural thickening.
2	_____	Pericarditis	Pericardial pain with at least 1 of the following: rub, effusion, or electrocardiogram or echocardiogram confirmation.
2	_____	Low complement	Decrease in CH50, C3, or C4 below the lower limit of normal for testing laboratory
2	_____	Increased DNA binding	Increased DNA binding by Farr assay above normal range for testing laboratory.
1	_____	Fever	>38° C. Exclude infectious cause.
1	_____	Thrombocytopenia	<100,000 platelets / x10 ⁹ /L, exclude drug causes.
1	_____	Leukopenia	< 3,000 white blood cells / x10 ⁹ /L, exclude drug causes.
TOTAL SLEDAI SCORE		_____	

Methodology

Patients consented to participate in the study. The demographic data of these patients were collected during this period using a standardized questionnaire. Following this, 5 ml of the venous blood sample was drawn from the patient's arm into red and purple top bottle tubes, and the tubes were taken to the Immunology Laboratory section of the Department of Paraclinical Sciences of the UWI for analysis.

Immunoassays (ELISA) for the determination of the serum concentration of IL-17, IL-23 or IL-33

The sera were extracted and stored at minus20°C until further analysis. Using commercially available sandwich ELISA kits (IBL International, Hamburg, Germany), the sera were tested to determine the presence of IL-23, 17, and 33 according to established immunochemical analysis and the manufacturer's instructions.

Three tests were performed using IBL human IL-17A, IL-23, and IL-33 ELISA kits. Ninety-six-well ELISA plates were lined with monoclonal anti-human antibodies against IL-17, IL-23, or IL-33. The patient samples were added to the plates and incubated. After washing, the wells were incubated with biotin-conjugated anti-human IL-17A, IL-23, or IL-33.

For the IL-17A ELISA kit, and IL-23 ELISA kit protocol followed the guidance as referenced in 5.

For the IL-33 ELISA kit, the wells were incubated with human serum for 1 h, the microplate was washed, and after incubation with the anti-human biotin conjugate for 1 h, the microplate was rewashed, and streptavidin-coupled horseradish peroxidase (HRP) was added. Unbound avidin-HRP was removed by washing, and a reactive substrate solution was added. The levels of IL-33 in the samples were proportional to the amount of colored product produced. The addition of 3M H₂SO₄ stopped the reaction, and the absorbance of the samples and controls was measured at 450 nm. Human IL-33 concentrations were determined using a standard curve drawn from seven human IL-33 standard dilutions.

Statistical analysis

Data were analyzed using Microsoft Excel and SPSS 22 software programs. The data were descriptive, and it was reported as a comparison of frequency distributions. A $p < 0.05$ was considered a statistically significant value. A linear regression test was used to determine the cytokine concentrations in the stated population.

Results

A total of 60 blood samples were drawn from 30 SLE patients and 30 healthy individuals. The crucial observation from the analysis of the age groups of the patients enrolled in this study, most of the patients who had SLE were in the 25 to 30 years age group. For the controls, and the highest frequency was those over 40 years of age. The

least number of participants recruited from the SLE group was under the 20-year age group.

Of all SLE patients, 15 (50%) were of Afro-Trinidadian descent, 12 (40%) were of Indo-Trinidadian descent, and 3 (10%) were of mixed descent or origin. Of all the SLE patients recruited in the investigation, one (3%) was a male patient and 29 (97%) were female. Nineteen healthy controls (63%) were women, and 11 (37%) were men.

These results were related to patient employment and SLE. Most SLE patients were unemployed (47%), while the second most common occupation was a 10% clerk. In non-SLE patients like SLE patients, it was found that many patients were unemployed as well at 57%.

Cytokine concentrations and SLEDAI scores in patients with and without SLE SLEDAI-2K scores ranged from 4 to 14. The mean SLEDAI score was 9.1, and the median score was 9.5. Table 1 shows that many patients (87%) experienced arthritis, and no patients experienced urinary or cardiovascular accidents. Visual disturbances accounted for 35% of patients, while headaches accounted for a significant number of patients at 68%. Vasculitis was seen in 32% of the patients, while renal symptoms such as proteinuria and hematuria accounted for 50% and 30%, respectively. Many patients experienced symptoms of myositis at 77% and alopecia at 60%. Pleurisy accounted for 43% of patients, while low complement and leukopenia values were 60% and 35%, respectively [5-7]. In the control group those manifestations accounted for less than 5%. There was a statistical significance of SLE manifestation percentages as compared with the control group as shown in Table 2.

Figure 1 shows that serum IL-17 and IL-23 levels were more significant in SLE patients than in controls ($P < 0.05$); however, there was no statistically significant difference between IL-33 levels between SLE patients and healthy controls ($P < 0.05$). There was no significant correlation between serum IL-17 levels and the 23 SLE patients with SLE ($r = 0.308$, $p > 0.05$). Serum IL-17 and IL-23 levels were positively correlated with SLEDAI score. Nevertheless, IL-33 levels showed no correlation with SLEDAI score.

Discussion

This study aimed to investigate the association between IL-17, 23, and 33 cytokines in SLE. It determines whether there is any correlation between these cytokines and the disease activity index. In a study conducted by Hegab et al., 2014, they looked at IL-23 serum levels in patients with lupus compared to healthy individuals to correlate the serum levels of cytokines with disease activity and its possible role in the pathogenesis of SLE. Serum levels of IL-23 were determined in all patients and controls using a quantitative enzyme-linked immunosorbent assay. They found that serum IL-23 concentration was significantly elevated in SLE patients compared to healthy controls, and it was significantly correlated with the disease activity index [6]. Their report supports our findings that serum IL-23 levels were higher in SLE patients than in healthy individuals and positively correlated with the SLEDAI score.

A study conducted by Mok et al., 2010 showed that higher serum IL-17 and IL-23 levels were found in SLE patients than in healthy individuals [3]. Another study conducted by Iwakura et al., 2006 showed that IL-23 induced the differentiation of naive CD4 (+) T cells into pathogenic helper T cells that produce IL-17 and other cytokines vital in the inflammatory response. In his paper, it was also reported that there was a correlation between IL-17 and IL-23 levels [8]. This was inconsistent with our results for the Pearson's correlation coefficient, which equaled 0.0308, and there was a p-value of 0.017; and these correlations were statistically significant.

Yang et al., 2011 examined whether IL-33 serum levels were associated with lupus. A total of 70 patients with diseases were recruited. Sera were obtained at their clinical visit and compared to the sera from 40 healthy controls. Serum IL-33 levels were significantly higher in patients with SLE than in healthy controls, but were lower than those found in patients with rheumatoid arthritis [4]. These results are different from the results obtained in our study, which found no statistically significant difference between serum IL-33 levels among SLE patients and healthy controls ($p > 0.05$).

Mok et al., 2010 examined the association between serum IL-33 levels in patients with SLE and disease activity. The SLEDAI was used to assess disease activity. Sandwich ELISA measured the IL-33 levels. Elevated serum IL-33 showed similar values among SLE patients and controls and showed no correlation with the SLEDAI score [9], which was similar to our results, which showed no correlation between IL-33 and disease activity index .

Pons-Estel et al., 2010 showed an increased risk among reproductive-age women who were African Americans. Moreover, SLE is two to four times more frequent and more severe among non-white populations worldwide and tends to be more severe in men, the pediatric population, and patients with late-onset lupus [10]. These results were similar to those obtained in our current analysis, where there was an increased incidence of SLE among female patients of reproductive age who were of African origin.

Izati et al. (2021) demonstrated that therapeutic inhibition of the IL-23/IL-17 axis receptors on Th cells, particularly IL-23R, is potentially relevant in SLE patients [11]. Dai et al. (2017) concluded in a study of cytokines in SLE that IL-23 accounts for the main aspects of human and murine lupus including the expansion of double negative T cells, increased IL-17 production, and decreased IL-2. They proposed that the blockade of IL-23 should have a therapeutic value in patients with SLE [12].

Conclusion

Cytokines IL-17, IL-23 and IL-33 were measured in adult patients with SLE in Trinidad and Tobago. A larger study is needed to corroborate the participation of these cytokines in SLE and how positively they correlate with the SLEDAI-2K score. However, using a sample size representative of the population it was interesting to find in this study that serum IL-17 and IL-23 levels were more significant in SLE patients than in controls, but not IL-33 levels, as well both IL-17 and IL-23 correlated best with the SLEDAI-2K score.

Disclaimer

This paper is an extended version of a **preprint** document of the same author.

The **preprint** document is available in this link: <https://www.medrxiv.org/content/10.1101/2020.09.27.20202762v2.full.pdf>

[As per journal policy, pre-print article can be published as a journal article, provided it is not published in any other journal]

Ethical Approval:

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

Consent

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

References

UNDER PEER REVIEW

1. Wong CK, Lit LC, Tam LS, Li EK, Wong PT, Lam CW. Hyperproduction of IL-23 and IL-17 in patients with systemic lupus erythematosus: implications for Th17-mediated inflammation in auto-immunity. *Clin Immunol* 2008;127:385-93.
2. Fabien B, Northcott M., Hoi A, Mackay F, Morand E. Clinical associations of serum IL-17 in systemic lupus erythematosus. *Arthritis Research & Therapy* 2013; 15:R97.
3. Mok MY, Wu HJ, Lo Y, Lau CS. The relation of IL-17 (IL-17) and IL-23 to Th1/Th2 cytokines and disease activity in systemic lupus erythematosus. *J Rheumatol* 2010; 37:2046-52.
4. Yang Z, Liang Y, Xi W, Li C, Zhong R. Association of increased serum IL-33 levels with clinical and laboratory characteristics of systemic lupus erythematosus in Chinese population. *Clin Exp Med* 2011;11:75-80.
5. Nobe A, Justiz-Vaillant A, Akpaka PE, Poon-King P (2016) Levels of Interleukin 17 and 23 in Patients with Systemic Lupus Erythematosus (SLE) in Trinidad and Tobago. *Immunochem Immunopathol* 2: 115. doi: 10.4172/2469- 9756.1000115
6. Gladman DD, Ibañez D, Urowitz MB. Systemic lupus erythematosus disease activity index 2000. *J Rheumatol.* 2002 Feb;29(2):288-91. PMID: 11838846.
7. Justiz-Vaillant AA, Ferrer-Cosme B, Ramirez-Hernandez N. A Female with Systemic Lupus Erythematosus and Streptococcal Pneumonia treated with Intravenous Immunoglobulins (Ivlg). *Preprint.org* 2020, DOI: 10.20944/preprints202009.0544.v1
8. Iwakura Y, Ishigame H. The IL-23/IL-17 axis in inflammation. *J Clin Invest* 2006; 116:1218-22.
9. Mok MY, Huang FP, Ip WK, Lo Y, Wong FY, Chan EY, Lam KF, Xu D. Serum levels of IL-33 and soluble ST2 and their association with disease activity in systemic lupus erythematosus. *Rheumatology (Oxford)* 2010; 49:520-7.
10. Pons-Estel G; Alarcon, Graciela S; Scofield, Lacie; Cooper, Glinda S. "Understanding the Epidemiology and Progression of Systemic Lupus Erythematosus". *Seminars in Arthritis and Rheumatism* 2010; 39: 257–68.
11. Izati AF, Mohd Shukri ND, Wan Ghazali WS, Che Hussin CM, Wong KK. Increased IL-23R⁺ Th Cells Population Exhibits Higher SLEDAI-2K Scores in Systemic Lupus Erythematosus Patients. *Front Immunol.* 2021;12:690908. Published 2021 Aug 17. doi:10.3389/fimmu.2021.690908
12. Dai H, He F, Tsokos GC, Kyttaris VC. IL-23 Limits the Production of IL-2 and Promotes Autoimmunity in Lupus. *J Immunol.* 2017 Aug 1;199(3):903-910. doi: 10.4049/jimmunol.1700418.




Figure 1 shows the levels of IL-17, IL-23, and IL-33 in patients with and without SLE. The levels of cytokines were significantly higher for IL-23 than for the other ILs. IL-17 levels were significantly higher than in the control group.

Table 2. Level of statistical significance of clinical manifestations based on the Systemic lupus erythematosus (SLE) disease activity index (SLEDAI) 2000 score.

Clinical Manifestation	Percentage	P-value
Arthritis	89%	0.000
Visual disturbances	35%	0.003
Headache	68%	0.000
Vasculitis	32%	0.005
Proteinuria	50%	0.000
Haematuria	30%	0.007
Fever	72%	0.000
Myositis	77%	0.000
Pleurisy	45%	0.001
Low complement levels	60%	0.000
Leukopenia	35%	0.003

Table 1 shows SLE Disease Activity Index (SLEDAI) 2000 score depicted that 89% of the SLE patients had lupus in activity, and therefore required treatment with immunosuppressors right away. All variables measured are statistically significant, $p < 0.05$.