

## Effect of Iron Nutrition on Growth, Quality and Yield of Soybean (*Glycine Max. L.*) Grown on Problematic Inceptisol.

### ABSTRACT

**Aims:** To study the effect of soil and foliar application of iron on growth parameters, **quality** and **yield** of soybean (*Glycine max.L.*).

**Study design:** The experiment was laid out in **randomised** block design with three replications.

**Place and duration of the study:** Agricultural Research Station, Kasbe Digraj, Dist: Sangli (MS) India. The study was conducted during **Kharif** 2018-19.

**Methodology applied:** **The present investigation consisted of eight treatments** such as absolute control, general recommended dose of fertilizer (GRDF), GRDF + soil application of  $\text{FeSO}_4$  @ 10 and 20  $\text{kg ha}^{-1}$  with and without two foliar sprays of chelated Fe @ 0.2 per cent at 30 and 50 days after sowing (DAS) , GRDF + soil application of  $\text{FeSO}_4$  @ 20  $\text{kg ha}^{-1}$ , GRDF +  $\text{FeSO}_4$  @ 10  $\text{kg ha}^{-1}$  + cow dung slurry @ 500 litre  $\text{ha}^{-1}$ .

**Results.** The basal application of  $\text{FeSO}_4$  @ 10 or 20  $\text{kg ha}^{-1}$  along with two foliar sprays of chelated Fe @ 0.2 per cent at 30 and 50 days after sowing improved growth parameters of soybean viz; chlorophyll content, average number of days to 50 per cent flowering, plant height, number of branches, number of pods and effective root nodules as compare to only soil application or foliar spray of iron. Significantly higher grain yield, straw **yield**, **100 grain weight**, **oil** and crude protein yield of soybean were obtained in **soil application** of  $\text{FeSO}_4$  @ 20  $\text{kg ha}^{-1}$  along with two foliar sprays of chelated Fe @ 0.2% at 30 and 50 days after sowing.

**Conclusion:** Integrated approach of iron nutrition in soybean both as soil and foliar application could result in better growth and **the** enhanced yield of soybean.

**Keywords:** Iron Growth, Quality, Soil and Foliar **fertilisation**, Soybean, Yield.

### 1.INTRODUCTION

Soybean (*Glycine max L.*) is a leguminous crop and it belongs to family papillionaceae, sub family of leguminoaceae, originally a crop of China. Soybean **has been** cultivated for more than 3000 years in South-Eastern Asia **Dwevedi and Kayastha** (2011) [1].

Soybean is the world's most popular edible oil, and it plays a vital role in the economy. Legumes serve an important part in human nutrition since they are high in protein, calories, and certain minerals and

vitamins. Soybean is the most abundant source of protein and vegetable oil containing polyunsaturated fatty acids, particularly Omega 6 and Omega 3. [2] Chauhan et al. (1988).

Soybeans are grown on a total of 124 million hectares around the world. After the United States, Brazil, China, and Argentina, India comes fifth in terms of area and output. All world estimated area and production of soybean in *Kharif*- 2017 was 10.60 million ha and 8.00 million MT respectively APE (2017) [3]. Due to factors such as soybean's short growing season (90-110 days), good market price, and higher yield than other pulses, the area under soybean production is increasing. It's simple to process for items like soy cheese, soy milk, soy protein, soy yoghurt, soybean oil, and soy nuts. Soybean is also used to make soy ink, soy paint, and soy molasses. It is a potential crop that can boost the food-processing industry in rural areas. Soybean production is affected by many factors such as climatic and edaphic factors, which severely affect production; According to Turner 1991, the performance of this crop is highly influenced by the availability of trace elements such as Molybdenum and Iron. Besides, iron deficiency of Mn and Zn can also affect the production of soybean crops Zayed, et al. (2017) [4]. Deficiency of micronutrients and low availability of other essential nutrients or imbalance use of fertilisers emerged as the important constraint in soybean production. Hence a balanced nutrient application is must to increase the productivity of the soybean crop. Among micronutrients, iron plays vital role in a structural component of porphyrin molecules, cytochromes, hemes, hematin, ferrichrome and leghaemoglobin. These substances are involved in oxidation-reduction reactions in respiration and photosynthesis. It's also found in enzymes like amino levulinic acid synthetase and co-protoporphyrin ogenoxidase, which are required for nitrogen fixation in nitrogen-fixing bacteria. The existence of cytochromes for executing various photosynthetic reduction activities, as well as ferredoxin as an electron acceptor, is reflected in the abundance of iron in chloroplasts. Ferredoxins are Fe-S proteins that are the photosynthetic electron transport chain's first stable redox molecule. Iron deficiency is usually observed in soybean grown in calcareous or alkaline soils. In calcareous soil, iron availability is restricted due to conversion of ferrous to ferric and showed deficiency of Fe manifest into yellowish inter-venal paling of younger leaves (commonly referred to as iron chlorosis) and soil conditions such as high soil pH found in large areas of the Great Plains may decrease the plant availability of some macro and micronutrients. This may be corrected through initially application at the time of sowing and foliar fertiliser application of a combination of starter and booster dose of fertiliser. Supplementary foliar application of N, P, K and micronutrients for deficient soils can help to enhance the crop yields under these conditions. Foliar application of micronutrients in high pH saline soils is more beneficial in terms of growth and yield of the crop Zayed, et al. (2017) [4]. Foliar application of micronutrients is more beneficial as compared to soil application as the application rate of the nutrient is comparatively lesser, nutrient absorption is more moreover, when roots cannot provide necessary nutrients, foliar application is always a compatible alternative Hanwate et al.(2018) [5]. There is an increasing interest from producers about the potential benefits of foliar application of nutrients as a compliment of their fertilisation programs to maximise yields. In view of this, the present investigation was undertaken to study the effect of soil and foliar application of iron on nutrient availability, uptake, yield and quality of soybean (*Glycine max.* L.).

## 2. MATERIAL AND METHODS

### 3.1 Experimental site and soil

The field experiment was conducted at Agricultural Research Station, Kasbe Digraj, Dist: Sangli Maharashtra, India during *kharif* season of the year 2018. This study area is located in Deccan plateau, hot semi-arid eco region, in the Western Maharashtra plane zone (Zone VI) and is situated at 16°08' North latitude, 74°08' East longitude and at an altitude of 580 m above mean sea level (MSL). The experimental soil (0-15 cm soil depth) had alkaline pH, electrical conductivity (EC) 0.18 dS m<sup>-1</sup>, calcium carbonate (CaCO<sub>3</sub>) 6.80 g kg<sup>-1</sup>, clayey in texture, bulk density (BD) 1.25 Mg m<sup>-3</sup> and organic carbon 4.50 g kg<sup>-1</sup>. The soil available nitrogen, and potassium contents were 170, 7.50, 433 kg ha<sup>-1</sup> respectively, and soil DTPA iron, zinc, copper and manganese contents were 4.05, 0.35, 0.40 and 2.52 ppm respectively.

### 3.2 Experimental details

The experiment was laid out in **randomised** block design with eight treatments and three replications in *kharif*, 2018. The treatments were absolute control (T<sub>1</sub>), general recommended dose of fertilizer (GRDF) *i.e.* 50:75:45 kg ha<sup>-1</sup> N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O + 10 t ha<sup>-1</sup> FYM (T<sub>2</sub>), GRDF + soil application of FeSO<sub>4</sub> @ 10 kg ha<sup>-1</sup> (T<sub>3</sub>), GRDF + soil application of FeSO<sub>4</sub> @ 20 kg ha<sup>-1</sup> (T<sub>4</sub>), GRDF + FeSO<sub>4</sub> @ 10 kg ha<sup>-1</sup> + cow dung slurry @ 500 liters ha<sup>-1</sup> (T<sub>5</sub>), GRDF + two foliar sprays of chelated Fe @ 0.2% at 30 and 50 days after sowing (DAS) (T<sub>6</sub>), GRDF + soil application of FeSO<sub>4</sub> @ 10 kg ha<sup>-1</sup> + two foliar sprays of chelated Fe @ 0.2% at 30 and 50 DAS (T<sub>7</sub>) and GRDF + soil application of FeSO<sub>4</sub> @ 20 kg ha<sup>-1</sup> + two foliar sprays of chelated Fe @ 0.2% at 30 and 50 DAS (T<sub>8</sub>). The soybean crop was **fertilised** with 50 kg N, 75 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup> and 45 K<sub>2</sub>O for treatment GRDF as a basal dose of N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O was applied through urea, single super phosphate and muriate of potash to treatment T<sub>2</sub> to T<sub>8</sub> at the time of sowing. **Ferrous sulfate was incubated with well decomposed FYM for four days and this incubated ferrous sulfate was used for soil application in treatments T<sub>3</sub>, T<sub>4</sub>, T<sub>7</sub> and T<sub>8</sub> at the time of sowing.** The cow dung slurry (125 kg cow dung + 500 liters water) with FeSO<sub>4</sub> @ 10 kg ha<sup>-1</sup> were incubated for one week and applied to the treatment T<sub>5</sub> during first irrigation. The seeds of soybean variety *Phule Sangam* (KDS 726) were inoculated with *Rhizobium* and phosphate **solubilising** bacteria @ 250 g per 10 kg of seeds and used for sowing. The three irrigations were given during the crop growth period. The soybean crop was sown in **kharif** season with 30 cm row spacing.

### 3.3. Growth parameters

**The observations of growth parameters were recorded by randomly selecting five soybean plants from each treatment plot.** The selected plants were marked by fixing pegs. **The days required for 50 per cent flowering were counted since the day after sowing.** The plant **heights** of five plants **was** measured from the ground level up to the growing point of **the** plant and the average height of **the** plant was expressed in centimeters. The number of branches plant<sup>-1</sup> **was** recorded at pod filling stage. Randomly selected five plants used for counting the number of pods plant<sup>-1</sup> at harvest. The pods from each plant were removed, separated, counted and recorded under the respective treatments and then the mean was computed. The observation on total number of root nodules and number of effective and non-effective nodules plant<sup>-1</sup> was recorded by carefully uprooting two randomly selected soybean plants from each treatment.

### 3.4. Chlorophyll content

The chlorophyll content was obtained by 30 DAS and 50 DAS of green plant samples. Chlorophyll of fresh plant leaves (4<sup>th</sup> leaf) at flowering stage **was** extracted in 85 per cent acetone and the absorbance values at 660 nm and 642.5 nm wavelength were recorded on spectro photometer **Arnon (1946)** [6].

### 3.5. Harvesting

The soybean crop was harvested at physiological maturity when the pods turned yellow colour with matured seeds. The border line plants were removed first to eliminate the border effect. The crop from the net plot was cut close to the ground and kept in respective plots for drying. The plot wise threshing of soybean was carried out. The grains were separated from the plant by a mechanical thresher. The straw yield and soybean grain yield were recorded by weighing as per treatments. One hundred seeds were randomly collected from the net plot yield, counted, weighed and expressed as test weight in grams.

### 3.7. Quality parameters:

The protein content in the seeds was analysed by indirect method. First, the per cent nitrogen content of the sample was estimated by microkjeldahl method PJA (1975) [7]. Then the nitrogen value was multiplied by a factor 5.71 to get the protein content of the sample and expressed in percentage FAO (2003) [8]. Oil percentage of grain was determined by Soxhlet extractor using petroleum ether as a solvent.

### 3.8 Statistical Analysis

The experimental data were analysed statistically by applying the "Analysis of variance" technique and significance was tested by variance ratio i.e. F value at 5 per cent level of significance SMA (1985) [9]. Standard error of the mean (S.Em.) and critical difference (CD) was worked out to evaluate differences between treatment means.

## 3 Results and Discussion

### 3.1 Chlorophyll content at 30 and 50 DAS

Chlorophyll content in leaves differed significantly due to soil application of iron and foliar sprays treatments. The soil application of  $\text{FeSO}_4$  @ 20 kg ha<sup>-1</sup> along with two foliar sprays of chelated Fe @ 0.2% (T<sub>8</sub>) had recorded the highest chlorophyll content significantly at 30 and 50 DAS (20.28 mg g<sup>-1</sup>) and (21.95 mg g<sup>-1</sup>) respectively, over the rest of treatments however, it was at par with soil application  $\text{FeSO}_4$  @ 10 kg ha<sup>-1</sup> along with two foliar sprays of chelated Fe @ 0.2% at 30 DAS (19.49 mg g<sup>-1</sup>) and 50 DAS (21.15 mg g<sup>-1</sup>) and soil application of  $\text{FeSO}_4$  @ 20 kg ha<sup>-1</sup>. The soil application of  $\text{FeSO}_4$  @ 10 or 20 kg ha<sup>-1</sup> and foliar application of chelated Fe 0.2% exhibited higher chlorophyll content at 30 and 50 DAS as compare to GRDF (T<sub>2</sub>) and absolute control (T<sub>1</sub>). This might be due to the beneficial effect of  $\text{FeSO}_4$  application to soil along with foliar spray increased iron availability in soil, and ferrous iron (Fe<sup>2+</sup>) uptake by plant leaves in foliar resulting in better absorption and translocation of iron. Which, in turn, might have helped the cellular activity and also directly or indirectly participated in the formation of chlorophyll and thus increasing photosynthesis. The results are in agreement with that of Kandoliya *et al.* (2018) [10], who reported an increase in total chlorophyll content due to iron nutrition in wheat crops.

### 3.2 Days to 50% flowering

Days to 50 per cent flowering also differed significantly due to different treatment application of iron through the soil and foliar. The significantly higher average number of days to 50 per cent flowering (44.33) was noticed in soil application of  $\text{FeSO}_4$  @ 20 kg ha<sup>-1</sup> along with two foliar sprays of chelated Fe @ 0.2% at 30 and 50 DAS (44.33) over the rest of the other treatments. The T<sub>8</sub> was at par with T<sub>7</sub> at 30 and 50 DAS (43.66). The treatments T<sub>2</sub> to T<sub>7</sub> were at par with each other for average number of days to 50 per cent flowering. The application of Fe fertiliser with FYM took longer period for average number of days to 50 per cent flowering. This might be due to the  $\text{FeSO}_4$  along with FYM treatments enhances the growth by mineralisation and availability of essential nutrients to soybean and increases the uptake of

nutrients and more vegetative growth of soybean. The findings of Maheswari and Karthik (2017) [11] also confirmed the results of present study.

### 3.3 Plant height

The plant height of the soybean crop was significantly influenced by iron application through the soil and foliar treatments. The treatment T<sub>2</sub> recorded significantly higher plant height as compared to control. The application of chemical fertilisers along with FYM resulted in the increase in growth attributes. This may be due to better uptake and translocation of plant nutrients to growing plants and more photosynthesis which in turn promoted more number of leaves, leaf area and dry matter production and found to have a more beneficial effect on plant height of soybean. The soil application of FeSO<sub>4</sub> @ 20 kg ha<sup>-1</sup> along with two foliar sprays of chelated Fe @ 0.2% at 30 and 50 DAS (T<sub>8</sub>) was shown to have a significantly higher plant height (86.33 cm) than the rest of the treatments. The treatment T<sub>8</sub> was on par with T<sub>7</sub> for plant height (85.00 cm), T<sub>4</sub> (83.33 cm) and T<sub>6</sub> (84.00 cm). It clearly indicated that GRDF along with soil application of FeSO<sub>4</sub> @ 10 and 20 kg ha<sup>-1</sup> with two sprays of chelated Fe 0.2 % treatments were found higher plant height. The application of Fe resulted in the increase in chlorophyll content and more photosynthesis also FYM had more beneficial and significant effect on plant height. Similar results were also reported by Balachander *et al.* (2003) [12].

### 3.4 Number of branches per plant

The number of branches per plant of soybean was influenced by various the treatments and was found to be statistically significant. The significant highest number of branches (11.0 plant<sup>-1</sup>) was recorded in soil application of FeSO<sub>4</sub> @ 20 kg ha<sup>-1</sup> along with two foliar sprays of chelated Fe @ 0.2% over the rest of the treatments. The treatments T<sub>8</sub>, T<sub>7</sub>, T<sub>4</sub>, T<sub>6</sub> and T<sub>5</sub> were at par with each other in terms of number of branches per plant. The higher number of branches in soybean exhibited by treatment receiving soil application of FeSO<sub>4</sub> @ 10 or 20 kg ha<sup>-1</sup> and two sprays of chelated Fe @ 0.2% might be due to the combined application of Fe with recommended dose of other major nutrients which in turn might have increased uptake of nutrients in the plants leading to enhanced chlorophyll content and carbohydrate synthesis and thus helped in increased number of branches per plant of soybean. Similar results were also noticed by Kumar *et al.* (2013) [13] and Kunjammal and Sukumar (2019) [14].

### 3.5 Number of pods per plant

The number of pods per plant differed significantly due to iron application through soil and foliar sprays. The soil application of FeSO<sub>4</sub> @ 20 kg ha<sup>-1</sup> along with two foliar sprays of chelated Fe @ 0.2% (T<sub>8</sub>) obtained the significantly more number of pods (44.0 plant<sup>-1</sup>) over the rest of treatments. The treatment T<sub>8</sub> was at par with T<sub>7</sub> (43.33 plant<sup>-1</sup>) and T<sub>4</sub> (41.33 plant<sup>-1</sup>) for number of pods. The highest pods plant<sup>-1</sup> due to the foliar application could be attributed to iron's significant effect on reproductive organs, such as stamens and pollens Siefi *et al.* (2011) [15]. The increase in the number of soybean pods confirmed the translocation of photosynthates to the productive sink. Application of Fe increases the number of pods per plant in mothbean Sachendra *et al.* (2006)[16].

### 3.6 Root nodules per plant

A significantly more number of effective root nodules (44.33) were found in soil application of FeSO<sub>4</sub> @ 20 kg ha<sup>-1</sup> along with two foliar sprays of chelated Fe @ 0.2% treatment than rest of the treatments. Iron is required for the synthesis of Fe containing proteins in the host, including leghaemoglobin and in bacteroids for nitrogenase and cytochromes of electron transport chain. Similar results were also observed by Shukla and Shukla (1994) [17]. The soil application of FeSO<sub>4</sub> @ 20 kg ha<sup>-1</sup> along with two

foliar sprays of chelated Fe @ 0.2% was found to have non-significant effect with soil application  $\text{FeSO}_4$  @ 10  $\text{kg ha}^{-1}$  along with two foliar sprays of chelated Fe @ 0.2% at 30 and 50 DAS ( $T_7$ ) and soil application of  $\text{FeSO}_4$  @ 20  $\text{kg ha}^{-1}$  ( $T_4$ ). It indicated that basal application  $\text{FeSO}_4$  @ 10 or 20  $\text{kg ha}^{-1}$  were increased the nodule count. The soil application of  $\text{FeSO}_4$  @ 10 or 20  $\text{kg ha}^{-1}$  and foliar application of chelated Fe 0.2% at 30 and 50 DAS treatments were found to have a higher number of effective root nodules at 30 and 50 DAS as compared to GRDF ( $T_2$ ) and absolute control ( $T_1$ ). The soil application of ferrous sulphate at 25  $\text{kg ha}^{-1}$  to soybean crop increased nodulation, and nodules dry weight per plant as compared to control Bhanavase (1994) [18]. The reverse trend was observed in respect of non-effective root nodules per plant of soybean.

### 3.7 100 grain weight

The findings regarding 100 grain weight revealed that the treatments receiving iron nutrition either through soil, foliar or combination of both recorded higher grain weight. Soil application of  $\text{FeSO}_4$  @ 20  $\text{kg ha}^{-1}$  along with two foliar sprays of chelated Fe @ 0.2% showed significantly higher 100 grain weight (19.20 g) over GRDF ( $T_2$ ) and control ( $T_1$ ) however, this value was at par with all the other treatments of iron nutrition. The reason for the increased of the 100 grain weight could be attributed to enhanced photosynthetic activity due to increased chlorophyll content in leaves due to iron application. This might have resulted in the production and accumulation of carbohydrates in 100 grain weight development. Similar results were also recorded by Mohammad *et al.* (2012) [19] foliar spray of micronutrient had a significant effect on 1000 seed weight safflower. Rubens *et al.* (2019) [20] reported an increased in soybean grain weight due to micronutrient fertilisation.

### 3.8 Grain yield

The significantly higher grain yield (24.93  $\text{q ha}^{-1}$ ) was observed with treatment receiving soil application of  $\text{FeSO}_4$  @ 20  $\text{kg ha}^{-1}$  along with two foliar sprays of chelated Fe @ 0.2% ( $T_8$ ) over the rest of the treatments which was estimated to be 14 per cent higher compared to  $T_2$  and 81 per cent over  $T_1$ . The results exhibited that treatments receiving iron nutrition irrespective of method used for iron nutrition demonstrated increment in soybean grain yield as compared treatments without iron supplement. This might be due to rapid availability of iron to plants due to foliar application of chelated Fe and soil applied  $\text{FeSO}_4$ . Moreover, application of FYM also might have resulted increased concentration of plant available iron and formation of metalo-organic complexes of higher extractability which in turn might have resulted in continuous supply of iron. The enhanced availability of iron could have increased chlorophyll content and accumulation more carbohydrates which is associated with increase in flowering and pod development ultimately increased grain yield of soybean. While foliar application of iron might have resulted in direct absorption of the foliage sprayed with Fe solution. The results conform to that of Sale *et al.* (2017) [21] who observed increased soybean yields due to foliar nutrition of Fe and Zn. Similarly, Moosavi and Ronaghi (2011) [22] also reported a substantial increase in soybean yield in response to foliar and soil iron nutrition.

### 3.9 Oil content and oil yield

The oil content in seed was not influenced significantly due to various treatments. Nevertheless, treatment receiving iron fertilisation along with major nutrients revealed slight increase in oil content. Significantly difference was noticed in the case of oil yield, application of  $\text{FeSO}_4$  @ 20  $\text{kg ha}^{-1}$  along with two foliar sprays of chelated Fe @ 0.2% ( $T_8$ ) recorded the significantly higher oil content (19.46%) and oil yield (485  $\text{kg ha}^{-1}$ ) over the rest of treatments. The soil application of  $\text{FeSO}_4$  @ 20  $\text{kg ha}^{-1}$  along with two

foliar sprays of chelated Fe @ 0.2% was at par with soil application of  $\text{FeSO}_4$  @  $10 \text{ kg ha}^{-1}$  along with two foliar sprays of chelated Fe @ 0.2% at 30 and 50 DAS for oil yield ( $T_7$   $446 \text{ kg ha}^{-1}$ ) and soil application of  $\text{FeSO}_4$  @  $20 \text{ kg ha}^{-1}$  ( $T_4$   $444 \text{ kg ha}^{-1}$ ). The soil application of  $\text{FeSO}_4$  @  $10$  or  $20 \text{ kg ha}^{-1}$  and foliar application of chelated Fe 0.2% at 30 and 50 DAS and GRDF treatments recorded higher oil yield over the control. It indicated that the application of iron through the soil and foliar was beneficial for increasing soybean oil yield. Ferrous sulphate also contains sulphur in addition to iron. Sulphur is one of the important secondary nutrients required by the crops, sulphur and iron might have helped to obtain a higher oil yield of soybean. Higher oil yield may be due to higher iron availability in alkaline soils, which ensured better biosynthesis of oil in groundnut Poonia *et al.* (2018) [23].

### 3.10 Crude protein content and protein yield

Protein content in seed and protein yield was significantly influenced by various treatments. The application of Fe either through soil as well as foliar sprays recorded higher values of crude protein content and protein yield of soybean as compare to control. Significantly lower crude protein (27.16%) and crude protein yield ( $372 \text{ kg ha}^{-1}$ ) were recorded in without FYM and fertilisers over the rest of treatments. However, the significantly higher crude protein (32.24%) and crude protein yield ( $807 \text{ kg ha}^{-1}$ ) were noticed in treatment soil application of  $\text{FeSO}_4$  @  $20 \text{ kg ha}^{-1}$  along with two foliar sprays of chelated Fe @ 0.2% as compare to rest of treatments. The treatments  $T_2$  to  $T_7$  were at par for crude protein content in soybean. The treatment  $T_8$  was at par with  $T_7$  ( $746 \text{ kg ha}^{-1}$ ) and  $T_4$  ( $746 \text{ kg ha}^{-1}$ ) for crude protein yield of soybean. It clearly indicated that application of iron is beneficial for increasing crude protein content and crude protein yield of soybean. This might be because iron is essential for nitrogen fixation and better availability of nitrogen and it absorption ultimately increases in protein content in grain of soybean. Similar results were close in conformity with Sale *et al.* [21].

## 4. Conclusions

The application of  $50:75:45 \text{ N:P}_2\text{O}_5:\text{K}_2\text{O} \text{ kg ha}^{-1} + 10 \text{ t FYM ha}^{-1}$  and soil application of  $\text{FeSO}_4$  @  $20 \text{ kg ha}^{-1}$  alongwith two foliar sprays of chelated Fe @ 0.2% at 30 and 50 DAS to soybean in iron deficient soil recorded higher growth parameters, grain and straw yield, quality parameters, nutrient uptake by soybean. The residual soil fertility was improved in treatments received GRDF and Fe as compared to initial soil fertility status.

## 5. REFERENCES

1. Dwevedi A. Kayastha AM. Soybean a multifaceted legume with enormous economic capabilities. Soybean-Biochemistry, Chemistry and Physiology, In Tech Europe University Campus, Croatia, 2011.
2. Chauhan GS Verma NS Basin GS. Effect of extrusion processing on the nutritional quality of protein in rice legume blends. *Nahrung* 1988; 32: 43-47.
3. Area production estimates of soybean in India Kharif. [sopa@sopa.org](mailto:sopa@sopa.org),2017.
4. Zayed BA Salem AKM El Sharkawy HM. Effect of different micronutrient treatments on rice (*oryza sativa* L.) growth and yield under saline soil conditions. *World J. Agric. Sci.* 2011; 7(2): 179-184.
5. Hanwate GR Giri SN Yelvikar NV. Effect of foliar application of micronutrients on nutrient uptake by soybean crop. *Inter. J. of Pure Application Biosci.* 2018; 6: 261-265.

6. Arnon DI. Copper enzymes in isolated chloroplasts polyphenol oxidase in (*Beta vulgaris*). *J. of Plant Physiol.*1949; 24: 1-15.
7. Parkinson JA Allen SE. A wet oxidation procedure suitable for the determination of nitrogen and other mineral nutrients in biological material. *Commun in Soil Sci and Plant Analy.*1975; 6: 7-11.
8. Food and Agriculture Organization of the United Nations *Viale delle Terme di* ISBN 92-5-104986-6. Printed in Italy 2003.
9. *Statistical method of Agricultural workers*, ed Panse VG Sukhatme PV publ.ICAR, New Delhi 1985.
10. Kandoliya RU Kunjadia BB.Effect soil and foliar application of zinc and iron on micronutrients uptake by wheat in calcareous soil of Saurashtra region. *European J. of Biotech and Biosci.* 2018; 6: 65-69.
11. Uma Maheswari MA Karthik Ajay Kumar R. Effect of foliar nutrition on growth, yield attributes and seed yield of pulse crops. *Int.J.Curr.Microbiol.App.Sci.* 2017; 6(11): 4134-4139. doi: <https://doi.org/10.20546/ijcmas.2017.611.484>
12. Balachandar D Nagarajan P Gunasekaran S. Effect of organic amendments and micronutrients on nodulation and yield of black gram in acid soil. *Leg Research J.* 2003; 26: 192-195.
13. Kumar V Dwivedi VN Tiwari DD. Effect of phosphorus and iron on yield and mineral nutrition in chickpea. *Annals Plant Soil Res.*2013; 11: 16-18.
14. Kunjammal P Sukumar J. Effect of different seed treatment on grain yield of **maize** (*Zea mays*.) under drought stress conditions. *Madras Agric J.*2019; 106(4–6): 384–387.
15. Seifi-Nadergholi M Yarnia M Rahimzade KF. Effect of Fe and Mn and their application method on yield and yield components of common bean (*Phaseolus vulgaris*). *Middle-East J of Sci Res.* 2011; 8: 859-865.
16. Sachendra BNM Singh J Bohra A Viyas A. Change in morphological and biochemical characteristics of Mothbean in Indian thar desert due to sulphur and iron nutrition. *America-Eurasian J Agric and Environ Sci.* 2006;1: 51-57.
17. Shukla V Shukla IC. Effect of Fe, Mo, Zn and P on symbiotic nitrogen fixation of chickpea. *Indian J Agric Chem.*1994; 32: 118-123.
18. Bhanavase DB Jadhav BR Kshirsagar CR Patil PL. Studies on chlorophyll, nodulation, N-fixation, soybean yield and their correlation as influenced by micronutrients. *Madras Agric J.*1994; 8:325-328.
19. Mohammad G Mahoud R Tavassoli A. Effect of micronutrients foliar application on yield and seed oil content of safflower (*Carthamus tinctorius* L.). *African J Agric Res.* 2012; 7: 482-486.
20. Rubens RS Larissa UR Rodrigo RF Álvaro JG Faria Vitor Nascimento L. Nutritional and morphophysiological responses of soybean to micronutrient fertilisation in soil. *Commun in Plant Sci.* 2019; 9: 93-99.
21. Sale RB Nazirkar RB Ritu ST Nilam BK. Effect of foliar spray of zinc, iron and seed priming with molybdenum on growth and yield attributes and quality of soybean in the rainfed condition of Vertisol. *Intern J. Chem Stud.* 2017;6: 828-831.

22. Moosavi AA Ronaghi A. Influence of foliar and soil applications of iron and manganese on soybean dry matter yield and iron-manganese relationship in a calcareous soil. *Australian J Crop Sci.* 2011; 5: 1550-1556.
23. Poonia T Bhunia SR Choudhary R. Effect of fertilisation on nitrogen and iron content, uptake and quality parameters of groundnut (*Arachis hypogaea* L.). *Internatl J Current Micro and Appl Sci.* 2018;7: 2297-2303.

**Table 1 Effect of soil and foliar application of iron on chlorophyll content and number of days to 50 percent flowering of soybean**

Treatments	Chlorophyll content (mg g <sup>-1</sup> )		Days to 50 percent flowering
	30 DAS	50 DAS	

T <sub>1</sub>	16.11	16.45	38.68
T <sub>2</sub>	17.25	17.26	42.00
T <sub>3</sub>	17.62	18.02	42.68
T <sub>4</sub>	18.21	18.88	43.00
T <sub>5</sub>	17.76	18.26	42.00
T <sub>6</sub>	16.92	20.05	41.66
T <sub>7</sub>	19.49	21.15	43.66
T <sub>8</sub>	20.28	21.95	44.33
SE <sub>±</sub>	0.71	0.59	0.54
CD at 5 %	2.17	1.81	1.66

**Table 2 Effect of soil and foliar application of iron on growth parameters of soybean**

Treatments	Plant height at flowering stage (cm)	Number of branches plant <sup>-1</sup> at	Number of pods plant <sup>-1</sup> at harvest	Number of root nodules plant <sup>-1</sup> at flowering	
				Effective	Non-effective

	<b>flowering stage</b>				
T <sub>1</sub>	75.00	8.00	29.33	31.33	16.33
T <sub>2</sub>	80.00	8.66	37.66	36.66	15.00
T <sub>3</sub>	81.00	9.33	38.66	38.00	13.00
T <sub>4</sub>	83.33	10.00	41.33	40.66	12.33
T <sub>5</sub>	81.66	9.66	40.66	38.33	11.33
T <sub>6</sub>	84.00	10.00	40.66	39.33	12.33
T <sub>7</sub>	85.00	10.66	43.33	42.33	7.00
T <sub>8</sub>	86.33	11.00	44.00	44.33	8.00
SE±	0.97	0.54	0.93	1.50	0.82
CD at 5 %	2.95	1.66	2.85	4.55	2.50

**Table 3** Effect of soil and foliar application of iron on grain, straw yield and yield contributing parameter of soybean

Treatments	Grain yield (q ha <sup>-1</sup> )	Straw yield (q ha <sup>-1</sup> )	100 grain weight (g)
T <sub>1</sub>	13.72	20.70	17.1
T <sub>2</sub>	21.80	33.05	17.7
T <sub>3</sub>	22.12	34.79	18.2
T <sub>4</sub>	23.14	36.33	18.9
T <sub>5</sub>	22.37	35.28	18.7
T <sub>6</sub>	22.19	35.20	18.3
T <sub>7</sub>	23.24	36.38	19.0
T <sub>8</sub>	24.93	37.79	19.2
SE <sub>±</sub>	0.70	1.52	0.40
CD at 5 %	2.12	4.62	1.23

**Table 4 Effect of soil and foliar application of iron on quality parameters of soybean**

<b>Treatments</b>	<b>Oil (%)</b>	<b>Oil yield (kg ha<sup>-1</sup>)</b>	<b>Crude protein (%)</b>	<b>Crude protein yield (kg ha<sup>-1</sup>)</b>
T <sub>1</sub>	18.40	252	27.16	372
T <sub>2</sub>	18.83	410	30.78	671
T <sub>3</sub>	19.00	420	30.88	684
T <sub>4</sub>	19.20	444	32.31	746
T <sub>5</sub>	19.40	434	31.38	705
T <sub>6</sub>	19.06	422	31.91	707
T <sub>7</sub>	19.20	446	32.17	746
T <sub>8</sub>	19.46	485	32.24	807
SE <sub>±</sub>	0.28	15.89	0.80	31.30
CD at 5 %	NS	48.20	2.43	94.94