

Case report

Case Report on Alcohol Dependence with Delirium Tremors

Abstract

Alcohol is a natural product made by reacting lactose fermentation with yeast spores. Because it includes calories. With no nutritional value, alcohol is used as a source of energy. Different sugar sources are used for fermentation to create a variety of alcoholic beverages. Patient Present History:-On September 3rd, 2020, a 32-year-old male patient was admitted to hospital with a specific complaint of abdominal pain for the past three years. He began drinking alcohol on a daily basis. Initially, he only drank at night, but later he began drinking during the day. **Clinical finding:** - Mood irritable, social withdrawal, mental confusion, seizure, vomiting, mood changes, lethargy and vomiting. **Diagnostic Evaluation:** - Hb - 11.9mg%, MCV - 87cum/micron, MCH- 28.9pico/gm, Total Bilirubin- 1.47.USG –He is appearance was natural once more. There was no localized lesion or oblivious mass lesion in the pelvis.

Keywords:-Alcohol Dependence Syndrome, delirium tremor, alcoholic beverages, social withdrawal

Introduction

Alcohol is a natural product made by reacting lactose fermentation with yeast spores. Because it includes calories with no nutritional value, alcohol is used as a source of energy. Different sugar sources are used for fermentation to create a variety of alcoholic beverages.¹ History of the Patient - On May 5, 2021, a 32-year-old male patient was admitted in hospital with the primary complaint of. Since the age of two. A previous psychiatric illness (DSM - IV and ICD - 10) in which a person is physically or mentally reliant on alcohol is known as alcohol dependency.² A chronic condition in which a person desires alcoholic beverages and is unable to control his or her consumption. A person with this disease often has to drink higher doses to achieve the name effect and may have symptoms.³

Patient Present History :-

On September 3rd, 2020, a 32-year-old male patient was admitted to hospital with a specific complaint of abdominal pain for the past three years. He began drinking alcohol on a daily basis. Initially, he only drank at night, but later he began drinking during the day. On a daily basis, he consumes 2 to 3 quarters. Eye opener dinking and abdominal discomfort are also reported by the patient. Intermianally he is in excruciating discomfort.

Content	Results
Mood and affect is impaired	Unstable and irritable mood
Content of thought is impaired	Delusions of persecution is present
Attention is intact	he count the number backwards
Concentration is intact	correct result
Judgment is intact	he giving correct answer

Table 1. Mental Status Examinations

Pathology	Results
Hb %	11.9mg %
MCV	87Cum / micron
MCH	28.9Pico / gm
Total Bilirubin	1.47
Table 2. Blood Investigation Report	

USG - His appearance was natural once more. There was no localised lesion or obvious mass lesion in the pelvis. The first impression is mild hepatomegaly. A thorough examination of the patient's background and mental state indicates that he or she is truly suffering from a racial oppression hallucination. The first impression is mild hepatomegaly. A thorough examination of the patient's background and mental state indicates that he or she is truly suffering from a racial oppression hallucination.

Therapeutic intervention -

Psychopharmacology

Antipsychotic and anticraving medications were given to the patient. Tab Neurobionfort OD, inj. Lorazepam 2 mg HS, inj. Thiamine 200mg IM. Treatment for a short period of time acute psychotic disorders necessitates prompt hospitalization, whether in a mental facility or a crisis line. These syndromes are used to treat a variety of medical conditions. The attending physician makes the choice to admit a patient to the hospital.

Continuation Treatment-

After a specific period of time has elapsed, the effectiveness of anti-craving medicine for psycho-pharmacotherapy is usually evident. Improved sleep, agitation regression, anxiety and paranoia reintegration, and the removal of psychotic symptoms were all seen in the first six weeks. If the patient's condition does not improve, another antipsychotic medication will be prescribed, or the first dose will be raised. A worsening of symptoms, severe side effects, or an insufficient pharmaceutical response are all common reasons for cognitive treatment. Treatment with opioids, mood stabilisers (lithium or valproate), or an anticonvulsant medicine (carbamazepine)⁴ may be prescribed if mood disorders or cyclic breakouts emerge. The distinction between post-narcoleptic depression and the development of an affective illness is crucial (schizoids). Prevention of recurrence Consider the likelihood that symptoms will resurface during the first two years of starting treatment, which should be continued for another one to two years after recovery. Periodic checkups and effective therapeutic therapy, including social and psychological counselling, are necessary during this long-term follow-up.

Nursing management:-

Nursing care was provided according to the requirements in order to maintain personal hygiene and avoid further complications. The patient, as well as his relatives, should be elicited as much information as feasible. The behavior pattern included posturing, psychomotor activity, and hygiene. Check to make sure the person is not hurt. Take notice of

the patient's emotional state and affect to see if it's appropriate. Assessed to discover if his thoughts were delusory. Determine the severity of the patient's hand tremor and suggest upper-extremity training.

Follow-up and Results:

At the time of release, the patient's condition was satisfactory. His main complaint when he was admitted to the hospital was hand tremor, abdominal pain, nausea, vomiting, and headache. His condition has improved since he received treatment. Their family was informed about the medication regimen and personal hygiene instructions, and they were told to return in 10 days for a routine examination to see how the sickness was progressing.

Discussion

Syndrome of Alcohol Dependence The Edwards and Gross alcohol - dependence syndrome (ADS) formulations from 1976 are included in the main classification systems for alcohol - use disorder (DSM - IV and ICD - 10). However, the criteria for distinguishing alcohol dependence (AD) from substance dependency (AA) in DSM - IV and hazardous use in ICD - 10 continue to be problematic. Furthermore, these worries stem from a shift in historical perspectives and problems associated with alcohol intake. Alcohol use disorders (AUDs) are characterised in terms of their historical evolution; improvements in the recognition of drug issues including alcohol.⁵

The amount of alcohol that can be detected is different. The degree to which alcohol is linked to such events varies.⁶ The Stroke fraction is influenced by alcohol, coronary heart disease, liver cirrhosis, and a variety of cancers.⁷ The amount of alcohol that causes various disorders varies.⁸ Alcohol consumption is the fourth leading cause of preventable death in the United States (after smoking, high blood pressure and obesity). Hazardous alcohol use killed nearly three million people in 2016, accounting for 5.3 percent of all deaths, according to a 2018 WHO estimate.⁹ Men are responsible for the majority of such deaths over the world. Ghogare had conducted study on the effects of alcoholism on the brain. In a study, Joshi et al compared total self-stigma in patients with schizophrenia and alcoholism. Kelkar and Parisha studied the prevalence of substance abuse among schizophrenia patients.¹⁰ Patel et al looked on the frequency of alcohol-related disorders in hospitalised male patients. Clinical comorbidity, severity, and liver enzyme impairment in alcohol-dependent patients were reported by Doctors.¹¹ 13 In rural Central India, Jonas et al. investigated the prevalence of depression, suicidal ideation¹³ and alcohol and nicotine consumption.

Strength

A 24-year-old male patient tolerates all medications well and responds effectively to the hospital's therapeutic therapy within seven days of receiving it.

Informed Consent

Patients and their relatives were notified before this case was adopted, and informed consent was obtained from both the patient and the relatives.

Conclusions:

Patient The patient was admitted to the hospital for 10 days with abdominal pain, nausea, and vomiting as the chief diagnosis. Tremor might also be felt in both hands. A member of the health team had begun rapid therapy now that the patient's condition had improved.

REFERENCES

1. Behere, P.B., Behere, A.P., Chowdhury, D., 2020a. Rise in cases of alcohol withdrawal following lockdown in dry district of wardha. *Journal of Neurosciences in Rural Practice* 11, 478–480. <https://doi.org/10.1055/s-0040-1712554>
2. Modi, L., Gedam, S.R., Shivji, I.A., Babar, V., Patil, P.S., 2018. Comparison of total self-stigma between schizophrenia and alcohol dependence patients. *International Journal of High Risk Behaviors and Addiction* 7. <https://doi.org/10.5812/ijhrba.61043>
3. Patel, T.V., Brahmabhatt, M.J., Vankar, G.K., 2018. Prevalence of alcohol use disorders in hospitalised male patients. *Archives of Psychiatry and Psychotherapy* 20, 47–55. <https://doi.org/10.12740/APP/99147>.
4. Acharya, Sourya, Vidyashree Hulkoti, Amol Andhale, Sugat Jawade, and Sneha Deshmukh. “Carbamazepine Induced Stevens-Johnson Syndrome (SJS).” *JOURNAL OF EVOLUTION OF MEDICAL AND DENTAL SCIENCES-JEMDS* 9, no. 51 (December 21, 2020): 3892–93. <https://doi.org/10.14260/jemds/2020/852>.
5. Aschaffenburg, G., 1911. *Handbuch der Psychiatrie*: hrsg. von G. Aschaffenburg (Vol. 4, No. 1). Deuticke.
6. Ghogare, Ajinkya Sureshrao, Pradeep Shriram Patil, and Ganpatlal Kodarbhai Vankar. “A Protocol of Case Control Study of Childhood Trauma and Alexithymia in Persons with Alcohol Dependence Syndrome.” *JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH* 14, no. 9 (September 2020). <https://doi.org/10.7860/JCDR/2020/44593.14018>.
7. Janicak, P., Davis, J., Preskorn, S., et al. (1997). Management of acute psychosis. In *Principles and practice of psychopharmacotherapy* (eds. P.G. Janicak, J.M. Davis, S.H. Preskorn, and F.J. Ayd) (2nd edn), pp. 110–39. Williams and Wilkins, Baltimore, MD.
8. Ting - Kai Li1, Brenda G Hewitt, the Alcohol Dependence Syndrome, 30 Years Later: A Commentary. The 2006 H. David Archibald Lecture, Epub 2007 Aug 4.
9. Kelkar, Parisha. “prevalence of substance abuse in patients of schizophrenia.” *Indian journal of Psychiatry* 61, no. 9, 3 (January 2019): S472–73.
10. Patel, Tejas, V, Mehul J. Brahmabhatt, and Ganpat K. Vankar. “Prevalence of Alcohol Use Disorders in Hospitalised Male Patients.” *Archives of Psychiatry and Psychotherapy* 20, no. 4 (December 2018): 47–55. <https://doi.org/10.12740/APP/99147>.
11. Gedam, Sachin Ratan, Ajab Dhabarde, Pradeep S. Patil, Animesh Sharma, Kanika Kumar, and Vijay Babar. “Psychiatric Comorbidity, Severity of Dependence and Liver Enzymes Dysfunction among Alcohol Dependent Individuals: A Cross - Sectional Study from Central Rural India.” *Journal of Clinical and Diagnostic Research* 13, no. 4 (April 2019): VC1–5. <https://doi.org/10.7860/JCDR/2019/40368.12759>.