

ADDITION OF COGNITIVE BEHAVIORAL THERAPY WITH MOTOR CONTROL EXERCISES ON PAIN AND FUNCTION IN ELDERLY INDIVIDUALS WITH CHRONIC LOW BACK PAIN

ABSTRACT

Introduction

Low back pain is a common, disabling, costly medical problem in most countries. CLBP is one of the most frequently reported conditions affecting the older adults. Numerous studies have also documented the associations of pain-related beliefs and appraisals with pain intensity and related problems, including depression, physical disability, and activity and social role limitations. Recently there is an increased attention to the role of cognitions in mood, anxiety, and other psychological disorders sparked interest in incorporating cognitive therapy techniques into behavioral therapies for chronic pain. CBT is a form of psychological therapy. It focuses on actions are the behavioral part. CBT is a feasible treatment for cognitively intact, older persons with CLBP, and may be efficacious as well. The aim of the study is to analyze the effect of cognitive behavioral therapy with Motor control exercises programme on CLBP

Methods

Study is an experimental study design includes 40 patients with CLBP with more than 6 months duration. 20 participants underwent motor control exercises with moist heat pack and 20 participants underwent motor control exercises with cognitive behavioral therapy and moist heat pack. Study was conducted for 8 weeks of duration and the outcome measures were pain and disability which was measured using numerical pain scale and Oswestry disability index.

Results

The between group comparison was made using the unpaired t test and the results were found to be 10.15 ± 0.502 , at the 95% of the Confidence intervals for disability. The between group comparison was made using the unpaired t test and the results were obtained are 7.47 ± 0.22 , at the 95% of the Confidence intervals for Pain.

Conclusion

The study concluded that motor control exercises with cognitive behavioral therapy were significantly effective in reducing pain and disability.

Key words: Chronic low back pain, motor control exercises, cognitive behavioral therapy, Pain, Oswestry disability index.

INTRODUCTION

Chronic low back pain in elders arises from age related changes in the lumbar spine and these changes were closely associated with the lifestyle of the individuals¹. Low back pain (LBP) is the most common health problem among older adults that results in pain and disability². Development of the low back pain has strongly associated with postural load conditions in elders. Prevalence of low back pain exceeded 70% of elders working in rural/ mountain areas and 42% in those in city. Poor posture, irregular movement of the lumbar spine with reduction of balance between the muscles enhances the nociceptive stimuli. Motion restriction due to pain leads to the contracture of intervertebral joints and the atrophy of the other lumbar spine structures, resulting in a vicious circle of pain³. Older adults aged 65 years or above, are the second most common age group to visit physicians for LBP⁴. Severe LBP usually results in poor treatment outcomes and functional disability⁵. Management of low back pain is very costly and there are a large increasing proportion of health care expenditures without evidence of corresponding improvements in outcomes⁶. The study showed that the major costs of LBP can be identified with the chronic and recurrent LBP. There are different ways in the management of low back pain that has thrived through decades of physiotherapy practice like Cryotherapy, Thermotherapy, TENS, Biofeedback, Ultrasound, SWD and traction^[7,8,9]. More recently a new protocol has been developed for retraining control of the stabilizing muscles around the spine and the pelvis. Spinal segmental stabilization exercises are also called as Motor control exercises which plays an important role in the management of back pain and neck pain are gaining the popularity¹⁰. Re-education of postural muscles may be achieved using specific stabilization exercises. The aim of the study is to correct imbalances of activity between more deeply placed stability muscles and more superficially placed mobilizing counterparts. The motor control exercise program is based on the treatment approach aimed to improve the function of specific muscles of the low back in frequent sessions^[11,12]. Behavioral Therapy (BT) and Cognitive Behavioral Therapy (CBT) for CLBP are terms for psychological interventions that often get applied inter-changeably in the CLBP literature. Therapies based upon these principles seek to help the patient with pain reduce symptom intensity, regain functioning, and reduce suffering.

Many techniques get incorporated into this form of therapy and rarely are single components applied in actual practice¹³. Numerous studies have conducted on the therapeutic efficacy of CBT techniques, which are a key component in most multidisciplinary/interdisciplinary pain management programs. The effectiveness of the CBT in addition to the motor control exercises was not well discussed. So this study aims to identify the effect of addition of the CBT with the motor control exercise on pain and disability in chronic low back pain in elders.

METHODOLOGY


Study was accepted by institutional ethical committee, it is an experimental study design with control group and experimental group of participants. 40 subjects with chronic low back pain more than 55 years were included in the study. A brief explanation was given to all the participants and consent was obtained from every individual. All subjects were divided into two equal groups. Subjects were chosen following selection criteria include both gender with age group of 55-68 years, have low back pain more than 1-year duration, not having any neurological involvement or cardiovascular, non-radiating pain or no weakness in the lower limb. Study was conducted for duration of 6 months. Motor control group (MCG) receives a set of exercises as per the protocol prescribed by Richardson et al., 1999. The exercises were done alternate days for 60 minutes of duration. The Cognitive behavioral therapy group (CBTG) receives a set of instructions which was prescribed based on protocols and motor control exercises for 60 minutes. Both the groups underwent moist heat therapy for 10 minutes. The individual participants underwent 8 weeks of exercise programme. Outcome measures are pain and functional disability, Pain was assessed using visual analog scale and functional disability was assessed using Oswestry disability index¹⁴.

RESULTS

Data analyzed using SPSS 20.1. Student 't' test was used to find out the difference between the groups, paired 't' was used to find out the difference between pretest and posttest between groups, whereas Unpaired 't' test was used to find out the difference between Posttest variables between the groups. The values were calculated using the paired t test to compare the values within the groups. The values of the disability measured by the Oswestry disability index for the Group A are 19.52 ± 0.540 , at the 95% of the Confidence intervals. The results of the study show that there was a significant difference obtained between the pretest and the posttest values. Same analysis was conducted for the Group B participants and the values were


23.89±0.647 at 95% of CI. The between group comparison was made using the unpaired t test and the results were found to be 10.15 ±0.502, at the 95% of the Confidence intervals. Figure I shows the graphical representation of the disability scores.

Figure 1: Disability



The values were calculated using the paired t test to compare the values within the groups. The values of the pain values measured by the numerical pain rating scale for the Group A are 8.37 ± 0.22 , at the 95% of the Confidence intervals. The results of the study show that there were significant differences obtained between the pretest and the posttest values. Same analysis was conducted for the Group B participants and the values were 19.71 ± 0.19 , at the 95% of the Confidence intervals. The between group comparison was made using the unpaired t test and the results were obtained are 7.47 ± 0.22 , at the 95% of the Confidence intervals. GraphII shows the graphical representation of the pain scores.

Figure 2: Pain



DISCUSSION

Like among young adults, the majority of LBP among older adults has no definite pathology. These patients experience LBP that is altered by posture, activity, or time of the day. Non-specific LBP may originate from different pain sources. Many older patients with chronic LBP display physical findings. Therapeutic exercise should be prescribed based on specialist evaluation of the pain reduction achieved by medication. Motor control exercises are multi component exercise program that aims at improving strength, flexibility and endurance. These active form of exercises are designed to strengthen muscles, to support the spine and helps to prevent low back pain¹⁵. Studies have supported that segmental stabilization and muscular strengthening in chronic low back pain. They suggested that both techniques were effective as they decreased pain and reduced disability¹⁶. Cognitive functional therapy helps people to identify abnormal patterns of posture and focusing on changing it so it enhances the effect of traditional rehabilitation of pain. Also, it replaces abnormal thoughts with more constructive ways of thinking¹⁷. The level of body control and awareness (body perception), as well as their ability to relax their trunk muscles and normalize pain provocative postural and movement behaviors are the main target of cognitive functional therapy and has an effect on pain management¹⁸. The significant decrease in back pain with more evidence in pain reduction in group B than in group A might attributed to cognitive functional therapy as it directly challenges the behaviors in a cognitively integrated, functionally specific and graduated manner. As in group B the study focused on changing the posture by cognitively awareness of it and also cognitively changing the perception of pain by focusing on good perception of it and posture after the treatment program¹⁹. Group A and Group B participants underwent motor control

exercises where both the groups has produced significant improvement as similar to the study²⁰. Similar studies postulated that the co-contraction of the TrA and LM muscles is the basis of the lumbo-sacral biomechanical stability and that these muscles act by reducing the compressive overloads, attenuating or eradicating pain perception²¹. The outcomes of the current study is also found that even after pain remission in patients with low back pain, proper deep muscle reestablishment often did not happen and that specific physical therapy focusing on those muscles was necessary²². The limitation of the study includes a small number of participants and the follow up was not made after the study, personal hygiene, activities are not under the control, the study not used any modalities, and there is no particular work population selected²³.

CONCLUSION

The study concluded that motor control exercises with cognitive behavioral therapy were significantly effective in reducing pain and disability.

References

1. Bressler HB, Keyes WJ, Rochon PA, Badley E. The prevalence of low back pain in the elderly: a systematic review of the literature. *Spine*. 1999 Sep 1;24(17):1813.
2. Podichetty VK, Mazanec DJ, Biscup RS. Chronic non-malignant musculoskeletal pain in older adults: clinical issues and opioid intervention. *Postgraduate medical journal*. 2003 Nov 1;79(937):627-33.
3. Yamamoto H. Low back pain due to degenerative disease in elderly patients. *the elderly*. 2003;45(13):42.
4. Cypress BK. Characteristics of physician visits for back symptoms: a national perspective. *American journal of public health*. 1983 Apr;73(4):389-95.
5. Cayea D, Perera S, Weiner DK. Chronic low back pain in older adults: What physicians know, what they think they know, and what they should be taught. *Journal of the American Geriatrics Society*. 2006 Nov;54(11):1772-7.
6. Frymoyer JW, Gordon SL. New perspectives on low back pain. Park Ridge, Illinois: American Academy of Orthopaedic Surgeons. 1989:191-3.
7. Foster NE, Thompson KA, Baxter GD, Allen JM. Management of nonspecific low back pain by physiotherapists in Britain and Ireland: a descriptive questionnaire of current clinical practice. *Spine*. 1999 Jul 1;24(13):1332.
8. Li LC, Bombardier C. Physical therapy management of low back pain: an exploratory survey of therapist approaches. *Physical therapy*. 2001 Apr 1;81(4):1018-28.
9. Gracey JH, McDonough SM, Baxter GD. Physiotherapy management of low back pain: a survey of current practice in Northern Ireland. *Spine*. 2002 Feb 15;27(4):406-11.

10. Richardson C, Jull G, Hides J, Hodges P. Therapeutic exercise for spinal segmental stabilization in low back pain. London: Churchill Livingstone; 1999.
11. BPhy SO, BPhy DF, MPhty GJ. Recent advances in therapeutic exercise for the neck: implications for patients with head and neck pain. *Australian Endodontic Journal*. 2003 Dec;29(3):138-42.
12. Kladny B, Fischer FC, Haase I. Evaluation of specific stabilizing exercise in the treatment of low back pain and lumbar disk disease in outpatient rehabilitation. *Zeitschrift fur Orthopadie und ihre Grenzgebiete*. 2003 Jul 1;141(4):401-5.
13. Patroncini M. Impaired Trunk Muscle Function in Sub-acute Neck Pain: Etiologic in the Subsequent Development of Low Back Pain?. *manuelletherapie*. 2005 Oct;9(04):187-8.
14. Macea DD, Gajos K, Calil YA, Fregni F. The efficacy of Web-based cognitive behavioral interventions for chronic pain: a systematic review and meta-analysis. *The Journal of Pain*. 2010 Oct 1;11(10):917-29.
15. Gatchel RJ, Rollings KH. Evidence-informed management of chronic low back pain with cognitive behavioral therapy. *The Spine Journal*. 2008 Jan 1;8(1):40-4.
16. Ali S, Sajjad AG, Keramat KU, Darian H. CHRONIC LOW BACK PAIN; EFFECTS OF THE LUMBAR STABILIZATION EXERCISES ON PAIN, RANGE OF MOTION AND FUNCTIONAL DISABILITY IN THE MANAGEMENT. *Professional Medical Journal*. 2017 Apr 1;24(4).
17. França FR, Burke TN, Hanada ES, Marques AP. Segmental stabilization and muscular strengthening in chronic low back pain: a comparative study. *Clinics*. 2010;65:1013-7.
18. Martel MO, Thibault P, Sullivan MJ. The persistence of pain behaviors in patients with chronic back pain is independent of pain and psychological factors. *PAIN®*. 2010 Nov 1;151(2):330-6.
19. O'Keeffe M, Purtill H, Kennedy N, O'Sullivan P, Dankaerts W, Tighe A, Allworthy L, Dolan L, Bargary N, O'Sullivan K. Individualised cognitive functional therapy compared with a combined exercise and pain education class for patients with non-specific chronic low back pain: study protocol for a multicentre randomised controlled trial. *BMJ open*. 2015 May 1;5(6):e007156.
20. Elshawi AM, Hamada HA, Zakaria HM, Ragab WM. The effect of cognitive functional therapy in low back pain due to postural scoliosis: A randomized controlled trial. *International Journal of Medical Research & Health Sciences*. 2016;5(12):14-22.
21. Arun B. Effect of Myofascial Release Therapy with Motor Control Exercises on Pain, Disability and Transversus Abdominis Muscle Activation in Chronic Low Back Pain. *A Journal of Health Professions*. 2013;3(3):28-32.
22. Snijders CJ, Ribbers MT, de Bakker HV, Stoeckart R, Stam HJ. EMG recordings of abdominal and back muscles in various standing postures: validation of a biomechanical model on sacroiliac joint stability. *Journal of Electromyography and Kinesiology*. 1998 Aug 1;8(4):205-14.

23. Hides JA, Richardson CA, Jull GA. Multifidus muscle recovery is not automatic after resolution of acute, first-episode low back pain. *Spine*. 1996 Dec 1;21(23):2763-9.

UNDER PEER REVIEW