

## Causes of Acute Abdominal Pain in Geriatrics; **Review Article**

### **Abstract:**

**Background:** Aging affects all functions of the gastrointestinal system (GIS). The elderly commonly has atypical presentation of a disease with more subtle symptoms; hence physicians who are not familiar with these might miss the opportunity to make a diagnosis of abdominal pain in timely manner. **The** main objective of our study is to summarize the current evidence regarding causes and management of acute abdominal pain in elderly.

**Methodology:** MEDLINE, Embolic, CINAHL, PsycInfo, and ASSIA were searched from 2000 until April 2021, and reference lists of included studies were searched. Studies were included that described causes and management of abdominal pain in elderly. No software has been utilized to analyze the data. The data was extracted based on specific form that contains (Author's name, publication year, country, methodology and results). **Results:** The study included 7 papers. 2 cross sectional, 2 prospective and 3 retrospective studies all reporting causes and management of acute abdominal pain in elderly. **Conclusion:** the most common causes of abdominal pain in elderly population were biliary disease, appendicitis and bowel obstruction. Renal colic, hernia and ischemia were also reported in different rates. Diagnosis and management of abdominal pain especially in elderly should be immediate to avoid potential complications.

Keywords: abdomen, pain, causes, management, acute, elderly, geriatrics

## **Introduction:**

Aging affects all functions of the gastrointestinal system (GIS): motility, enzyme and hormone secretion, digestion, and absorption. While there is no GI disease that is specific and limited to advanced age, some illnesses are more prevalent in this age group and may require different management [1].

An acute abdomen is a condition that demands urgent attention and treatment. The acute abdomen may be caused by an infection, inflammation, vascular occlusion, or obstruction. Gastrointestinal (GI) changes in the elderly are common, and despite some GI disorders being more prevalent in the elderly, there is no GI disease that is limited to this age group [2]. While some changes associated with aging GI system are physiologic, others are pathological and particularly more prevalent among those above age 65 years. Abdominal pain in older adults is a concerning symptom common to a variety of diagnoses with high morbidity and mortality [3]. While abdominal pain is about 10% of all complaints presenting to emergency department (ED), 20% of these patients are known to be geriatric patients.[4]

The causes of an acute abdomen in this population are similar to those found in younger patients with some differences in frequency. In each case, a sense of urgency to make the proper diagnosis and then institute appropriate therapy is critical [5]. The elderly commonly has atypical presentation of a disease with more subtle symptoms; hence physicians who are not familiar with these might miss the opportunity to make a diagnosis in timely manner [6, 7]. Polypharmacy and medication side effects further contribute to the complexity of the clinical picture and can derail treating physicians in the wrong direction. Additionally, polypharmacy and co morbidities predispose elderly patients to a more complicated clinical course and increase the probability for development of complications [8].

Acute abdominal pain in the elderly patient presents a significant and challenging problem. In the elderly population, rapid recognition is particularly important; because the patient may have delayed seeking medical care and may have uncommon illnesses complicating the ultimate course [9]. Diagnostic accuracy is lower, and mortality far higher, than in younger patients. Reasons for these differences are multifactorial: the case mix is different, the evolution and prognosis of specific diseases are different, and the ways in which diseases present are also different in elderly patients [10, 11].

## **Study Objective:**


Therefore, the main objective of our study is to summarize the current evidence regarding causes and management of acute abdominal pain in elderly.

## Methodology:

**Search strategy:** Search strategies were combined for papers on abdominal pain causes, geriatric GI disorder, and emergency management of abdominal pain in elderly and most prevalent causes of abdominal pain in elderly together with qualitative methodological filters. MEDLINE, Embolic, CINAHL, PsycInfo, and ASSIA were searched from 2000 until April 2021, and reference lists of included studies were searched. As showed in Figure (1).


**Study selection:** Studies were included that described causes and management of abdominal pain in elderly. Papers were included that either focused on causes or management. Papers were excluded that examined other specific objectives. Paired reviewers independently screened titles and abstracts of all identified references. Paired reviewers independently assessed full-text articles. Disagreements were resolved by discussion. Non-English studies were excluded.

**Statistical Analysis:** No software has been utilized to analyze the data. The data was extracted based on specific form that contains (Author's name, publication year, country, methodology and results). These data were reviewed by the group members to determine the initial findings, and the current evidence regarding causes and management of acute abdominal pain in elderly. Double revision of each member's outcomes was applied to ensure the validity and minimize the mistakes.



El  
ig  
ib  
ili  
ty

n  
cl  
u  
d  
e  
d



**Figure (1): PRISMA chart showing data extraction process**

**Results:**

The search of the mentioned databases returned a total of 107 studies that were included for title screening. 94 of them were included for abstract screening, which lead to the exclusion of 33 articles. The remaining 61 publications full-texts were reviewed. The full-text revision lead to the exclusion of 54 studies, and 7 were enrolled for final data extraction (Table 1).The included studies had different study designs,

While abdominal pain is about 10% of all complaints presenting to emergency department (ED), 20% of these patients are known to be geriatric patients.[4]

**Table (1) show studies conducted by Pinar HENDEN ÇAM et al [12];** 48.2% were male, and 51.8% were female. An internal disease was detected in 76.8% of the patients as an origin of abdominal pain. 258 patients had a medical source (76.8%) and 78 had a surgical source (23.2%). Of the patients with medical diagnoses, 53.5% were in 65–74 years group and 46.5% were in 75 years and above group. Of the patients with surgical diagnoses, 48.7% were in 65–74 years group and 51.3% were in 75 years and above group. The most frequent finding accompanying abdominal pain was vomiting. The most frequent chronic disease accompanying abdominal pain was hypertension in both age groups.48.8% of the patients with abdominal pain

were hospitalized and they were hospitalized mostly by gastroenterology ward (24.8%). **Laurell H et al. found that [13];** Hospital stay increased from 170 days per 100 emergency admissions in the control group to 320 and 458 days in the younger and older study groups, respectively. At the emergency department, older patients were more often misdiagnosed than control patients (52 vs. 45%;  $p = 0.002$ ). At discharge the diagnosis was more accurate in the control group (86 vs. 77%;  $p < 0.0001$ ). Hospital mortality was higher among older patients (23/831 vs. 2/1,458;  $p < 0.001$ ). The admission-to-surgery interval was increased (1.8 vs. 0.9 days,  $p < 0.0001$ ) in patients  $\geq 65$  years. Rebound tenderness ( $p < 0.0001$ ), local rigidity ( $p = 0.003$ ) and rectal tenderness ( $p = 0.004$ ) were less common in the older than in the control patients with peritonitis. In patients  $\geq 65$  years, C-reactive protein did not differ between patients operated on and those not, contrary to the finding in patients  $< 65$  years ( $p < 0.0001$ ).

**Table (2) show studies of Espinoza, Ricardo et al. who reported that [14];** 66% of elderly patients had concomitant diseases, that were multiple in 63%. Thirty one percent had postoperative complications. Compared with their younger counterparts, elderly patients required significantly ( $p < 0.05$ ) more admissions to intensive care units (2.7 and 24.2% respectively), more connections to mechanical ventilation (1.4 and 8.9% respectively) and longer hospital stays (5.4 $\pm$ 7.4 and 12.4 $\pm$ 10.9 days, respectively). In this series overall mortality was 6.7%, being 0.6% for young patients and 11.1% for the surgical group over 65 years old. **Osterwalder, Isabelle et al. found that [15];** thirty-day mortality was comparable to that of all other ED patients (2.2% vs. 2.1%). Patients with abdominal pain had a low risk of representation, and the majority of representations due to missed diagnoses were of benign origin. The high incidence of extra-abdominal causes is noteworthy, as this may induce change to differential diagnosis of abdominal pain.

**In table (3); Cervellin, Gianfranco et al. [16] reported that** Renal colic was the most frequent cause of **emergency department** (ED) admission in men, whereas **nonspecific abdominal pain** (NSAP) was more prevalent in women. Urinary tract infection was higher in women. Overall, 885 patients (16.57%) were hospitalized. Four hundred and eighty-five patients had repeated ED visits throughout the study period. Among these, 302 patients (6.46%) were readmitted within 30 days, whereas 187 patients (3.82%) were readmitted within 5 days. Renal colic was the first cause for ED readmission, followed by NSAP. In 13 cases readmitted to the ED within 5 days, and in 16 cases readmitted between 5–30 days the diagnosis was changed. **Gardner, Carly S et al. [17] reported that;** Utilization of abdominopelvic CT in geriatric patients presenting to the ED with acute abdominal symptoms strongly influences clinical management and significantly affects disposition. Diagnoses were clinically unsuspected prior to CT in 43% ( $p < 0.05$ ), with significant difficulty in diagnosing SBO ( $p < 0.05$ ), diverticulitis ( $p < 0.01$ ), and colonic obstruction ( $p < 0.01$ ). Positive CT results influenced treatment plans in 65%, surgical in 48%, and medical in 52%. Disposition from the ED was significantly affected by CT ( $p < 0.001$ ), 65% of admissions with positive CT ( $p < 0.001$ ) and 63% of discharges with negative CT ( $p < 0.001$ ). **Costamagna, D et al. [18] reported that;** mean age of the patients was 78 years and the

male/female ratio was 149/142. A total of 126 patients (43%) had one or more associated disease. Mesenteric ischaemia and secondary peritonitis were the most important causes of fatal outcome (respectively 42 and 17% of mortality).

**Table 1: Cross sectional studies:**

<b>Author, year,</b>	<b>Country</b>	<b>Study type</b>	<b>Sample Size</b>	<b>Outcomes</b>
<b>Pinar HENDEN ÇAM et al. 2018</b>	Turkey	Hospital-based cross-sectional study	336 patients	Most common Prediagnosis were biliary diseases and diseases related to biliary tract followed by nonspecific abdominal pain, abdominal pain secondary to malignity, illus, and acute gastroenteritis, respectively. The most frequent finding accompanying abdominal pain was vomiting. Surgical treatments were applied to the 17.6% of the patients with abdominal pain.
<b>Laurell, H et al. (2006)</b>	Sweden	Cross sectional	557 patients aged 65-79 years and 274 patients aged > or = 80 years were registered	A specific diagnosis, i.e. other than 'nonspecific abdominal pain', was established in 76 and 78% of the patients aged 65-79 and > or = 80 years respectively. Older patients were more often misdiagnosed than control patients.

**Table (2): Prospective studies:**

<b>Author, year,</b>	<b>Country</b>	<b>Study type</b>	<b>Sample Size</b>	<b>Outcomes</b>

<b>Espinoza, Ricardo et al. (2004)</b>	Spain	Prospective study	45 patients	The causes accounting for 71% of acute abdominal pain were bilio-pancreatic diseases (31.1%), intestinal adhesive obstruction (17.7%), complicated abdominal wall hernia (13.7%), and complications of peptic ulcer disease (8.9%). Sixty four percent required surgical treatment.
<b>Osterwalder, Isabelle et al. 2020</b>	Switzerland	Prospective monocentric all-comer study	3960 screened presentations, 480 (12.1%) were due to Abdominal Pain	Among 63 (13.1%) related representations, the most prevalent causes were cholelithiasis, gastroenteritis, and urinary retention. A missed diagnosis was attributed to 27 (5.6%) presentations. Extra-abdominal causes were identified in 162 (43%) presentations.

**Table (3): Retrospective Studies:**

<b>Author, year,</b>	<b>Country</b>	<b>Study type</b>	<b>Sample Size</b>	<b>Outcomes</b>
<b>Cervellin, Gianfranco et al. (2016)</b>	Italy	Retrospective analysis	5,340 cases	The most frequent causes were nonspecific abdominal pain (NSAP) (31.46%), and renal colic (31.18%). Biliary colic/cholecystitis, and diverticulitis were more prevalent in patients aged >65 years (13.17% vs. 5.95%, and 7.28% vs. 2.47%, respectively). Appendicitis

				(i.e., 4.54% vs. 1.47%) and renal colic (34.48% vs. 20.84%) were more frequent in patients aged <65 years. NSAP was the most common cause in both age classes.
<b>Gardner, Carly S et al. (2015)</b>	USA	Retrospective study	464 patients	The most common diagnoses were SBO (18%), diverticulitis (9%), non-ischemic vascular-related emergency (6%), bowel ischemia (4%), appendicitis (3%), and colonic obstruction (2%).
<b>Costamagna, D et al. (2009)</b>	Italy	Retrospective study	291 consecutive patients	The most common causes for an emergency operation were mechanical bowel obstruction (45%), hollow viscous perforation (18%) and strangulated hernia (18%). 234 patients (80%) recovered and were free from major complications. The remaining 57 (20%) developed at least one major complication (including death). The 30-days postoperative deaths were 33 (11%).

### **Discussion:**

As the life expectancy of the community increases, clinicians can expect an increase in geriatric patients presenting with abdominal pain .Compared with younger patients, this patient population is less likely to present with classic symptoms, physical examination findings, and laboratory values of abdominal disease [19].

Conditions causing acute abdominal pain may vary, from conditions needing immediate intervention, to relatively mild presentations needing careful observation to avoid over investigation and unnecessary interventions [20]. Common causes of an acute abdomen include acute appendicitis, cholecystitis, pancreatitis, and diverticulitis. Acute peritonitis is a cause of acute abdomen and can result from rupture of a hollow viscus or as a complication of inflammatory bowel disease or malignancy. Vascular events causing an acute abdomen include mesenteric ischemia and ruptured abdominal aortic aneurysm. Urologic conditions including ureteral colic and pyelonephritis can also present as acute abdominal pain. Many authors include small bowel obstruction as a cause of acute abdomen [21].

As age progresses, contraction ability of the gallbladder, in response to cholecystokinase enzyme, decreases. Additionally, increased cholesterol and phospholipid content of the bile causes gallbladder stones and increased biliary tract diameter results in biliary diseases [22]. Among the studies included in our results; three studies reported biliary disorder as the first cause of abdominal pain in elderly [11, 13, 15]. The mortality rate of elderly patients diagnosed with cholecystitis is approximately 10%. Cholecystitis is acalculous in approximately 10% of elderly patients with the condition. Classically, the diagnosis requires the presence of right upper quadrant pain associated with fever and leukocytosis [23]. Appendicitis is a less common cause of abdominal pain in elderly patients than in younger patients, but the incidence among elderly patients appears to be rising. Only approximately 10% of cases of acute appendicitis occur in patients older than 60 years, whereas one half of all deaths from appendicitis occur in this age group [24]. Cervellin, Gianfranco et al. [16] reported that appendicitis is more common cause of abdominal pain in elderly than other age groups. The initial diagnosis is incorrect in 40-50% of patients in this age range. Espinoza, Ricardo et al. [14] reported intestinal adhesive obstruction to be the cause of abdominal pain in (17.7%) of all studies cases. This was higher than a figure reported in previous study that bowel obstruction accounts for approximately 12% of cases of abdominal pain in elderly patients [25].

Diagnostic delay, late management and the risk of clinical worsening are the leading concerns of many EPs during the evaluation of patients with AAP. A comprehensive physical examination, close observation and serial diagnostic testing were found to be effective means to lower the risk of adverse outcome [26]. This conclusion is supported by the data of a systematic review and meta-analysis of the scientific literature, showing that the length of hospital stay and the rate of complications or readmission were not significantly different when comparing active observation with early laparoscopic intervention [27]. Additional evidence was brought that an observational period of 10 hours enhanced the ability to diagnosing appendicitis in patients with an intermediate probability [28]. In another prospective study including 220 patients of all ages admitted with AAP, a substantial decrease (i.e., from a 33% to a 5%) of negative findings on laparotomy was observed at the end of follow-up [29]. Interestingly, the data of our retrospective analysis confirms a large use of active clinical observation during ED stay, since 52% of our patients were discharged after more than 4 hours of LOS in the ED [30].

## Conclusion:

After summarizing the current studies regarding causes and management of abdominal pain in geriatrics; the most common causes of abdominal pain in elderly population were biliary disease, appendicitis and bowel obstruction. Renal colic, hernia and ischemia were also reported in different rates. Diagnosis and management of abdominal pain especially in elderly should be immediate to avoid potential complications.

## The already known about this topic:

It is recurrently confusing, which changes in GI function represent a part of normal aging processes and which of them are pathological results of a disease process.

## What this study adds:

- This study will summarize the current evidence regarding causes of acute abdominal pain in elderly.
- This study will also explore the current evidence regarding management of acute abdominal pain in elderly.

## References:

1. Dunic, Igor et al. Gastrointestinal Tract Disorders in Older Age. *Canadian journal of gastroenterology & hepatology* vol. 2019 6757524. 17 Jan. 2019. doi:10.1155/2019/6757524
2. Bhutto A., Morley J. E. The clinical significance of gastrointestinal changes with aging. *Current Opinion in Clinical Nutrition & Metabolic Care*. 2008;11(5):651–660. doi: 10.1097/MCO.0b013e32830b5d37.
3. D'Souza A. L. Ageing and the gut. *Postgraduate Medical Journal*. 2007;83(975):44–53. doi: 10.1136/pgmj.2006.049361.
4. Fagbohun C. F., Toy E. C., Baker B., III The evaluation of acute abdominal pain in the elderly patient. *Primary Care Update for OB/GYNS*. 1999;6(6):181–185. doi: 10.1016/S1068-607X(99)00021-9. [CrossRef] [Google Scholar].

5. Kauvar, D R. "The geriatric acute abdomen." *Clinics in geriatric medicine* vol. 9,3 (1993): 547-58.
6. Lasch H., Castell D. O., Castell J. A. Evidence for diminished visceral pain with aging: studies using graded intraesophageal balloon distension. *American Journal of Physiology-Gastrointestinal and Liver Physiology*. 1997;272(1):G1–G3. doi: 10.1152/ajpgi.1997.272.1.G1.
7. Hankey G. L., Holmes G. K. T. Coeliac disease in the elderly. *Gut*. 1994;35(1):65–67. doi: 10.1136/gut.35.1.65.
8. Durazzo M., Campion D., Fagoonee S., Pellicano R. Gastrointestinal tract disorders in the elderly. *Minerva Medica*. 2017;108(6):575–591. doi: 10.23736/S0026-4806.17.05417-9.
9. Jutel A. Sociology of diagnosis: A preliminary review. *Sociology of Health and Illness*. 2009;31(2):278–299
10. Tulchinsky TH, Varavikova EA. Measuring, Monitoring, and Evaluating the Health of a Population. *The New Public Health*. 2014;91-147. doi:10.1016/B978-0-12-415766-8.00003-3
11. de Dombal FT. Acute abdominal pain in the elderly. *J ClinGastroenterol*. 1994 Dec;19(4):331-5. doi: 10.1097/00004836-199412000-00016. PMID: 7876517.
12. Pınar HENDEN ÇAM et al. "Investigation of Geriatric Patients with Abdominal Pain Admitted to Emergency Department." *Current gerontology and geriatrics research* vol. 2018 9109326. 30 May. 2018, doi:10.1155/2018/9109326
13. Laurell H et al. "Acute abdominal pain among elderly patients." *Gerontology* vol. 52,6 (2006): 339-44. doi:10.1159/000094982
14. Espinoza, Ricardo et al. "Abdomen agudo en el adulto mayor" [Acute abdomen in the elderly]. *Revistamedica de Chile* vol. 132,12 (2004): 1505-12. doi:10.4067/s0034-98872004001200008
15. Osterwalder I, Özkan M, Malinovska A, Nickel CH, Bingisser R. Acute Abdominal Pain: Missed Diagnoses, Extra-Abdominal Conditions, and Outcomes. *J Clin Med*. 2020;9(4):899. Published 2020 Mar 25. doi:10.3390/jcm9040899
16. Cervellin, Gianfranco et al. "Epidemiology and outcomes of acute abdominal pain in a large urban Emergency Department: retrospective analysis of 5,340 cases." *Annals of translational medicine* vol. 4,19 (2016): 362. doi:10.21037/atm.2016.09.10
17. Gardner, Carly S et al. "Impact of CT in elderly patients presenting to the emergency department with acute abdominal pain." *Abdominal imaging* vol. 40,7 (2015): 2877-82. doi:10.1007/s00261-015-0419-7
18. Costamagna, D et al. "Acute abdomen in the elderly. A peripheral general hospital experience." *Il journal di chirurgia* vol. 30,6-7 (2009): 315-22.
19. Magidson, Phillip D, and Joseph P Martinez. "Abdominal Pain in the Geriatric Patient." *Emergency medicine clinics of North America* vol. 34,3 (2016): 559-74. doi:10.1016/j.emc.2016.04.008

20. Cartwright, Sarah L, and Mark P Knudson. "Evaluation of acute abdominal pain in adults." *American family physician* vol. 77,7 (2008): 971-8.`z
21. Li PH, Tee YS, Fu CY, Liao CH, Wang SY, Hsu YP, Yeh CN, Wu EH. The Role of Noncontrast CT in the Evaluation of Surgical Abdomen Patients. *Am Surg*. 2018 Jun 01;84(6):1015-1021.
22. Rossetti B., Spizzirri A., Migliaccio C., et al. Acute pancreatitis in the elderly: Our experience. *BMC Geriatrics*. 2009;9(1, article no. A47) doi: 10.1186/1471-2318-9-S1-A
23. Choung RS, Locke GR 3rd, Schleck CD, ZinsmeisterAR, Talley NJ. The effects of ageing on the onset and disappearance of unexplained abdominal pain: a population-based study. *Aliment PharmacolTher*. 2014 Jan. 39(2):217-25.
24. Xu Y, Jeffrey RB, Chang ST, DiMaio MA, Olcott EW. Sonographic differentiation of complicated from uncomplicated appendicitis: implications for antibiotics-first therapy. *J Ultrasound Med*. 2017 Feb. 36(2):269-77.
25. Reginelli A, Russo A, Pinto A, et al. The role of computed tomography in the preoperative assessment of gastrointestinal causes of acute abdomen in elderly patients. *Int J Surg*. 2014. 12 Suppl 2:S181-6..
26. Ditillo MF, Dziura JD, Rabinovici R. Is it safe to delay appendectomy in adults with acute appendicitis? *Ann Surg* 2006;244:656-60.
27. Agresta F, Ansaloni L, Catena F, et al. Acute appendicitis: position paper, WSES, 2013. *World J EmergSurg* 2014;9:26.
28. Maggio AQ, Reece-Smith AM, Tang TY, et al. Early laparoscopy versus active observation in acute abdominal pain: systematic review and meta-analysis. *Int J Surg* 2008;6:400-3.
29. Graff L, Radford MJ, Werne C. Probability of appendicitis before and after observation. *Ann Emerg Med* 1991;20:503-7.
30. Thomson HJ, Jones PF. Active observation in acute abdominal pain. *Am J Surg* 1986;152:522-5.