

FREQUENCY OF OVERWEIGHT AMONG WOMEN OF OVARIAN ENLARGEMENT PRESENTED WITH PELVIC PAIN COMPLAINS

ABSTRACT

Objective: To determine the frequency of overweight among females of ovarian enlargement and presented with pelvic pain complaints.

Methodology: This descriptive study was conducted at physiology department of Sindh University with collaboration of different health facilities of Tando Muhammad Khan and Hyderabad Sindh. All the females presented with ovarian enlargement with pelvis pain and age >18 years were included. Body mass index (BMI) was calculated as the ratio of an individual's weight in kilograms divided by the height in meters squared ($BMI = \text{kg/m}^2$). All the data was collected via study proforma. Data was analyzed by using SPSS version 20.

Result: A total of 239 women of ovarian enlargement were studied, their mean age was 32.79 ± 9.31 years. Neoplastic ovarian lesions were seen among 12.54% of the cases. Cystic ovarian lesions were among 30.54% of the females, 5.01% had ovarian mass and 21.33% females had mixed pathologies. BMI was normal among 62.76% cases, while 20.08% women were overweight, 03.34% were obese and 13.8% females were underweight.

Conclusion: Overweight females were observed to be highly frequent among females of ovarian enlargement. Cystic enlargement is more common as compared to solid or mixed enlargement.

Keywords: Ovarian Enlargement, BMI, Ultrasonography.

INTRODUCTION

The lower abdominal/pelvic pain is more common in women related to all age groups. Its diagnosis can be challenging because many symptoms and signs are insensitive and nonspecific.¹ Women are usually suffering from pelvic pain are mostly related to gynecological problems e.g ovarian enlargement, pelvic inflammatory diseases, etc. This is a very more common problem nowadays because of hormonal changes or imbalances, stress, hypertension, obesity, diabetes, and malignancies. However, a woman presenting with an abdominal or pelvic mass might complain of various symptoms but a significant majority will have no guiding/obvious symptoms at presentation although the final diagnosis may be life-threatening.² Ovarian cysts, also known as ovarian masses or adnexal masses, are frequently found incidentally in asymptomatic women.³ Ovarian cysts are sometimes found in the course of evaluating women for pelvic pain though the cysts may or may not be the cause of the pain.³ Polycystic ovary syndrome is a heterogeneous condition characterized by oligo-ovulation or anovulation, hyperandrogenism, menstrual irregularities and subfertility.^{4,5} Approximately 30–75% of women with PCOS are obese, with central obesity noted in 50–60% of women with PCOS, irrespective of BMI. Obesity encourages the phenotypic expression of PCOS and causes deterioration of hormonal and metabolic parameters in those who are already symptomatic.^{4,6} Obese women are more likely to have menstrual irregularity and anovulatory infertility than normal-weight women. In reproductive-age women, the relative risk of anovulatory infertility increases at a BMI of 24 kg/m^2 and

continues to rise with increasing BMI. Consistent with a pathophysiologic role for obesity, weight reduction can restore regular menstrual cycles in these women.⁷ Obesity exacerbates hormonal and clinical features of PCOS and women with PCOS appear at higher risk of obesity, with multiple underlying mechanisms linking the conditions. Lifestyle intervention is the first line in the management of PCOS to both prevent weight gain and induce weight loss; however improved engagement and sustainability remain challenges with the need for more research.⁸ Hence this study has been conducted to determine the frequency of overweight among females of ovarian enlargement and presented with pelvic pain complaints.

MATERIAL AND METHOD

This descriptive study was conducted at physiology department of Sindh University with collaboration of different health facilities of Tando Muhammad Khan and Hyderabad Sindh. All the females presented with ovarian enlargement with pelvis pain and age >18 years were included. All the women with ovarian cancer and those who were not agree to participate in the study were excluded. After taking informed consent a complete medical history and clinical examination was done. BMI was calculated as the ratio of an individual's weight in kilograms divided by the height in meters squared ($BMI = \text{kg}/\text{m}^2$). BMI was categorized as underweight ($BMI < 18 \text{ kg}/\text{m}^2$), normal weight ($BMI < 18.9-24.8 \text{ kg}/\text{m}^2$), overweight ($BMI 25.0-29.9 \text{ kg}/\text{m}^2$) and obese ($BMI > 30.0 \text{ kg}/\text{m}^2$). All the data was collected via study proforma. Data was analyzed by using SPSS version 20.

RESULTS

A total of 239 women of ovarian enlargement were studied, their mean age was 32.79 ± 9.31 years. Out of all neoplastic ovarian lesions were found in 28(12.54%) of the cases. Cystic ovarian lesions were among 73(30.54%) of the females, 10(5.01%) had ovarian mass and 51(21.33%) females had mixed pathologies. Majority of the females 150(62.76%) had normal BMI, while 48(20.08%) women were overweight, 08(03.34%) were obese and 33(13.8%) females were underweight as showed in table.1

Table: 1. Descriptive statistics of the age, ovarian lesions and BMI n=239

Variables		Statistics
Age	Mean±SD	32.79±9.31 years
Neoplastic enlargement	Yes	28(12.54%)
	No	211(87.46%)
	Cystic	105(43.80%)
	Mass	10(5.01%)
	Mixed	51(21.33%)
	Others	73(30.54%)
BMI	Underweight ($BMI < 18 \text{ kg}/\text{m}^2$)	33(13.8%)
	Normal BMI ($18.9-24.9 \text{ kg}/\text{m}^2$)	150(62.76%)
	Overweight ($BMI 25.0-29.9 \text{ kg}/\text{m}^2$)	48(20.08%)

	Obese (BMI >30.0kg/m ²)	08(03.34%)
--	-------------------------------------	------------

DISCUSSION

Overweight and obesity are common among adolescent girls and adult women with PCOS. In response to nutrient excess, adipocytes can enlarge (hypertrophy) or form new adipocytes (hyperplasia).⁹ Anti-Mullerian Hormone (AMH) concentrations reflect ovarian reserve and are correlated with the number of growing follicles and AMH concentrations were found to be higher in girls with obesity with PCOS compared to girls with obesity without PCOS of comparable age and pubertal status.⁹ In this study average age of the females was 32.79±9.31 years. Similarly, Rehman R et al¹⁰ reported that the mean age of obese PCO women was 32.02 ± 4.81 years. Although, Ramanand SJ et al¹¹ reported that the mean age of the females was 22.05±4.649 years, this average was lower as compared to our findings and this may be because of they studied only young PCOS patients. However, Yasin M et al¹² reported that the mean age of the PCO women was 24.93±5.67 years.

In this study neoplastic ovarian lesions were found in 28(12.54%) of the cases.

Cystic ovarian lesions were among 73(30.54%) of the females, 10(5.01%) had ovarian mass and 51(21.33%) females had mixed pathologies. Lee HJ et al¹³ found that approximately 6% of hens with ovarian tumors had polycystic ovarian condition, suggesting that polycystic ovarian condition may be a potential risk factor for the development of spontaneous ovarian cancer.

In this study 20.08% women were overweight and 08(03.34%) were obese. Some other studies stated that the women with PCOS have an increased prevalence of obesity and a consensus on the proportion of overweight and obese women with PCOS is not clear, with estimates of between 40% and 60%.^{14,15} Although Yasin M et al¹² reported that the 13% women were obese among PCO females. However, obesity is also associated with low-grade inflammation. Thus, it is possible that long-standing unresolved low-grade inflammation may be associated with the development of PCOS. However, it is unknown whether low-grade inflammation is a cause or an effect of PCOS.¹³ The majority of women with PCOS have insulin resistance and/or are obese and their elevated insulin levels increase GnRH pulse frequency and either contribute to or cause the abnormalities seen in the hypothalamic–pituitary–ovarian axis that lead to PCOS.¹⁶

Conclusion

Overweight females were observed to be highly frequent among females of ovarian enlargement. Cystic enlargement is more common as compared to solid or mixed enlargement. Person with normal weight is more suffered from right side ovarian pathology than left side Pathology. According to the sonographic findings study shows cystic ovarian enlargement are common in all age group however very in common in 20-30 age groups.

Ethical Approval:

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

Consent

As per international standard or university standard, patients' written consent has been collected and preserved by the author(s).

REFERENCES

1. Kruszka P, Kruszka SJ. Evaluation of acute pelvic pain in women. *American family physician*. 2010 Jul 15;82(2):141-7.
2. Pathiraja P. Abdominal Masses in Gynecology. *A Textbook of Gynecology for Less-Resourced Locations*. 2012;page 100-111
3. Ross E, Fortin C. Ovarian Cysts. *The Cleveland Clinic Foundation*. 2016. clevelandclinicmeded.com/medicalpubs/diseasemanagement/womens-health/ovarian-cysts/
4. Pandey S, Bhattacharya S. Impact of obesity on gynecology. *Women's Health*. 2010 Jan;6(1):107-17.
5. Pasquali R, Gambineri A. Polycystic ovary syndrome: a multifaceted disease from adolescence to adult age. *Annals of the New York Academy of Sciences*. 2006;1092(1):158-74.
6. Diamanti-Kandarakis E. Role of obesity and adiposity in polycystic ovary syndrome. *International journal of obesity*. 2007 Nov;31(2):S8-13.
7. Sam S. Obesity and polycystic ovary syndrome. *Obesity management*. 2007 Apr 1;3(2):69-73.
8. Barber TM, Franks S. Obesity and polycystic ovary syndrome. *Clinical endocrinology*. 2021 Oct;95(4):531-41.
9. Witchel SF, Oberfield SE, Peña AS. Polycystic ovary syndrome: pathophysiology, presentation, and treatment with emphasis on adolescent girls. *Journal of the Endocrine Society*. 2019 Aug;3(8):1545-73.
10. Rehman R, Mehmood M, Ali R, Shaharyar S, Alam F. Influence of body mass index and polycystic ovarian syndrome on ICSI/IVF treatment outcomes: A study conducted in Pakistani women. *International Journal of Reproductive Bio Medicine*. 2018 Aug;16(8):529.
11. Ramanand SJ, Ghongane BB, Ramanand JB, Patwardhan MH, Ghanghas RR, Jain SS. Clinical characteristics of polycystic ovary syndrome in Indian women. *Indian journal of endocrinology and metabolism*. 2013 Jan;17(1):138.
12. Yasin M, Yasmeen F. Polycystic ovarian syndrome; pattern of disease in patients. *Professional Med J* 2014;21(1): 179-184.
13. Lee HJ, Bahr JM, Bitterman P, Basu S, Sharma S, Abramowicz JS, Barua A. Polycystic ovarian condition may be a risk factor for ovarian tumor development in the laying hen model of spontaneous ovarian cancer. *Journal of immunology research*. 2018 Nov 25;2018.
14. Lo JC, Feigenbaum SL, Yang J, Pressman AR, Selby JV, Go AS. Epidemiology and adverse cardiovascular risk profile of diagnosed polycystic ovary syndrome. *J Clin Endocrinol Metab* 2006; 91: 1357–1363.
15. Pasquali R, Gambineri A, Pagotto U. The impact of obesity on reproduction in women with polycystic ovary syndrome. *BJOG* 2006; 113: 1148
16. Ruan X, Kubba A, Aguilar A, Mueck AO. Use of cyproterone acetate/ethinylestradiol in polycystic ovary syndrome: rationale and practical aspects. *The European Journal of Contraception & Reproductive Health Care*. 2017 May 4;22(3):183-90.