

Original Research Article

Microwave assisted extraction of Papaya leaves and investigation on antioxidant activity

Shobitharai¹, Divya Jyothi^{1*}, Swathi Das¹, Sumayya CM¹, Thabsheer A¹

¹Department of Pharmacognosy, NGSM Institute of Pharmaceutical Sciences (NGSMIPS), Nitte (Deemed to be University), Deralakatte, Mangaluru-575018, Karnataka, India.

ABSTRACT:

Microwave assisted extraction (MAE) has gained lot of attention due to its advantages such as less solvent consumption, short time period, higher extraction efficiency, therefore serves as better alternative for conventional extraction methods of plant materials. Plant phenolic compounds are important constituents responsible for reducing the oxidative stress that induces tissue damage which is the one of the major causative factors associated with the chronic disease. Papaya plant is a medicinal plant which became popular for the treatment of dengue fever due to its property. Considering the current medicinal importance of the papaya plant, the present study was aimed at microwave assisted extraction of phenolic content from papaya leaf using ethanol, water as solvent and investigate their antioxidant potential.

In order to compare the extraction efficiency of phenolic compounds, conventional extraction and microwave assisted extraction method was used to prepare the extracts. Then extracts were subjected to preliminary phytochemical analysis followed by the estimation of total phenolic content by using Folin-Ciocalteu method. Antioxidant activity was investigated by 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay.

The alcoholic and aqueous extracts of papaya leaf showed the presence of steroids, alkaloids, saponins, carbohydrates, phenolic compounds by preliminary phytochemical analysis. FTIR spectrum of both aqueous and ethanolic extract showed characteristic peak at 3314.62 cm^{-1} , 1635 cm^{-1} which provide evidence for presence of phenolic compounds. The total phenolic content of the alcoholic and aqueous leaf extracts from MAE was found to be 43.58mg and 80.58 mg/g papaya leaf powder of the Gallic acid equivalent (GAE), respectively. Aqueous solvent was found to be suitable for extraction of phenolic content from papaya leaf and Microwave assisted extracts showed higher phenolic content and therefore potential antioxidant activity. Therefore, papaya leaf is a good candidate to be used as a natural antioxidant for the treatment of various diseases.

Key words: Papaya, Anti-oxidant, Total phenolic content, phytochemical analysis, UV spectroscopy, FTIR, DPPH assay

INTRODUCTION

In recent years, global trend is increasing towards the use of natural antioxidant in the area of food science and complementary medicines in comparison with synthetic antioxidants which are toxic to human health [1]. Plants acts as rich source of natural antioxidants due to the presence of secondary metabolites mainly poly-phenolic compounds and flavonoids. Phenolic compounds acts as a reducing agents due to their redox potential, thereby acting as antioxidants, thereby have important role in lipid peroxidation [2]. Several extraction techniques such as Microwave extraction (MAE), supercritical fluid extraction, solvent extraction, Soxhlet extraction, refluxation methods are used for extraction of antioxidant constituents such as Polyphenolic compounds from Plant [3-6]. Among these MAE is popularly being used due to its higher efficiency of extraction [7]. Many reports have shown that MAE has more extraction potential than conventional method of extraction [8,9].

Carica papaya, commonly known as papaya belonging to the family Caricaceae, is a small, sparsely branched tree which is usually five to ten m tall with a single stem, spirally arranged leaves at the top of the trunk. . Young leaves of papaya are used for the treatment of jaundice, urinary complains, urinary tract infection, gonorrhoea, dressing wounds, vermifuge in colic, fever, beriberi etc [10]. Papaya leaf extract were also reported to have antimalarial and anti-plasmodial activities [11]. Significant use of leaf juice is found to be the capability to increase platelets and WBC and also repairs the liver [12]. Papaya leaf consist of many active constituents such as papain, chymopapain, cystatin, tocopherol, ascorbic acid, cyanogenic glucoside, flavonoids and glucosinolates [13]. Carpaine, dihydrocarpaine, and cyanogenic glycoside are the other active components of alkaloid family. Bitter taste of papaya leaves is due to the presence of pseudocarpaine and dehydrocarpaine [14]. Papaya also contains important flavonoids namely kaempferol, myricetin, quercetin etc [15].

The extract and isolated constituents of plant are known to possess various biological activities such as anti-hyperlipidemic, antidiabetic, anti-inflammatory and also act as free-radical scavengers. Further it has been proven that free radicals play a key role in the development of metabolic disorders and thereby affects the quality of life [16]. Various researchers have reported that phenolic compounds and flavonoids are one of the important secondary metabolites which act as very potent free radical scavengers. Phenolic compounds act as a reducing agent due to their redox potential, thereby acting as antioxidants [17]. Hence phytoconstituents with high amount of phenolic compounds are known to show protective effects in biological system against oxidative stress. Hence considering the medicinal importance of papaya leaf, in the current study an effort was made to extract the phenolic compound by MAE including investigation of antioxidant potential of the extract.

MATERIALS AND METHODS:

Collection and preliminary processing of plant material

Fresh green leaves of *Carica papaya* were collected from locality of Surathkal, Mangalore. The plant was identified and authenticated by an expert botanist at St. Aloysius College, Mangalore. The leaves were washed thoroughly with distilled water and then chopped into pieces.

Extraction of plant material

Extracts of the papaya leaves were prepared separately by conventional method (Refluxation) and by microwave assisted extraction (MAE) method.

Aqueous and ethanolic extracts were prepared separately by subjecting 10gms of chopped leaves to refluxation for a period of 8 hours with 200 ml of distilled water and ethanol respectively as the solvents.

Microwave Assisted Extraction (MAE) was done by using 10 grams of the leaves was in a microwave oven (CATA-R) working at a 800W irradiation power and 2450MHz frequency. MAE was done using ethanol and water as solvent at a temperature of 50°C for a period of 5mins [18].

After the extraction, solutions were filtered, filtrate was evaporated and concentrated using rotary flash evaporator to get dry extracts. The extracts obtained by soxhlation and MAE compared for the percentage yield and amount of phenolic content, thereby indicating their antioxidant activity.

After the extraction, solutions were filtered; filtrate was evaporated and concentrated using rotary vacuum evaporator to get dry extracts. The percentage yields of aqueous and ethanolic extracts were calculated.

Preliminary Phytochemical Screening (Qualitative Analysis)

All the papaya leaf extracts of were subjected to various phytochemical tests to determine the presence of various phyto-constituents [19].

Estimation of phenolic content

Phenolic content in papaya leaf extracts were estimated by Folin-Ciocalteu method [20]. Test solution was prepared by taking 100mg of extract was dissolved in 100ml of phosphate buffer (pH 6.8). 10ml of the above solution and diluted up-to 100ml with phosphate buffer. From this 4ml was transferred to 25ml volumetric flask to which 1.25ml of FC reagent and 2.5ml of sodium carbonate was added. The final mixture volume was adjusted with distilled water. For the calibration curve, Gallic acid was used as standard and standard solution of 50µg/ml was prepared in phosphate buffer (pH 6.8). Aliquots of 2, 4, 6, 8, 10 and 12 ml were taken in 6 different 25ml volumetric flasks from the stock solution. Into each, 1.25 ml of Folin-Ciocalteu reagent and 2.5ml of 20% sodium carbonate. The resulting blue colour was evaluated for absorbance at 765 nm on UV-Visible spectrophotometer (Shimadzu 1700) after keeping in dark at room temperature for 30min. Using the linear equation obtained from the standard plot, the phenolic content was estimated and depicted as gallic acid equivalent per gram of the plant material.

Fourier transfer Infrared spectroscopy

Fourier Transform Infrared spectroscopy can be considered as a powerful tool for identifying functional groups present in compounds. Extract from MAE was encapsulated in KBr pellet of FT-IR to prepare translucent sample discs. The spectrum of these samples was recorded using Bruker FTIR spectrophotometer.

Determination of antioxidant activity by DPPH assay

The radical scavenging activity was determined by the use of DPPH free radical [21]. Ascorbic acid was used as a standard by dissolving 10mg in 10ml of methanol as diluent. Serial dilutions were prepared using 10 µl, 20 µl, 30 µl, 40 µl and 50 µl of this standard and the volume made up-to 50 µl using methanol. To these 100 µl of DPPH was added and the absorbance was noted at a wavelength of 517nm after 30 minutes of incubation against blank taken as 50 µl of the diluent (methanol). The test solutions were also prepared in a similar

manner using 10mg of the 4 plant extracts and dissolving in 10ml of methanol. 100 µl of DPPH reagent and 50 µl of methanol were used as the control. Using the following equation, the percentage inhibition activity was calculated:

$$\% \text{ Inhibition} = [(A_o - A_I) / A_o] \times 100 \dots \dots \dots (i)$$

Where A_o stands for the absorbance of the control, and A_I denotes the absorbance of the extract/ standard .

RESULTS AND DISCUSSION

Extraction of plant material and Phytochemical Analysis

Aqueous and ethanolic extracts of papaya leaf were prepared by refluxation method. The yield of ethanolic and aqueous extract was found to be 4.05% and 6.6% respectively. The yield of the leaf extract of from MAE was found to be 26.67% and 46.45% using ethanol and water as solvents respectively. MAE produced the higher yield in comparison to refluxation method because MAE offers a rapid delivery of energy to a total volume of the solid matrix, efficiently and homogenously. Because natural moisture present within the plant absorbs microwave energy, cell disruption is promoted by internal superheating, which facilitates desorption of active constituents from the matrix thus improving the final yield [22]

Preliminary phytochemical analysis of papaya leaf extract confirmed the presence of steroids, alkaloids, saponins, carbohydrates, phenolic compounds

Estimation Total phenolic content by Folin-Ciocalteu method

Amount of polyphenolic content in papaya leaf extracts were estimated by applying the Folin-Ciocalteu method where gallic acid is opted as the standard. Absorbance of different concentration of gallic acid solutions was measured at 765 nm for preparation of standard plot which is shown in Fig.1.




Fig. 1: Standard plot of gallic acid in phosphate buffer pH 6.8 at 765nm

From the calibration equation, Refluxation method resulted in extraction of 39.58 mg and 65.58 mg/g Gallic acid equivalent (GAE) of polyphenolic content from papaya leaf extracts by using ethanolic and aqueous solvents respectively.

Ethanollic and Aqueous extracts obtained from MAE produced 43.58mg and 80.58 mg/g Gallic acid equivalent (GAE)of polyphenolic content respectively. Hence MAE using aqueous solvent was found to be superior for getting high extraction efficiency of phenolic content than refluxation method. The MAE system has been used in the extraction of several phyto-chemicals, including polyphenols, where it seems to provide a good yield of polyphenols in less time and consuming fewer solvents. MAE technique for polyphenols extraction depends on the type of material, solvent type and purity, power and time of microwave application, available sample surface area, as well as the operating temperature. The most critical factor is the type of sol-vent as its effects cut across the whole process, ranging from the solubility of the target components to the process efficiency. Hence, the solvent must be selected with care by considering both its affinity to the target compounds and its microwave energy absorption capability [23].Numerous reports state that aqueous mixtures of organic solvents are the most suitable for extraction of phenolic compounds from plant sources. Different plant material requires different solvent type for maximum extraction of phenolic compounds [24]. The aqueous solvent was found to be suitable for extraction of phenolic compounds from papaya leaves.

Fourier transfer infrared spectroscopy: The FTIR analysis of extracts obtained from MAE was done to determine the important functional groups present. FTIR spectrum of Parijata and Tamarind leaf extracts are shown in Fig 2&3




Fig.2: FTIR spectrum of Papaya leaf ethanolic extract obtained by MAE




Fig.3: FTIR spectrum of Papaya leaf aqueous extract obtained by MAE

The FTIR spectrum of both aqueous and ethanolic leaf extract shows characteristic band at 3342cm^{-1} corresponds stretching vibration of -OH and -H bonded alcoholic and phenolic groups. Band at 1638 mainly due to aromatic C=C and C=O vibrations [25].

DPPH radical scavenging activity:

Free radical scavenging activity using the DPPH method of different concentration of papaya leaf extracts are shown in **Table 1**. Result showed that as the concentration of the extracts was increased, DPPH radical scavenging activity also increased. At highest concentration ($100\mu\text{g}/\mu\text{l}$), papaya aqueous extracts obtained from MAE showed higher percentage of DPPH scavenging activity i.e.75.08%. Extracts. The extracts from MAE showed higher polyphenolic content and this can be correlated with their antioxidant activity. The antioxidant property of plant extracts is generally attributed to phenolic compounds because they can neutralize the free radicals by their multifunctional properties such as hydrogen donating, reducing, metal chelating, and singlet oxygen quenching properties. They also terminate the free radical chain reaction by forming a relatively stable phenoxy-radical intermediate [26].

Table 1: Antioxidant activity of Papaya leaf extracts by DPPH assay

Samples	Concentration $\mu\text{g}/\mu\text{l}$	% DPPH Scavenging activity
Aqueous extracts from MAE	20	32.30
	40	40.90
	60	67.47
	80	73.19
	100	75.08
Ethanolic Extracts from MAE	20	9.20

	40	23.78
	60	37.24
	80	49.11
	100	68.53
Aqueous Extracts from Refluxation	20	12.11
	40	32.29
	60	38.60
	80	49.46
	100	65.34
Ethanollic Extracts from Refluxation	20	8.207
	40	21.60
	60	35.67
	80	41.78
	100	56.78

CONCLUSIONS

Papaya leaves contain appreciable quantity of phenolic content, and act as potent free radical scavenger, hence it can be used as a source of natural antioxidants which will have higher potential in the treatment of various diseases arising due to involvement of free radical and hence could lead to a new field of future research. In comparison to refluxation method, MAE showed higher extraction efficiency for phenolic compound and hence showing higher antioxidant potency. It has been proved that polarity of the solvent, nature of the extracted compounds and extraction process highly affects therapeutic activities of the plant extracts. The antioxidant behaviour of the plant mainly depends on the amount of phenolic content and it enhances the total antioxidant capacity of medicinal plants.

ACKNOWLEDGEMENT

I am thankful to Nitte University and NGSM institute of pharmaceutical sciences for providing all the facilities and requirement for carrying out this work.

CONFLICT OF INTEREST: No conflict of interest exists.

REFERENCES:

1. Sofidiya MO, Odukoya OA, FAMILONI OB, Inya-Agha SI. Free radical scavenging activity of some Nigerian medicinal plants. *Planta Medica*. 2006 ;72(11):171.
2. Correia Da Silva TB, Souza VK, Da Silva AP, Lyra Lemos RP, Conserva LM. Determination of the phenolic content and antioxidant potential of crude extracts and isolated compounds from leaves of *Cordia multispicata* and *Tournefortia bicolor*. *Pharmaceutical biology*. 2010 ;48(1):63-9.
3. Sandeep DS, Nayak P, Nayana K, Nasiha, Alfish N, Kumar N, Kumar A. Developing anti-dandruff shampoo formulations using different Indian plant Herbs- An Eco-friendly Hair care Cosmetic. *Journal of Xi'an Shiyou University, Natural Science* Ed.2021; 17(9):684-693.

4. Reshma R, Achala B , Rajesh KS, Harish , Raman R. Evaluation of Anti-venom Activity of Leaf Extract of *Wedelia trilobata*. Journal of Xi'an Shiyou University, Natural Science Ed.2021; 17(9):609-613.
5. Hashif A, Khandige PS, Nayak P. Evaluation of antidepressant activity of *Garcinia cambogia* on experimentally induced depression in Mice. Journal of Xi'an Shiyou University, Natural Science Ed.2021; 17(9):55-60
6. Olivia J, Varghese, Shetty P, Sharanya M. Skeletal muscle relaxant potential of *Annona reticulata* L. leaf extract in swiss albino mice a preclinical study. Journal of Xi'an Shiyou University, Natural Science Ed.2021; 17(9):673-683
7. Ganzler K., Salgo A., Valko K. Microwave extraction-a novel sample preparation method for chromatography. J. Chromatogr. 1986;371:299-306.
8. Pan X., Niu G., Liu H. Microwave-assisted extraction of tea polyphenols and tea caffeine from green tea leaves. Chem. Eng. Process. 2003;42:129-133.
9. Hong N., Yaylayan V.A., Raghavan G.S.V., Paré J.R.J., Bélanger J.M.R. Microwave assisted extraction of phenolic compounds from grape seed. Nat. Prod. Res. 2001;15:197-204.
10. Kirtikar KR, Basu BD. Indian Medicinal Plants. 2nd ed. Dehradun. International Book Distributors. 1998:1097-99.
11. Udoh OD, East S. Determination of essential and nonessential metals concentration in papaya (*Carica papaya*) seeds, leaves and supporting soil of odoshakiso district in South East Oromia region, Ethiopia. Int J Res in PharmChem. 2014;4(1):202-16.
12. Noriko O, Nam HD, Emi K, Akira K, Sathoshi I, Chikao M. Aqueous extract of *Carica papaya* leaves exhibit anti-tumour activity and immunomodulatory effects. J Ethnopharmacol. 2010;27:760-67.
13. Vyas SJ, Khatri TT, Ram VR, Dave PN, Joshi HS. Biochemical constituents in leaf of *Carica papaya* - ethnomedicinal plant of Kachchh region. Int Lett Natural Sci. 2014;12:16-20.
14. Prasetya AT, Mursiti S, Maryani S, Jati NK. Isolation and identification of active compounds from papaya plants and activities as antimicrobial. Materials Sci Eng. 2018;349:1-6.
15. Nugroho A, Heryani H, Choi JS, Park HJ. Identification and quantification of flavonoids in *Carica papaya* leaf and peroxynitrite-scavenging activity. Asian Pacific J Tropical Biomed. 2017;7:208-13.
16. Gupta M, Mazumder U, Gomathi P, Selvan VT. Anti-inflammatory evaluation of leaves of *Plumeria acuminata*. BMC Complement. Altern. Med. 2006; 6(1):1-36.
17. Sofidiya M, Odukoya O, Familoni O, Inya-Agha S. Free radical scavenging activity of some Nigerian medicinal plant extracts. Pak. J. Biol. Sci. 2006; 9(8):1438-41.
18. Jyothi D, Khanum S, Sultana R. Optimisation of microwave assisted extraction of withanolides from roots of *Ashwagandha* and its comparison with conventional extraction method. International Journal of Pharmacy and Pharmaceutical Sciences 2010;2(4): 46-50.
19. Kokate CK, Purohit AP, Gokhale SB. Pharmacognosy. 10th ed. New Delhi: Nirali Prakashan Pvt Ltd; 1998,92-4.
20. Bukhari SB, Bhangar MI, Memon S. Antioxidative activity of extracts from fenugreek seeds (*Trigonella foenum-graecum*). Pak J Anal Environ Chem. 2008;9(2):78-83.

21. Mendonca V , Jyothi D, Gajinkar V, Murthy SR. Microwave assisted extraction of phenolic compounds and investigation on antioxidant activity. Journal of Xi'an Shiyou University, Natural Science Edition 2021; 17(10): 232-239
22. Jyothi D, X Khanam S, Sultana R. Optimization of microwave assisted extraction of withanolides from roots of ashwagandha and its comparison with conventional extraction method. Int J Pharm Pharm Sci,2010; 2(4):46-50
23. Zhang HF, Yang XH, Wang Y. Microwave assisted extraction of secondary metabolites from plants: current status and future directions. Trends Food Sci.Technol. 2011;22:672–688
24. Thamizhiniyan Venkatesan, Young-Woong Choi, Young-Kyoon Kim. Impact of Different Extraction Solvents on Phenolic Content and Antioxidant Potential of *Pinus densiflora* Bark Extract. BioMed Research International, 2019; Article ID 3520675. <https://doi.org/10.1155/2019/3520675>
25. Metrouh-Amir H, Duarte CMM, Maiza F. Solvent effect on total phenolic contents, antioxidant, and antibacterial activities of *Matricaria pubescens*. Ind Crop Prod. 2015;67:249–56.
26. Leopoldini M, Marino T, Russo N, Toscano. Antioxidant properties of phenolic compounds: H-atom versus electron transfer mechanism. The Journal of Physical Chemistry A, 2004;108 (22):4916–4922.