

Herbal approaches for Alzheimer disease: A review

Abstract

"Plague of the Twenty-First Century," or "Alzheimer disease" is a neuro-degenerative disease that influenced the world's elderly population. The prevalence of Alzheimer's disease is expected to increase to one hundred thirty-five million by 2050, with no treatment(s) currently available to cure or monitor the disease's progression. The current treatment(s) have less capability to manage symptoms or delay disease development, and they can cause serious side effects. The cost of overall treatment is high for both the patients and their caregivers or family members. As an example, there is a serious requirement to find safer alternative treatments for better management of Alzheimer's disease. The various Indian herbal medicines such as *Centella asiatica*, *Celastrus paniculatus*, *Curcuma longa*, *Clitoria ternatea*, *Bacopa monnieri*, *Withania somnifera*, has been reviewed in this article. *Eclipta alba*, *Desmodium gangeticum*, *Evolvulus alsinoides*, *Moringa oleifera*, and *Convolvulus pluricaulis* are among the plants that have shown promising effect in the treatment related to cognitive disability.

Keywords: Ayurveda, herbal medicine, memory, brahmi.

Introduction

A German psychiatrist in 1906, and neuropathologist (Alois Alzheimer), first identified the neurodegenerative disease, Alzheimer's disease [1]. It causes a person's intellect and memory to deteriorate, eventually leading to the loss of their ability to carry out day-to-day tasks independently. It is the most prevalent form of dementia, accounting for 60–70% of cases [2]. The onset of the disease is associated with ageing and is seldom found in the people of young age [3]. It is generally seen in people around the age of 65 or more. Currently, there is no treatment for Alzheimer's disease, which is considered the "Plague of the Twenty-First Century" since the number of cases is probable to rise from 48 million to 135 million by 2050 [4]. To successfully control and minimise the disease's proliferation, it is essential to find a cure or successful methods of prevention.

Alzheimer disease results in neuronal damage and death due to the disruption of some essential processes such as communication in neurons and with brain cells, repair, and regeneration and metabolism. In most cases, the condition gets worse with the increase in age. In the early phase, memory-related neurons in the allocortex area of brain and hippocampus are lost. After this stage, this disease progresses to the next stages, affecting regions of the cerebral cortex responsible for thought, language, and social behaviour. This is a fatal disorder which also spreads to the other part of the brain, eventually making the patient unable to function and live independently which leads to death.

Molecular mechanisms of Alzheimer

1. **Oxidative stress:** This term describes a condition in which the oxidative stress in the body is caused by ROS (reactive oxygen species), which is particularly prevalent in disorders related to age like AD. In the early phase of AD, oxidative damage develops in the brain. It has an effect well before plaque pathology begins, as it occurs before amyloid-beta (A β) deposition and formation of intracellular

neuro-fibrillary tau protein tangles are irregular. Reactive Oxygen Species play a role in neuronal death at the cellular and tissue levels and neurodegenerative processes [5].

2. **Occurrence of A β plaques:**The degradation of the A β PP results in the formation of A β proteins. These proteins start to stick together and lead to formation of plaques between neurons at abnormally high levels. Cellular function is eventually disturbed as a result of these aggregates [6].
3. **Neurofibrillary tangles formation:** When abnormally high levels of hyperphosphorylated tau protein build up within the neurons, neurofibrillary tangles form. In healthy neurons, the tau protein is important for attaching to and stabilising microtubules. [7]. after tau detaches from microtubules, it links to other tau molecules and creates threads that connect to form tangles, which impair synaptic connection between neurons in AD [8]. The cleavage of a reversing barrier at the axon beginning section causes tau proteins in the axonal area of neurons to be missorted to the somato-dendritic section [9].
4. **Microglial cells:** Microglial cells act as brain macrophages, removing cell debris such as A β plaques, harmful chemical and damaged neurons. Furthermore, astrocytes help remove cell debris and plaques from the brain. In AD, these brain microglial cells gather around the neurons but are unable to fulfil their role of clearing debris. As a result of their accumulation, chemicals are released into the brain, causing chronic inflammation [10].
5. **Vascular causes:** Alzheimer disease can also lead to A build-up of plaque in the brain's arteries, as well as artery hardening (atherosclerosis), ministrokes, and other conditions [11], may retard flow of the blood and oxygen supply of oxygen to the brain and results in a degeneration and complicated loop i.e. a cause and consequence of Alzheimer's disease [12].
6. **Brain atrophy:** In AD, neurons are damaged and die. As a result, the neural network can break down, cause shrink in brain size [13].

The hippocampus of the medial temporal lobe, which is predominantly engaged in memory and emotion, suffers from the loss of brain function. The deterioration then accelerates impacting brain glucose metabolism and cause a decrease in neuronal processing in the parietal and temporal lobes on both sides [14]. The parietal lobe controls, self-awareness, memory retrieval, attention, sensation, and theory of mind [15–17]. The temporal lobe is involved in episodic memory, feelings, and mood [18]. As a result, memory, emotion, and perceptual control are all controlled by the brain regions afflicted by Alzheimer's disease.

Treatments available currently for Alzheimer's disease

Currently only few medications are available to cure Alzheimer's disease, and most of them cantreat the symptoms only. Symptoms such as memory loss, on the other hand, are slowed, extending life and improving quality of life for those who are affected. Donepezil, rivastigmine, and galantamine are some of the drugs that are used to treat mild to severe Alzheimer's disease. These cholinesterase inhibitors, which stops acetylcholine from being broken down found essential for memory and learning. Memantine is an NMDA antagonist that is used to treat AD [19]. Memantine and donepezil is used for this in combination. Depression, rage, restlessness, and anxiety are all indications of AD, and other drugs are used to treat them. Among these are imipramine, mirtazapine, bupropion, citalopram, sertraline and duloxetine.

Alternative therapies

As the world's population grows and people live longer, AD cases are on increasing all over the world. According to report conducted by WHO on dementia reveals that around 50 million people throughout the nation are suffering from dementia and every year new cases of 10 million are cont. adding to this no. which reflects that a person worldwide developing dementia in every three sec [21]. This fast onset of AD requires the advancement of new prevention and treatment methods for Alzheimer's disease. The new therapies are ineffective, have a high number of adverse effects, and suffer from low patient cooperation. These therapies are also costly for patients and providers, as the costs of continuing presence and hospital expenses regularly increasing year after year [22]. This necessitates the introduction of safer alternative of therapies that are being used as anti-AD therapies since ancient times and have been shown to be efficient. Ayurveda, traditional Chinese rituals, meditation, and exercise are all examples of such therapies. Many therapies focused at healing function of synapses and memory has shown to be helpful in reducing the progression of Alzheimer disease and increase effect of current therapy.

Herbal drugs

Ayurveda, a Sanskrit term that means "scripture of immortality," is the philosophy of life and health that promotes a holistic approach to care to keep the body, mind, and soul in balance. The 5 fundamental tatva of the universe: fire, space, air, earth and water [23] are the foundations of this traditional medication system. Its roots can be found in India's Indus Valley Civilization and the Indian subcontinent (about 3,000 BC).

Several herbal medicines that has been used to treat neuro-degenerative diseases all over the world. For example *Salvia officinalis* and *Salvia lavandulaefolia* are used in European regions to improve memory since 16th century [24] and found to be affected with the help of different clinical trials [25,26]. The Ayurvedic tradition has long used *Bacopa monniera* (water hyssop) to aid memory and cognitive abilities. The rejuvenative tonic *Withania somnifera* root is also mentioned in Traditional Books like Ayurveda for the cognition improvement. [27-30]. Various traditional remedies are attaining popularity as a result of their perceived efficacy, protection, and affordability. Indeed, experimental studies have only recently begun to provide indication and justification of choosing herbal therapy in cognitive disabilities.

When offered as a prophylactic treatment, numerous Indian herbal medicines that are CNS active such as *Withania somnifera*, *Celastrus paniculatus*, *Centella asiatica*, and *Bacopa monnieri* possess activity of cognitive improvement in different pharmacological experiment models of AD [31-35].

Brahmi



Fig 1: Leaves of *Bacopamonniera*

Biological source: *Bacopa monnieri* (*B. monniera*) is a minor, ever growing creeping herb with a number of branches, leaves are oblong, and flowers can be light purple or can also be white belonging to Scrophulariaceae. In India it is popularly known as Brahmi and can be used for its revitalising and nootropic properties; it also improves cognition power and intellectual abilities. It also has been utilised to cure a range of diseases by practitioners of India's traditional medicine for thousands of years.

Chemical constituent: Bacosides, which are triterpenoid saponins, are the most important chemical constituents of *B. monniera*. This plant also contains the alkaloids brahmine, nicotine, and herpestine. Bacopasides I–XII, which are new saponins, have also been discovered.




Fig2: Nicotin




Fig 3: Bacopasides I

System	Pharmacological activity	References
CNS	Anti-convulsant, Anti-depressant, Anxiolytic	[37]
Respiratory	Bronchodilator	[38]
GIT	Hepato-protective	[39]
Other	Analgesic, antioxidant, antimicrobial, anticancer.	[40]

Table1: Pharmacological activity of Brahmi

Effect of Brahmi in AD: According to a study on olfactory bulbectomized mice by Le et al., this plant helps in by protecting cholinergic nerve cells in the medial septal nucleus that extend in hippocampus of our brain. A rat model of Alzheimer's disease showed that alcoholic Brahmi extracts inhibited cholinergic neuron degeneration, implying that the medication could be used therapeutically. According to these findings, the plant extracts have a cholinergic effect comparable to existing therapies such as donepezil, rivastigmine, and galantamine [37,38]. The plant extract diminishes the deposition of A β and hippocampus damage due to stress and signalling potential therapeutic effect in Alzheimer's disease[39,40].

Gotu kala



Fig 4: Leaves of Gotu kala

Biological source: Centella asiatica, an annual plant, belonging to Apiceae that is also known as jalbrahmi or mandukparni, is found throughout India. It bears small oval fruit and possess green leaves with fan like shape and the flowers are white or light purple and can also be pink [41].

The Ayurvedic method of medicine has used the leaves of mandukparni as a memory enhancer [42]. Its use has also been recorded in traditional Chinese medicine and the African medical system. It's used to slow down the ageing process, avoid memory problems, and improve memory when combined with milk [43].

Chemical constituents: Asiaticosides, madasiatic, asiatic acid, madecassoside, and acid are the chief chemical components of C. asiatica. and centelloside, brahminoside, thankuniside, isothankuniside, Brahmoside and are among the chemical compounds extracted from Gotu kala.




Fig5: Asiatic acid




Fig 6: Madasiatic acid




Fig 7: Asiaticoside acid

System	Pharmacological activity	Reference
CNS	Neuroprotective Antidepressant Anticonvulsant Sedative Nootropic	[40]
GIT	Antiulcer Hepatoprotective antidiabetic	[40]
CVS	Cardioprotective	[40]
Immune system	Immunostimulant, antiviral	[40]
General	Antibacterial, antifungal, insecticide, antitumor, anti- psoriatic.	[40]

Table 2: Pharmacological activity of Gotu kala.

Effect of Gotu kala in AD: Due to its high antioxidant properties, this neurodegenerative adaptogen aids in the prevention of cognitive impairments and the alleviation of oxidative stress associated with AD [44]. It's a better alternative to AChE inhibitors, which is used as the 1st-line therapy for AD[45]. It can also help lower phospholipase A2 levels. This herb's antioxidant activity may affect a natural process that contributes to dementia, mitochondrial dysfunction[46]. Its active ingredients are saponins and Asiatic side derivatives, which substantially reduce death of cell caused by H₂O₂, lower the level of free radical, and prevent neural cell death mediated by A β (in vitro). As a result, it may be useful in preventing and treating A β toxicity as well as AD [47, 48].

A study of 28 stable elderly people found that Gotu kola extracts could help with mood disorders and age-related cognitive decline [49]. It's generally safe to use; however, over-dose can cause lethargy [50].

Ashwagandha



Fig 8: Leaves of ashwagandha

Biological source: *Withania somnifera* (W. somnifera) is a small woody shrub that belongs to the Solanaceae family and is widely cultivated in India. The herb is also known as Indian ginseng, winter cherry, or ashwagandha. It has greenish or yellowish flowers that are about a centimetre long [51,52].

Chemical constituents: It contains withaferins, isopelletierine, anferine, sitoindoside 7 and 8, and anferine, withanoloides are some of the most important phytoconstituents found in W. somnifera. somniferine, Withanine, cuscohygrine, withanane, pseudo-withanine, tropine, somniferine, pseudo-tropine, 3- α -gloyloxytropine, somnine, choline, and are some of the other chemical compounds [53–55].




Fig 9: Isopelletierine




Fig10: Withaferin




Fig11: Anaferine

System	Pharmacological activity	References
CNS	Anti-Parkinson Anxiolytic Antistress Antidepressant	[56]
CVS	Anti-Ischemiac	[56]
GIT	Antiulcer Nephroprotective Diabetes	[56]
Respiratory	Anti-hypoxiac	[56]
Other	Anti-inflammatory Antioxidative stress Anticancer	[56]

Table3: pharmacological activity of Ashwagandha

Effect of Ashwagandha in AD:the roots helps in the suppression of the nuclear factor (NF-B), and regulates the expression of genes involved in inflammatory mediators and oxidative stress parameters and thereby

producing anti-inflammatory and antioxidative effect by preventing the development of A β . It helps in the regeneration of synaptic function and the decrease apoptosis. It encourages nuclear factor related to E2-related factor 2 (Nrf2) to migrate to the nucleus, resulting in an increase in antioxidant enzyme expression and neuroprotective proteins like haeme oxygenase-1, demonstrating antioxidant effects that are beneficial in the treatment and prevention of AD [57,58].

Turmeric



Fig 12: leaves of Turmeric

Biological source: *Curcuma longa* (*C. longa*) Linn also known as turmeric, is a perennial plant in the Zingiberaceae family. It is derived from the plant's rhizome and is widely used as a food flavouring and colouring agent in India. In South and Southeast Asia, it is grown for commercial purposes.[59-61]

Chemical constituents: Curcuminoids, which include curcumin (diferuloyl methane), demethoxycurcumin, and bisdemethoxycurcumin, are the major chemical constituents of turmeric. Alpha- and beta-tumerone, artumerone, alpha- and γ -atlantone, curlone, zingiberene, and curcumol are some of the other chemical constituents found in the plant [62].




Fig 13: Curcumin




Fig14: Demethoxycurcumin




Fig15: Bisdemethoxycurcumin.

System	Pharmacological activity	References
CNS	Neuroprotective	[63]
GIT	Antihepatotoxic Antidiabetic Antidiarrheal carminative	[63]
Urinary system	Diuretic	[63]
CVS	Anti-thrombolytic Hypotensive	[63]
Immune system	Antiviral, antirheumatic,	[63]
Other	Antimicrobial Antioxidant Larvicidal Antiproliferative Antivenomous Anti-tyrosinase	[63]

Table 4: Pharmacological activity of turmeric

Effect of Turmeric in AD: Curcumin has anti-inflammatory properties by inhibiting the development of TNF- α , IL-1 β , and further pro-inflammatory cytokines in astrocytes and microglia, such as IL-8, MIP1 β , and MCP-1. By modifying the Nrf2-Keap1 (Kelch-like ECH-associated protein 1) pathway, the oxidative damage is reduced and improves functions that are cognitive, particularly in the case of ageing. By modulating the Nrf2-Keap1 (Kelch-like ECH-associated protein 1) pathway, it reduces oxidative damage and improves cognitive function, particularly in the case of ageing [64]. When curcumin binds to Keap1, Nrf2 attaches to Keap1 and present in the cytoplasm is released.

A study found that daily curry eaters beat those who had never or only sometimes eaten it on a typical Montreal Cognitive Assessment (Mini-Mental State Examination, MMSE) [65]. It can also limit microglia development, lower the level of cholesterol, inhibit AChE, modulate the signalling pathway of insulin, prevent brain dysregulation induced by bio-metal poisoning, and inhibit acetylcholinesterase [66,67].

The extensive consumption of turmeric in the Indian population has also been linked to lower rates of AD, implying a neuroprotective effect for haldi [68]. It is also used in food preparation and is completely healthy, along with the exception of diarrhoea, nausea, dizziness, or stomach upset if eaten in large amounts.

Shankpushpi



Fig 16: leaves and flower of shankpushpi

Biological source: *Convolvulus pluricaulis* is a species of *Convolvulus* (*C. pluricaulis*) Choisy is a perennial, wild, prostrate herb that is innate to Northern India and belongs to the *Convolvulaceae* family. In the ancient system of medicine, it also known as shankpushpi and have activity as a tonic to boost memory and brainpower [69].

Chemical constituents: It contains Alkaloids that is shankpushpine and convolamine, volatile oils, flavonoid-kampferol, phytosterol, amino acids, fatty acids, scopoletin, and beta-sitosterol that is SethiyaNKare the main chemical components [70].




Fig17: Convolamine




Fig18: Scopoletin

System	Pharmacological activity	References
CVS	Cardioprotective	[70]
CNS	Nootropic Inhibit stress Antidepressant	[71-74]

	Anti-convulsant Anxiolytic Sedative Inhibit Amnesia Catatonic	
GIT	Antiulcer	[74]

Table5: Pharmacological activity of Shankhpushpi

Effect of Shankhpushpi in AD: Its aqueous extracts protect a *Drosophila* model of AD against early mortality by neutralising human microtubule-associated tau protein mediated neurotoxicity. The tau protein in tauopathy is greatly reduced by Shankhpushpi extracts. increases antioxidant enzyme activity and reduces oxidative- stress mediated by tau protein [75]. The ethanolic extracts of Shankhpushpi increase neurite outgrowth, which improves memory [76]. The action of *C.pluricaulis* can be triggered by adding it to daily food. Its small leaves, flowers, and roots, as well as the paste produced from them, can be eaten on a regular basis.

DRUMSTICK TREE



Fig 19: leaves of drumstick tree

Biological source: *Moringa oleifera* (*M. oleifera*), a member from the Moringaceae family, is the utmost widely distributed plant. It's commonly referred to as a drumstick. This plant is instinctive to India, and trees can grow up to height of 10 metres in length.

Chemical constituents:It contains Vitamins i.e.vitamin A and C, alkaloids, isothiocyanates, tannins, and saponins and polyphenols i.e. flavonoids, chlorogenic acid,glucosinolates, and phenolic acids are the main chemical constituents in *M. oleifera* [77].




Fig20: Chlorogenic acid




Fig21: Retinol




Fig22: Glucosinolates

System	Pharmacological activity	References
CNS	Nootropic Memory Enhancer	[78,79]
Other	Anti-inflammatory Anti-diabetic Inhibit obesity Anticancer	[80-82]

Table6: Pharmacological effect of Drumstick

Effect of drumstick tree in ADS: It reversed the effects of colchicine on NE, 5-HT, and DA levels in the brain. Its antioxidant effect has been linked to improvements in learning and memory. Other studies have also shown that *M. oleifera* protects in contrast to memory loss in laboratory models for dementia.

Conclusion

Alternative medicine is used in the treatment of Alzheimer's disease from ancient times, and many extractions of medicinal plants and herbal formulae showed potential. Medicinal plants provide fertile ground for novel drug development due to the availability of diverse chemical ingredients and their capacity to work on a different type of biological targets. The no. of persons suffering by age-related diseases such as Alzheimer's disease will substantially increase as the world's population grows. The number of AD cases is expected to increase to 135 Million by 2050 in the ageing population. As a result, advanced research and development of effective preventive and therapeutic methods are gaining increasing interest. This paper has mentioned some of herbs related to the AD and their effect. However, more work is still needed in order to convert this potential into actual treatment. Multi-centre clinical trials should be done to confirm the effectiveness of the herbal medications alone or in the form of formulations for the treatment of AD.

NOTE:

The study highlights the efficacy of "Ayurveda " which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

References

1. Berchtold NC, Cotman CW (1998). Evolution in the conceptualization of dementia and Alzheimer's disease: Greco-Roman period to the 1960s. *Neurobiol Aging* 19, 173-189.
2. World Health Organization (2012). Dementia: A public health priority, <https://www.who.int/mentalhealth/publications/dementia-report-2012/en/>.
3. Guerreiro R, Bras J (2015). The age factor in Alzheimer's disease. *Genome Med* 7, 1-3.
4. Reddy VS, Bukke S, Dutt N, Rana P, Pandey AK (2017). A systematic review and meta-analysis of the circulatory, erythrocellular and CSF selenium levels in Alzheimer's disease: A metal meta-analysis (AMMA study-I). *J Trace Elem Med Biol* 42, 68-75.
5. Evans PH, Klinowski J, Yano E, Urano N (1989). Alzheimer's disease: A pathogenic role for aluminosilicate-induced phagocytic free radicals. *Free Radic Res* 6, 317-321.
6. Greenfield JP, Tsai J, Gouras GK, Hai B, Thinakaran G, Checler F, Sisodia SS, Greengard P, Xu H (1999). Endoplasmic reticulum and trans-Golgi network generate distinct populations of Alzheimer - amyloid peptides. *Proc Natl AcadSciUSA* 96, 742-747.
7. Binder LI, Frankfurter A, Rebhun LI (1985). The distribution of tau in the mammalian central nervous system. *J Cell Biol* 101, 1371-1378.
8. Ballatore C, Lee VM-Y, Trojanowski JQ (2007). Tau-mediated neurodegeneration in Alzheimer's disease and related disorders. *Nat Rev Neurosci* 8, 663-672.
9. Li X, Kumar Y, Zempel H, Mandelkow E-M, Biernat J, Mandelkow E (2011). Novel diffusion barrier for axonal retention of Tau in neurons and its failure in neurodegeneration. *EMBO J* 30, 4825-4837.
10. Katsumoto A, Takeuchi H, Takahashi K, Tanaka F (2018). Microglia in Alzheimer's disease: Risk factors and inflammation. *Front Neurol* 9, 978.
11. Gupta A, Iadecola C (2015). Impaired A clearance: A potential link between atherosclerosis and Alzheimer's disease. *Front Aging Neurosci* 7, 115.
12. Canobbio I, Abubaker AA, Visconte C, Torti M, Pula G (2015). Role of amyloid peptides in vascular dysfunction and platelet dysregulation in Alzheimer's disease. *Front Cell Neurosci* 9, 65.
13. National Institute on Aging. What Happens to the Brain in Alzheimer's Disease? <https://www.nia.nih.gov/health/what-happens-brain-alzheimers-disease>
14. Newberg AB, Serruya M, Wintering N, Moss AS, Reibel D, Monti DA (2014). Meditation and neurodegenerative diseases. *Ann N Y Acad Sci* 1307, 112-123.
15. Mesulam MM(1983). The functional anatomy and hemispheric specialization for directed attention. The role of the parietal lobe and its connectivity. *Trends Neurosci* 6, 384-387.
16. Saxe R, Kanwisher N (2003). People thinking about thinking people. The role of the temporo-parietal junction in "theory of mind". *Neuroimage* 19, 1835-1842.
17. Wagner AD, Shannon BJ, Kahn I, Buckner RL (2005). Parietal lobe contributions to episodic memory retrieval. *Trends Cogn Sci* 9, 445-453.

18. Campbell S, Macqueen G (2004). The role of the hippocampus in the pathophysiology of major depression. *J Psychiatry Neurosci* 29, 417-426.
19. Cummings JL, Morstorf T, Zhong K (2014). Alzheimer's disease drug-development pipeline: Few candidates, frequent failures. *Alzheimers Res Ther* 6, 37. [20] National Institute on Aging. How Is Alzheimer's Disease Treated? <https://www.nia.nih.gov/health/howalzheimerers-disease-treated>
20. Alzheimer's Disease International. Dementia statistics. <https://www.alz.co.uk/research/statistics>
21. Alzheimer's Association (2019). Alzheimer's Disease Facts And Figures. <https://www.alz.org/med>
22. Le XT, Pham HTN, Do PT, Fujiwara H, Tanaka K, Li F, Van Nguyen T, Nguyen KM, Matsumoto K (2013). Bacopa monnieri ameliorates memory deficits in olfactory bulbectomized mice: Possible involvement of glutamatergic and cholinergic systems. *Neurochem Res* 38, 2201- 2215.
23. Meena DK, Upadhyay D, Singh R, Dwibedy BK (2015). A critical review of fundamental principles of Ayurveda. *Int Ayurvedic Med J* 3, 2075-2083.
24. Perry, N.; Court, G.; Bidet, N.; Court, J.; Perry,(1996). E. European Herbs with Cholinergic Activities: Potential in Dementia Therapy. *Int. J. Geriatr. Psychiatry*, 11, 1063–1069.
25. Perry, N.S.; Bollen, C.; Perry, E.K.; Ballard,(2003). C. Salvia for dementia therapy: Review of pharmacological activity and pilot tolerability clinical trial. *Pharm. Biochem. Behav.*, 75, 651–659.
26. Akhondzadeh, S.; Noroozian, M.; Mohammadi, M.; Ohadinia, S.; Jamshidi, A.H.; Khani,(2003). M. Salvia officinalis extract in the treatment of patients with mild to moderate Alzheimer's disease, a double blind, randomized and placebo-controlled trial. *J. Clin. Pharm. Ther.*, 28, 53–59.
27. Manyam,(1999) B.V. Dementia in Ayurveda. *J. Altern. Complement. Med.*, 5, 81–88.
28. Kuboyama, T.; Tohda, C.; Komatsu, K.(2005). Neuritic regeneration and synaptic reconstruction induced by withanolide A. *Br. J. Pharm.*, 144, 961–971. .
29. Abourjaily,(2001). P. American Herbal Pharmacopoeia and Therapeutic Compendium (A botanical supplement monograph series). *Nutr. Clin. Care*, 4, 221–222. ..
30. Holcomb, L.A.; Dhanasekaran, M.; Hitt, A.R.; Young, K.A.; Riggs, M.; Manyam,(2006).B.V. Bacopa monniera extract reduces amyloid levels in PSAPP mice. *J. Alzheimer's Dis.*, 9, 243–251. .
31. Kuboyama, T.; Tohda, C.; Komatsu,(2006).K. Withanoside IV and its active metabolite, sominone, attenuate Ab (25–35)-induced neurodegeneration. *Eur. J. Neurosci.*, 23, 1417–1426. .
32. Kumar, M.; Gupta, (2002).Y. Antioxidant property of *Celastrus paniculatus* Willd: A possible mechanism in enhancing cognition. *Phytomedicine*, 9, 302–311. .
33. Shinomol, G.K.; Bharath,(2011). M.M. Exploring the Role of “Brahmi” (*Bacopa monnieri* and *Centella asiatica*) in Brain Function and Therapy. *Recent Pat. Endocr. Metab. Immune Drug Discov.*, 5, 33–49. .

34. Uabundit, N.; Wattanathorn, J.; Mucimapura, S.; Ingkaninan, (2010).K. Cognitive enhancement and neuroprotective effects of Bacopa monnieri in Alzheimer's disease model. *J. Ethnopharmacol.*, 127, 26–31. .
35. Kumar, M.H.V.; Gupta, (2003). Y. Effect of Centella asiatica on cognition and oxidative stress in an intracerebroventricular streptozotocin model of Alzheimer's disease in rats. *Clin. Exp. Pharm. Physiol.*, 30, 336–342.
36. Aguiar, S.; Borowski,(2013). T. Neuropharmacological Review of the Nootropic Herb Bacopa monnieri. *Rejuvenation Res.* 2013, 16, 313–326. .
37. Le XT, Pham HTN, Do PT, Fujiwara H, Tanaka K, Li F, Van Nguyen T, Nguyen KM, Matsumoto K (2013). Bacopa monnieri ameliorates memory deficits in olfactory bulbectomized mice: Possible involvement of glutamatergic and cholinergic systems. *Neurochem Res* 38, 2201- 2215.
38. Uabundit N, Wattanathorn J, Mucimapura S, Ingkaninan K (2010). Cognitive enhancement and neuroprotective effects of Bacopa monnieri in Alzheimer's disease model. *J Ethnopharmacol* 127, 26-31.
39. Mathew M, Subramanian S (2012). Evaluation of the anti-amyloidogenic potential of nootropic herbal extracts in vitro. *Int J Pharm Sci Res* 31, 4276-4280.
40. Holcomb LA, Dhanasekaran M, Hitt AR, Young KA, Riggs M, Manyam B V (2006). Bacopa monniera extract reduces amyloid levels in PSAPP mice. *J Alzheimers Dis* 9, 243-251.
41. Gohil, K.J.; Patel, J.A.; Gajjar,(2010). A.K. Pharmacological review on Centella asiatica: A potential herbal cure-all. *Indian J. Pharm. Sci.*, 72, 546–556. .
42. Rao, K.G.M.; Rao, S.M.; Rao, (2006) S.G. Centella asiatica (L.) Leaf Extract Treatment during the Growth Spurt Period Enhances Hippocampal CA3 Neuronal Dendritic Arborization in Rats. *Evid. Based Complement. Altern. Med.*, 3, 349–357.
43. Manyam, (1999). B.V. Dementia in Ayurveda. *J. Altern. Complement. Med.*, 5, 81–88.
44. Veerendra Kumar MH, Gupta YK (2003). Effect of Centella asiatica on cognition and oxidative stress in an intracerebroventricular streptozotocin model of Alzheimer's disease in rats. *Clin Exp PharmacolPhysiol* 30, 336-342.
45. Puttarak P, Dilokthornsakul P, Saokaew S, Dhippayom T, Kongkaew C, Sruamsiri R, Chuthaputti A, Chaiyakunapruk N (2017). Effects of Centella asiatica (L.) Puttarak, P. et al. (2017) Effects of

Centella asiatica (L.) Urb. on cognitive function and mood related outcomes: A systematic review and meta-analysis. *Sci Rep* 7, 10646.

46. Prakash A, Kumar A (2013). Mitoprotective effect of *Centella asiatica* against aluminum-induced neurotoxicity in rats: Possible relevance to its anti-oxidant and antiapoptosis mechanism. *Neurol Sci* 34, 1403-1409.
47. Gray N, Morre J, Kelley J, Maier C, Stevens F, Quinn J, Soumyanath A (2013). *Centella asiatica* protects against the toxic effects of intracellular beta-amyloid accumulation. *Planta Med* 79, PH6.
48. Gray NE, Sampath H, Zweig JA, Quinn JF, Soumyanath A (2015). *Centella asiatica* attenuates amyloid-induced oxidative stress and mitochondrial dysfunction. *J Alzheimers Dis* 45, 933-946.
49. Wattanathorn J, Mator L, Muchimapura S, Tongun T, Pasuriwong O, Piyawatkul N, Yimtae K, Sripanidkulchai B, Singkhoraard J (2008). Positive modulation of cognition and mood in the healthy elderly volunteer following the administration of *Centella asiatica*. *J Ethnopharmacol* 116, 325-332
50. Deshpande PO, Mohan V, Thakurdesai P (2015). Preclinical safety assessment of standardized extract of *Centella asiatica* (L.) urban leaves. *Toxicol Int* 22, 10-20.
51. Bone, (1996). K. *Clinical Applications of Ayurvedic and Chinese Herbs. Monographs for the Western Herbal Practitioner*; Phytotherapy Press: Queensland, Australia, 1996; pp. 137–141.
52. Chatterjee, A.; Pakrashi (1995). S.C. *The Treatise on Indian Medicinal Plants. Council for Scientific and Industrial Research; Publications & Information Directorate: New Delhi, India, 1995; Volume 4, pp. 208–212.*
53. Dar, N.J.; Hamid, A.; Ahmad M. (2015). Pharmacologic overview of *Withania somnifera*, the Indian Ginseng. *Cell. Mol. Life Sci.* 2015, 72, 4445–4460. . [PubMed]
54. Mirjalili, M.H.; Moyano, E.; Bonfill, M.; Cusido, R.M.; Palazón J. (2009). Steroidal Lactones from *Withania somnifera*, an Ancient Plant for Novel Medicine. *Molecules* 2009, 14, 2373–2393. . [PubMed]
55. Mishra, L.C.; Singh, B.B.; Dagenais (2000). S. Scientific basis for the therapeutic use of *Withania somnifera* (ashwagandha): A review. *Altern. Med. Rev.* 2000, 5, 334–346.
56. Kumar, V.; Dey, A.; Hadimani, M.B.; Marcović, T. (2015). Emerald, M. Chemistry and pharmacology of *Withania somnifera*: An update. *Tang (Humanit. Med.)* 2015,

57. Sandhir R, Sood A (2017). Neuroprotective potential of withania somnifera (ashwagandha) in neurological conditions. In *Science of Ashwagandha: Preventive and Therapeutic Potentials*. Springer International Publishing, pp. 373-387.
58. Sun GY, Li R, Cui J, Hannink M, Gu Z, Fritsche KL, Lubahn DB, Simonyi A (2016). Withania somnifera and its withanolides attenuate oxidative and inflammatory responses and up-regulate antioxidant responses in BV-2 microglial cells. *Neuromolecular Med* 18, 241- 252.
59. Majeed, M.; Badmaev, V.; Murray(1996). *F. Turmeric and the Healing Curcuminoids*; Keats Publishing, Inc.: New Canaan, CT, USA.
60. Perrone, D.; Ardito, F.; Giannatempo, G.; Dioguardi, M.; Troiano, G.; Russo, L.L.; De Lillo, A.; Laino, L.; Muzio(2015). L.L. Biological and therapeutic activities, and anticancer properties of curcumin. *Exp. Med.*, 10, 1615–1623.
61. Sharifi-Rad, J.; El Rayess, Y.; Rizk, A.A.; Sadaka, C.; Zgheib, R.; Zam, W.; Sestito, S.; Rapposelli, S.; Neffe-Skocińska, K.; Zielińska, D.; et al. (2020). Turmeric and Its Major Compound Curcumin on Health: Bioactive Effects and Safety Profiles for Food, Pharmaceutical, Biotechnological and Medicinal Applications. *Front. Pharm.* 2020, 11.
62. Goel A, Kunnumakkara AB, Aggarwal BB (2008). Curcumin as “Curecumin”: From kitchen to clinic. *BiochemPharmacol* 75, 787-809
63. Ng T-P, Chiam P-C, Lee T, Chua H-C, Lim L, Kua E-H (2006). Curry consumption and cognitive function in the elderly. *Am J Epidemiol* 164, 898-906.
64. Mishra S, Palanivelu K (2008). The effect of curcumin (turmeric) on Alzheimer’s disease: An overview. *Ann Indian AcadNeurol* 11, 13-19.
65. Tang M, Taghibiglou C (2017). The Mechanisms of action of curcumin in Alzheimer’s disease. *J Alzheimers Dis* 58, 1003-1016.
66. Shoba G, Joy D, Joseph T, Majeed M, Rajendran R, Srinivas PSSR (1998). Influence of piperine on the pharmacokinetics of curcumin in animals and human volunteers. *Planta Med* 64, 353-356.
67. Adams, M.; Gmünder, F.; Hamburger(2007), M. Plants traditionally used in age related brain disorders—A survey of ethnobotanical literature. *J. Ethnopharmacol.*, 113, 363–381.
68. Sethiya, N.K.(2009). An update on Shankhpushpi, a cognition-boosting Ayurvedic medicine. *J. Chin. Integr. Med.* 2009, 7, 1001–1022.
69. Ahmad, S.; Zafar(2007). R.-U.; Shahid, M. Anticonvulsant potential of callus cultures of *Convolvulus microphyllus*Sieb. *Orient. Pharm. Exp. Med.*, 7, 46–50.
70. Dhingra, D.(2007).Valecha, R. Evaluation of the antidepressant-like activity of *Convolvulus pluricaulischoisy* in the mouse forced swim and tail suspension tests. *Med. Sci. Monit.*, 13, BR155–BR161.

71. Dubey, G.P.; Pathak, S.R.(1994). Gupta, B.S. Combined effect of Brahmi (*Bacopa monniera*) and Shankhpushpi (*Convolvulus pluricaulis*) on cognitive functions. *Pharmacopsychocol* 1994, 3, 249–251.
72. Sharma, K.; Arora, V.; Rana, A.C.(2009). Bhatnagar, M. Anxiolytic effect of *Convolvulus pluricaulis* petals on elevated plus maze model of anxiety in mice. *J. Herb. Med. Toxicol.* 2009, 1, 41–46.
73. Kizhakke PA, Olakkaran S, Antony A, Tilagul KS, Hunasanahally PG (2019). *Convolvulus pluricaulis* (Shankhpushpi) ameliorates human microtubule-associated protein tau (hMAP τ) induced neurotoxicity in Alzheimer's disease *Drosophila* model. *J Chem Neuroanat* 95, 115-122.
74. Rai KS, Murthy KD, Karanth KS, Nalini K, Rao MS, Srinivasan KK (2002). Clitoriaternatea root extract enhances acetylcholine content in rat hippocampus. *Fitoterapia* 73, 685-689.
75. Vergara-Jimenez, M.; AlMatrafi, M.M.(2017). Fernandez, M.L. Bioactive Components in *Moringa Oleifera* Leaves Protect against Chronic Disease. *Antioxidants*, 6, 91.
76. Caceres, A.; Saravia, A.; Rizzo, S.; Zabala, L.; De Leon, E.(1992). Nave, F. Pharmacologic properties of *Moringa oleifera*. 2: Screening for antispasmodic, antiinflammatory and diuretic activity. *J. Ethnopharmacol.*, 36, 233–237.
77. Faizi, S.; Siddiqui, B.S.; Saleem, R.; Siddiqui, S.; Aftab, K.(1995). Gilani, A.-U.-H. Fully acetylated carbamate and hypotensive thiocarbamate glycosides from *Moringa oleifera*. *Phytochemistry* 1995, 38, 957–963.
78. Ghasi, S.; Nwobodo, E.(2000)Ofili, J. Hypocholesterolemic effects of crude extract of leaf of *Moringa oleifera* Lam in high-fat diet fed wistar rats. *J. Ethnopharmacol.*, 69, 21–25.
79. Mohan, M.; Kaul, N.; Punekar, A.; Girnar, R.; Junnare, P.; Patil(2005). L. Nootropic activity of *Moringa oleifera* leaves. *J. Nat. Remed.* 2005, 5, 59–62.
80. Verma, A.R.; Vijayakumar, M.; Mathela, C.S.; Rao(2009), C.V. In vitro and in vivo antioxidant properties of different fractions of *Moringa oleifera* leaves. *Food Chem. Toxicol.* 2009, 47, 2196–2201.
81. Mehla J et al ., (2020). Indian formulation for Alzheimer's disease, from Traditional Knowledge to scientific Assessment. *Brain science*.