

PREPARATION OF MOUTHWASH USING BLUE TEA AND ITS ANTIOXIDANT ACTIVITY

Running title: Blue tea mediated mouthwash and its antioxidant activity

ABSTRACT:

Introduction: *Clitoria ternatea* also known as the "blue pea," is a perennial twinning herbaceous plant in the Fabaceae family. The medicinal properties of this plant have been scientifically validated, especially at an international level, and it has been recorded to have a variety of biological activities, including antioxidant, antidiabetic, and hepatoprotective properties.

Aim: The aim of this study was to evaluate the antioxidant property of blue tea extract based mouthwash.

Materials and Methods: The study was performed as an in vitro study under a laboratory setting. Synthesis of the mouthwash was performed using dried leaves of the butter pea plant subsequently the mouthwash was tested for its antioxidant activity and was assessed via the DPPH assay. The obtained values were compared with that of a known standard. The obtained data was sorted in Microsoft Excel and statistically analyzed using Statistical Package for Social Sciences (SPSS Software, Version 23.0). Unpaired t test was done for the comparison of blue tea extract with the standard and one way ANOVA followed by Tukey's post hoc test was done to compare various concentrations with standard.

Results: The blue tea extract mouthwash was tested with 10 μ L initially and then was increased to 20 μ L, 30 μ L, 40 μ L and 50 μ L. Based on DPPH assay, as the concentration of the extract solution was increased by 10 μ L each time tested, their antioxidant activity also increased over a scale. It was seen that the blue tea extract mouthwash had significantly less antioxidant capacity when compared to the standard ($p < 0.05$).

Conclusion: The blue tea extract mouthwash is found to be a potent antioxidant agent however the antioxidant property was less effective when compared to the standard.

Keywords: Blue tea, Antioxidant, Mouthwash, Green synthesis, Innovative.

INTRODUCTION:

Natural products have always been the go-to option for consumers all around the world in all sections of need let it be pharmaceuticals or daily use products.(1-2) The hype around such products are of the highest order due(3-4) to its properties which have less or no side effects when compared to the current chemical products which have at least one or more agents ,which if taken at higher doses can turn lethal.(5-7)

The 'Butterfly Pea' (*Clitoria ternatea*), also known as Asian Pigeon Wings, Blue Bell Vine, Blue Pea, Kordofan Pea, and Darwin Pea(8), is an amazing brain power enhancing herb native to tropical equatorial Asia. *Clitoria Ternatea* has been used as a memory enhancer, brain booster, anti-stress, and calming agent in traditional Chinese(9) and Ayurvedic medicine for centuries. Butterfly Pea has commonly been used as a vegetable in cooking, to colour deserts, or to make a vividly bright coloured tea because of its luminous indigo colour. Butterfly Pea(10) is high in health-promoting antioxidants, flavonoids, and peptides, and has shown promise in animal studies as a natural remedy for a variety of ailments. Because of the effects of the flavonoid quercetin on skin and hair, Butterfly Pea has been used in a variety of beauty products. Many health benefits have been attributed to Butterfly Pea(11) in both Chinese and Ayurvedic medicine, many of which have been confirmed by recent clinical studies. The herb has shown efficacy in studies for its brain-boosting properties and a wide range(12) of neurological benefits, including depression, anxiety, and fever reduction.

Furthermore, the medicinal properties of this plant have been scientifically validated, especially at an international level, and it has been recorded to have a variety of biological(13,14) activities, including antioxidant, antidiabetic, and hepatoprotective properties. Medicinal plants are a good natural source for finding cures for existing non-communicable diseases around the world. Furthermore, numerous studies have shown that antioxidant-rich foods are essential in the prevention and management of a variety of oxidative stress-related chronic diseases. Scavenging of free radicals(15), inhibition of oxidative enzymes, chelation of metal ions, and acting as antioxidant enzyme cofactors are all mechanisms that antioxidants use to manage oxidative stress in biological systems. Since oxidative stress plays a major role in the development of diabetes mellitus and its complications, diets rich in antioxidants may be a better alternative source for

managing diabetes mellitus. And since Blue pea has proven to have its antioxidant properties, using it for various other areas can also be proven to be helpful and efficient. Mouthwashes are known for keeping the breath fresh and preventing the build-up of bacteria(20–22) in between the teeth. Mouthwashes come in a variety of formulations, including everyday-care formulas, alcohol-free versions, and herbal blends, all of which are intended to encourage oral health, and good hygiene.

Since use of herbal mouthwashes have been in practice since a long time, introduction of the same but with blue tea extracts will only be addressed if it shows properties better than those currently in market. Constant check must be kept to ensure product value to make it more than the other mouthwashes being preferred. The extract can not only be incorporated in mouthwashes alone but also in topical gels which can be used at home or even as mouth sprays.

The aim of this study is to bring out a natural product i.e blue pea flower known to everyone but in form of a mouthrinse or even in form of topical gels. This would help in widening the range of areas where the blue tea extract can be used and also reach out to a wider population.

Our team has extensive knowledge and research experience that has translated into high quality publications(23–35),(36–40)(41)(42). In this context, the aim of this study was to prepare a mouthwash using blue tea and study its antioxidant properties.

MATERIALS AND METHODS:

The study was performed in the Blue Lab, Saveetha Dental College and Hospitals, Chennai. Ethical clearance required for the study was obtained from the institutional committee. The study was performed from January 2021 to February 2021.

Preparation of the mouthwash:

5g of the dried leaves of *Clitoriaternatea* (blue tea) plant was taken and 100ml of water was added and boiled until the colour changed into dark blue. The solution was then taken and filtered and concentrated to 10%. In a separate beaker, 0.3 grams sucrose was taken as the sweetening agent, 0.001 grams of preservative and 0.01 grams of sodium lauryl sulfate were

taken as the foaming agent. To this mix, 1ml of the 10% concentrated blue tea extract and 10 ml of distilled water was added and mixed well to obtain the blue tea extract mouthwash.

Antioxidant Activity:

The 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay was used to test the antioxidant activity of the blue tea extract mouthwash. Different concentrations (10 μ l, 20 μ l, 30 μ l, 40 μ l and 50 μ l) of mouthwash having the blue tea extract was mixed with 1 ml of 0.1 mM DPPH in methanol solution and 450 μ l of 50 mM TrisHCl buffer (pH 7.4) and incubated for 30 minutes. After incubation, the reduction in the number of DPPH free radicals was measured based on the wavelength at 517 nm. BHT (butylated hydroxytoluene) was used as control. The percentage inhibition was calculated from the following equation:

$$\% \text{ inhibition} = \frac{[\text{Absorbance of control} - \text{Absorbance of sample}] \times 100}{\text{Absorbance of Control}}$$

Absorbance of Control

Statistical analysis:

The obtained data was sorted in Microsoft Excel and statistically analyzed using Statistical Package for Social Sciences (SPSS Software, Version 23.0). Unpaired t test was done for the comparison of the standard and one way ANOVA followed by Tukey's post hoc test. Results depicted in graphs and tabulations.

RESULTS:

The blue tea extract mouthwash was tested with 10 μ L initially and then was increased to 20 μ L, 30 μ L, 40 μ L and 50 μ L. Based on DPPH assay, as the concentration of the extract solution was increased by 10 μ L each time tested, their antioxidant activity also increased over a scale. It was seen that the blue tea extract mouthwash had significantly less antioxidant capacity when compared to the standard ($p < 0.05$). (Figure 1)

One way ANOVA followed by post hoc analysis, revealed that there was a concentration dependent increase in the antioxidant capacity of the extract mouthwash.

Antioxidant capacity was least at 10 μ L concentration and it was more at 20 μ L which was statistically significant as observed in this study. The antioxidant activity was found to be increased significantly with increase in concentration of the extract mouthwash which was statistically significant, however it was not as effective as the standard. (Table 1)




Figure 1: Graph depicts the antioxidant activity of blue tea extract mouthwash compared to the standard. The Y axis depicts the mean absorbance value of the samples and the X axis denotes the various concentrations of the standard and samples. The blue bar depicts the standard while the green bar depicts the blue tea extract based mouthwash. Blue tea extract based mouthwash had less antioxidant activity as compared to the standard at all concentrations, however there was a significant increase in absorbance values with increase in concentration, $p < 0.05$, statistically significant (Unpaired t test, error percentage has been set upto 95%).

Table 1: Table depicts the statistical comparison of antioxidant activity at various concentrations using one way ANOVA analysis followed by post hoc. ($p < 0.05$)

Concentration (I)	Concentration (J)	Significance (p value)
10 μ L	20 μ L	0.000*
	30 μ L	0.000*
	40 μ L	0.000*
	50 μ L	0.000*
20 μ L	30 μ L	0.000*
	40 μ L	0.000*

	50 μ L	0.000*
30 μ L	40 μ L	0.000*
	50 μ L	0.000*
40 μ L	50 μ L	0.000*

*(p<0.05)

DISCUSSION:

The purpose of this study was to evaluate whether the prepared blue tea extract mouthwash had antioxidant activity and to compare its efficacy to existing standards.

The antioxidant activity of five different concentrations of plant samples was analyzed and compared to the very same concentration of standard drugs. The activity of the formulated blue tea mouthwash was convincingly demonstrated in the concentration-dependent assay. The results revealed that the sample mouthwash had antioxidant properties. Despite the fact that the blue tea extract mouthwash demonstrated a concentration-dependent increase from lower to higher concentrations at all levels, it was significantly less effective than the standard. Although the antioxidant activity of the blue tea extract mouthwash was not superior to that of the standard, the antioxidant activity of the blue tea extract mouthwash increased significantly with increasing concentrations from 10 μ L to 20 μ L, and the effectiveness of the standard mouthwash was almost the same.

The *C. ternatea* extract mouthwash may have had more promising results if the components were used to its full potential rather than just depending on concentration levels. With increase in levels of certain components such as triterpenoids, flavanol glycosides and anthocyanins at varied scales may have brought a different result which could have been satisfactory. Studies show that certain components when extracted from the blue pea flower petals have eye opening results. The blue flower petals of *C. ternatea* contain quercetin glycosides and ternatin anthocyanins, which may be useful in the development of medications or nutraceuticals to protect against chronic inflammatory illnesses(11) by reducing the excessive synthesis of pro-inflammatory mediators by macrophage cells.

Certain authors have stated that the addition of the blue pea flower extract into daily consumed food items like yogurt has significantly increased the antioxidant properties of the yoghurt(43). The flower extract of *C. ternatea* serves as a direct antioxidant that may protect against free radicals produced by external or endogenous biological events(44). It could be because of increased concentration of the extract and may also be due to certain chemical components additions which can increase the efficacy of the blue pea extract. Since the *C. ternatea* plant is widely grown in various habitats, the extraction of its components which are important are not much of a tedious task. A study shows that it also helps in regeneration of the nitrogen level in the soil and also as a great source of novel phytochemicals. Butelase-1, a biotechnological tool(45) for peptide ligation and cyclization generated from *C. ternatea* pods, is also generating a lot of discussion.

The plant is not under observation for its antioxidant properties alone but also for its antimicrobial properties. Antimicrobial activity against *Staphylococcus aureus*, *Bacillus cereus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Streptococcus agalactiae*, and *Aeromonas hydrophila* has been found in all sections of *C. ternatea*(12). Authors of various studies have shown that the *C. ternatea* extracts have antibacterial, antipyretic, anti-inflammatory, analgesic, diuretic, local anaesthetic, antidiabetic, insecticidal, blood platelet aggregation-inhibiting, and vascular smooth muscle relaxing capabilities, among other things.

This plant has a long history of usage in traditional Ayurvedic medicine for a variety of ailments, and scientific research has(46) proved so in the current times.

Previous studies(47) done to check for various characteristics of the *C. ternatea* flower extract show that when used as a commercial product and also as a folkloric medicine(48), it does show promising results of antioxidant activity but not as much as the other commercial products which were currently in use. The results of the present study are in accordance with the previous studies. However, more clinical trials need to be conducted and hence can be used for medicinal purposes.

CONCLUSION:

The blue tea extract mouthwash is found to be a potent antioxidant agent however the antioxidant property was less effective when compared to the standard(49)(50)(51)(52)(53)(54)(55)(56)(57)(58)(59) ((59,60)(61)(62)

NOTE:

The study highlights the efficacy of "ayurveda" which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

REFERENCES:

1. Sagana M, Rajasekar A, Rajeshkumar S. Antifungal Activity Of Grape Seed Extract Mediated Zinc Oxide Nanoparticles - An Invitro Study. *Plant Cell Biotechnology And Molecular Biology*. 2020 Aug 25;21(29-30):14–20.
2. Yuvashree CS, Rajasekar A, Rajeshkumar S. Cytotoxic effect of titanium dioxide nanoparticles synthesized using grape seed extract: an in vitro study. *Plant cell biotechnology and molecular biology*. 2020 Aug 26;21(31-32):120–6.
3. Shivani N, Rajasekar A, Rajeshkumar S. Antifungal Activity Of Grape Seed Extract Mediated Titanium Oxide Nanoparticles Against *Candida albicans*: An in vitro study. *Plant*

cell biotechnology and molecular biology. 2020 Aug 26;21(35-36):8–15.

4. Devi BV, Rajasekar A, Rajeshkumar S. Antiinflammatory activity of zinc oxide nanoparticles synthesised using grape seed extract: an in vitro study. *Plant cell biotechnology and molecular biology*. 2020 Aug 26;21(33-34):6–16.
5. Pereira WD, Rajasekar A, Rajeshkumar S. Green synthesis of selenium nanoparticles (senps) using aqueous extract of clove and cinnamon. *Plant cell biotechnology and molecular biology*. 2020 Aug 25;21(29-30):85–91.
6. PRANATI, Rajasekar A, Rajeshkumar S. Anti inflammatory and cytotoxic effect of clove and cinnamon herbal formulation. *Plant cell biotechnology and molecular biology*. 2020 Aug 25;21(29-30):69–77.
7. Anjum AS, Rajasekar A, Rajeshkumar S. Synthesis And Characterization Of Grape Seed Mediated Titanium Dioxide Nanoparticles: An In Vitro Study. *Plant Cell Biotechnology And Molecular Biology*. 2020 Aug 26;21(33-34):17–23.
8. Albert H, Salt-Works Press. *Blue Tea* [Internet]. 1973. 22 p. Available from: https://books.google.com/books/about/Blue_Tea.html?hl=&id=yoAxHQAACAAJ
9. Lakshan SAT, Jayanath NY, Abeysekera WPKM, Abeysekera WKSM. A Commercial Potential Blue Pea (*L.*) Flower Extract Incorporated Beverage Having Functional Properties. *Evid Based Complement Alternat Med* [Internet]. 2019 May 20;2019:2916914. Available from: <http://dx.doi.org/10.1155/2019/2916914>
10. Gaytos CEG, Lumagbas NAA. Acceptability of Asian Blue Pea Flower (*Clitoria Ternatea*) Ice Cream [Internet]. *SSRN Electronic Journal*. Available from: <http://dx.doi.org/10.2139/ssrn.3517731>
11. Nair V, Bang WY, Schreckinger E, Andarwulan N, Cisneros-Zevallos L. Protective Role of Ternatin Anthocyanins and Quercetin Glycosides from Butterfly Pea (*Clitoria ternatea* Leguminosae) Blue Flower Petals against Lipopolysaccharide (LPS)-Induced Inflammation in Macrophage Cells. *J Agric Food Chem* [Internet]. 2015 Jul 22;63(28):6355–65. Available from: <http://dx.doi.org/10.1021/acs.jafc.5b00928>
12. Jamil N, Pa'ee F. Antimicrobial activity from leaf, flower, stem, and root of *Clitoria ternatea* – A review [Internet]. 2018. Available from: <http://dx.doi.org/10.1063/1.5050140>
13. *Clitoria ternatea* Linn [Internet]. SpringerReference. Available from: http://dx.doi.org/10.1007/springerreference_68312
14. Paranjpe P. *Indian Medicinal Plants: Forgotten Healers : a Guide to Ayurvedic Herbal Medicine with Identity, Habitat, Botany, Photochemistry, Ayurvedic Properties, Formulations & Clinical Usage* [Internet]. 2001. 316 p. Available from: https://books.google.com/books/about/Indian_Medicinal_Plants.html?hl=&id=iKVFAAAA YAAJ

15. Priya P, Elumali K, Shakila D, Geetha K, Dinesh Karthik A. Facile approach to synthesize, compared to MgO & ZnO nanoparticles by using *Clitoria ternatea*/*Tecoma castanifolia* flower [Internet]. Vol. 29, *Materials Today: Proceedings*. 2020. p. 1217–22. Available from: <http://dx.doi.org/10.1016/j.matpr.2020.05.479>
16. Oral Cancer and Mouthwash Use [Internet]. *JNCI: Journal of the National Cancer Institute*. 1983. Available from: <http://dx.doi.org/10.1093/jnci/71.6.1103>
17. Widatiningsih S, Rofiah S, Ristiyanawati. The Use of Brewed Herbs Mouthwash for Overcoming Morning Sickness [Internet]. *Proceedings of the 5th International Conference on Health Sciences (ICHS 2018)*. 2019. Available from: <http://dx.doi.org/10.2991/ichs-18.2019.38>
18. Mukhtar K. The possible beneficial role of the regular use of potent mouthwash solutions in the treatment of COVID-19 [Internet]. <http://isrctn.com/>. Available from: <http://dx.doi.org/10.1186/isrctn10197987>
19. Ramesh A, Varghese S, Jayakumar ND, Malaiappan S. Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study. *J Periodontol* [Internet]. 2018 Oct;89(10):1241–8. Available from: <http://dx.doi.org/10.1002/JPER.17-0445>
20. Paramasivam A, Priyadharsini JV, Raghunandhakumar S, Elumalai P. A novel COVID-19 and its effects on cardiovascular disease. *Hypertens Res* [Internet]. 2020 Jul;43(7):729–30. Available from: <http://dx.doi.org/10.1038/s41440-020-0461-x>
21. S G, T G, K V, Faleh A A, Sukumaran A, P N S. Development of 3D scaffolds using nanochitosan/silk-fibroin/hyaluronic acid biomaterials for tissue engineering applications. *Int J Biol Macromol* [Internet]. 2018 Dec;120(Pt A):876–85. Available from: <http://dx.doi.org/10.1016/j.ijbiomac.2018.08.149>
22. Del Fabbro M, Karanxha L, Panda S, Bucchi C, Nadathur Doraiswamy J, Sankari M, et al. Autologous platelet concentrates for treating periodontal infrabony defects. *Cochrane Database Syst Rev* [Internet]. 2018 Nov 26;11:CD011423. Available from: <http://dx.doi.org/10.1002/14651858.CD011423.pub2>
23. Paramasivam A, Vijayashree Priyadharsini J. MitomiRs: new emerging microRNAs in mitochondrial dysfunction and cardiovascular disease. *Hypertens Res* [Internet]. 2020 Aug;43(8):851–3. Available from: <http://dx.doi.org/10.1038/s41440-020-0423-3>
24. Jayaseelan VP, Arumugam P. Dissecting the theranostic potential of exosomes in autoimmune disorders. *Cell Mol Immunol* [Internet]. 2019 Dec;16(12):935–6. Available from: <http://dx.doi.org/10.1038/s41423-019-0310-5>
25. Vellappally S, Al Kheraif AA, Divakar DD, Basavarajappa S, Anil S, Fouad H. Tooth implant prosthesis using ultra low power and low cost crystalline carbon bio-tooth sensor with hybridized data acquisition algorithm. *Comput Commun* [Internet]. 2019 Dec 15;148:176–84. Available from:

<https://www.sciencedirect.com/science/article/pii/S0140366419307017>

26. Vellappally S, Al Kheraif AA, Anil S, Assery MK, Kumar KA, Divakar DD. Analyzing Relationship between Patient and Doctor in Public Dental Health using Particle Memetic Multivariable Logistic Regression Analysis Approach (MLRA2). *J Med Syst* [Internet]. 2018 Aug 29;42(10):183. Available from: <http://dx.doi.org/10.1007/s10916-018-1037-z>
27. Varghese SS, Ramesh A, Veeraiyan DN. Blended Module-Based Teaching in Biostatistics and Research Methodology: A Retrospective Study with Postgraduate Dental Students. *J Dent Educ* [Internet]. 2019 Apr;83(4):445–50. Available from: <http://dx.doi.org/10.21815/JDE.019.054>
28. Venkatesan J, Singh SK, Anil S, Kim S-K, Shim MS. Preparation, Characterization and Biological Applications of Biosynthesized Silver Nanoparticles with Chitosan-Fucoidan Coating. *Molecules* [Internet]. 2018 Jun 12;23(6). Available from: <http://dx.doi.org/10.3390/molecules23061429>
29. Alsubait SA, Al Ajlan R, Mitwalli H, Aburaisi N, Mahmood A, Muthurangan M, et al. Cytotoxicity of Different Concentrations of Three Root Canal Sealers on Human Mesenchymal Stem Cells. *Biomolecules* [Internet]. 2018 Aug 1;8(3). Available from: <http://dx.doi.org/10.3390/biom8030068>
30. Venkatesan J, Rekha PD, Anil S, Bhatnagar I, Sudha PN, Dechsakulwatana C, et al. Hydroxyapatite from Cuttlefish Bone: Isolation, Characterizations, and Applications. *Biotechnol Bioprocess Eng* [Internet]. 2018 Aug 1;23(4):383–93. Available from: <https://doi.org/10.1007/s12257-018-0169-9>
31. Vellappally S, Al Kheraif AA, Anil S, Wahba AA. IoT medical tooth mounted sensor for monitoring teeth and food level using bacterial optimization along with adaptive deep learning neural network. *Measurement* [Internet]. 2019 Mar 1;135:672–7. Available from: <https://www.sciencedirect.com/science/article/pii/S0263224118311333>
32. PradeepKumar AR, Shemesh H, Nivedhitha MS, Hashir MMJ, Arockiam S, Uma Maheswari TN, et al. Diagnosis of Vertical Root Fractures by Cone-beam Computed Tomography in Root-filled Teeth with Confirmation by Direct Visualization: A Systematic Review and Meta-Analysis. *J Endod* [Internet]. 2021 Aug;47(8):1198–214. Available from: <http://dx.doi.org/10.1016/j.joen.2021.04.022>
33. R H, Ramani P, Tilakaratne WM, Sukumaran G, Ramasubramanian A, Krishnan RP. Critical appraisal of different triggering pathways for the pathobiology of pemphigus vulgaris-A review. *Oral Dis* [Internet]. 2021 Jun 21; Available from: <http://dx.doi.org/10.1111/odi.13937>
34. Ezhilarasan D, Lakshmi T, Subha M, Deepak Nallasamy V, Raghunandhakumar S. The ambiguous role of sirtuins in head and neck squamous cell carcinoma. *Oral Dis* [Internet]. 2021 Feb 11; Available from: <http://dx.doi.org/10.1111/odi.13798>
35. Sarode SC, Gondivkar S, Sarode GS, Gadbail A, Yuwanati M. Hybrid oral potentially

- malignant disorder: A neglected fact in oral submucous fibrosis. *Oral Oncol* [Internet]. 2021 Jun 16;105390. Available from: <http://dx.doi.org/10.1016/j.oraloncology.2021.105390>
36. Kavarthapu A, Gurumoorthy K. Linking chronic periodontitis and oral cancer: A review. *Oral Oncol* [Internet]. 2021 Jun 14;105375. Available from: <http://dx.doi.org/10.1016/j.oraloncology.2021.105375>
 37. Vellappally S, Abdullah Al-Kheraif A, Anil S, Basavarajappa S, Hassanein AS. Maintaining patient oral health by using a xeno-genetic spiking neural network. *J Ambient Intell Humaniz Comput* [Internet]. 2018 Dec 14; Available from: <https://doi.org/10.1007/s12652-018-1166-8>
 38. Aldhuwayhi S, Mallineni SK, Sakhamuri S, Thakare AA, Mallineni S, Sajja R, et al. Covid-19 Knowledge and Perceptions Among Dental Specialists: A Cross-Sectional Online Questionnaire Survey. *Risk Manag Healthc Policy* [Internet]. 2021 Jul 7;14:2851–61. Available from: <http://dx.doi.org/10.2147/RMHP.S306880>
 39. Sutakwa A, Nadia LS, Suharman S. Addition of blue pea flower (*Clitoria ternatea* L.) extract increase antioxidant activity in yogurt from various types of milk. *Jurnal Agercolere* [Internet]. 2021 May 21 [cited 2021 Aug 26];3(1):31–7. Available from: <https://faperta.unisan.ac.id/jurnal/index.php/jac/article/view/123>
 40. Adisakwattana S, Pasukamonset P, Chusak C. Chapter 18 - *Clitoria ternatea* beverages and antioxidant usage. In: Preedy VR, editor. *Pathology* [Internet]. Academic Press; 2020. p. 189–96. Available from: <https://www.sciencedirect.com/science/article/pii/B9780128159729000184>
 41. Oguis GK, Gilding EK, Jackson MA, Craik DJ. Butterfly Pea (*Clitoria ternatea*), a Cyclotide-Bearing Plant With Applications in Agriculture and Medicine. *Front Plant Sci* [Internet]. 2019 May 28;10:645. Available from: <http://dx.doi.org/10.3389/fpls.2019.00645>
 42. Mukherjee PK, Kumar V, Kumar NS, Heinrich M. The Ayurvedic medicine *Clitoria ternatea*—From traditional use to scientific assessment. *J Ethnopharmacol* [Internet]. 2008 Dec 8;120(3):291–301. Available from: <https://www.sciencedirect.com/science/article/pii/S0378874108004911>
 43. Kamkaen N, Wilkinson JM. The antioxidant activity of *Clitoria ternatea* flower petal extracts and eye gel [Internet]. Vol. 23, *Phytotherapy Research*. 2009. p. 1624–5. Available from: <http://dx.doi.org/10.1002/ptr.2832>
 44. Jadhav V, Deshmukh S, Mahadkar S. Evaluation of antioxidant potential of *Clitoria ternatea* L. *Int J Pharm Pharm Sci* [Internet]. 2013;5(2):595–9. Available from: https://www.researchgate.net/profile/Varsha-Jadhav-rathod/publication/257356228_Varsha_JadhavSwati_Deshmukh_and_Shivprasad_Mahadkar_2013Evaluation_of_antioxidant_potential_of_Clitoria_ternateaL_Int_J_Pharm_and_Pharmaceutical_sci52595-599/links/5b4097ce0f7e9bb59b10fbff/Varsha-Jadhav-Swati-Deshmukh-and-Shivprasad-Mahadkar-2013Evaluation-of-antioxidant-potential-of-Clitoria-ternateaL-Int-J-Phar-and-Pharmaceutical-sci52595-599.pdf

45. Danda AK. Comparison of a single noncompression miniplate versus 2 noncompression miniplates in the treatment of mandibular angle fractures: a prospective, randomized clinical trial. *J Oral Maxillofac Surg* [Internet]. 2010 Jul;68(7):1565–7. Available from: <http://dx.doi.org/10.1016/j.joms.2010.01.011>
46. Robert R, Justin Raj C, Krishnan S, Jerome Das S. Growth, theoretical and optical studies on potassium dihydrogen phosphate (KDP) single crystals by modified Sankaranarayanan–Ramasamy (mSR) method [Internet]. Vol. 405, *Physica B: Condensed Matter*. 2010. p. 20–4. Available from: <http://dx.doi.org/10.1016/j.physb.2009.08.015>
47. Krishnan V, Lakshmi T. Bioglass: A novel biocompatible innovation. *J Adv Pharm Technol Res* [Internet]. 2013 Apr;4(2):78–83. Available from: <http://dx.doi.org/10.4103/2231-4040.111523>
48. Soh CL, Narayanan V. Quality of life assessment in patients with dentofacial deformity undergoing orthognathic surgery—A systematic review [Internet]. Vol. 42, *International Journal of Oral and Maxillofacial Surgery*. 2013. p. 974–80. Available from: <http://dx.doi.org/10.1016/j.ijom.2013.03.023>
49. Lekha L, Kanmani Raja K, Rajagopal G, Easwaramoorthy D. Schiff base complexes of rare earth metal ions: Synthesis, characterization and catalytic activity for the oxidation of aniline and substituted anilines [Internet]. Vol. 753, *Journal of Organometallic Chemistry*. 2014. p. 72–80. Available from: <http://dx.doi.org/10.1016/j.jorganchem.2013.12.014>
50. Dhinesh B, Isaac JoshuaRamesh Lalvani J, Parthasarathy M, Annamalai K. An assessment on performance, emission and combustion characteristics of single cylinder diesel engine powered by *Cymbopogon flexuosus* biofuel [Internet]. Vol. 117, *Energy Conversion and Management*. 2016. p. 466–74. Available from: <http://dx.doi.org/10.1016/j.enconman.2016.03.049>
51. PradeepKumar AR, Shemesh H, Jothilatha S, Vijayabharathi R, Jayalakshmi S, Kishen A. Diagnosis of Vertical Root Fractures in Restored Endodontically Treated Teeth: A Time-dependent Retrospective Cohort Study. *J Endod* [Internet]. 2016 Aug;42(8):1175–80. Available from: <http://dx.doi.org/10.1016/j.joen.2016.04.012>
52. Vijayakumar GNS, Nixon Samuel Vijayakumar G, Devashankar S, Rathnakumari M, Sureshkumar P. Synthesis of electrospun ZnO/CuO nanocomposite fibers and their dielectric and non-linear optic studies [Internet]. Vol. 507, *Journal of Alloys and Compounds*. 2010. p. 225–9. Available from: <http://dx.doi.org/10.1016/j.jallcom.2010.07.161>
53. Kavitha M, Subramanian R, Narayanan R, Udhayabanu V. Solution combustion synthesis and characterization of strontium substituted hydroxyapatite nanocrystals [Internet]. Vol. 253, *Powder Technology*. 2014. p. 129–37. Available from: <http://dx.doi.org/10.1016/j.powtec.2013.10.045>
54. Sahu D, Kannan GM, Vijayaraghavan R. Size-Dependent Effect of Zinc Oxide on Toxicity and Inflammatory Potential of Human Monocytes [Internet]. Vol. 77, *Journal of Toxicology*

and Environmental Health, Part A. 2014. p. 177–91. Available from:
<http://dx.doi.org/10.1080/15287394.2013.853224>

55. Neelakantan P, Cheng CQ, Mohanraj R, Sriraman P, Subbarao C, Sharma S. Antibiofilm activity of three irrigation protocols activated by ultrasonic, diode laser or Er:YAG laser in vitro [Internet]. Vol. 48, International Endodontic Journal. 2015. p. 602–10. Available from: <http://dx.doi.org/10.1111/iej.12354>
56. Lekha L, Kanmani Raja K, Rajagopal G, Easwaramoorthy D. Synthesis, spectroscopic characterization and antibacterial studies of lanthanide(III) Schiff base complexes containing N, O donor atoms [Internet]. Vols. 1056-1057, Journal of Molecular Structure. 2014. p. 307–13. Available from: <http://dx.doi.org/10.1016/j.molstruc.2013.10.014>
57. Gopalakannan S, Senthilvelan T, Ranganathan S. Modeling and Optimization of EDM Process Parameters on Machining of Al 7075-B4C MMC Using RSM [Internet]. Vol. 38, Procedia Engineering. 2012. p. 685–90. Available from: <http://dx.doi.org/10.1016/j.proeng.2012.06.086>
58. Parthasarathy M, Isaac Joshua Ramesh Lalvani J, Dhinesh B, Annamalai K. Effect of hydrogen on ethanol-biodiesel blend on performance and emission characteristics of a direct injection diesel engine. Ecotoxicol Environ Saf [Internet]. 2016 Dec;134(Pt 2):433–9. Available from: <http://dx.doi.org/10.1016/j.ecoenv.2015.11.005>