

Original Research Article

CLINICAL PRESENTATION OF THE EPILEPTIC PATIENTS AT PEOPLE MEDICAL COLLEGE HOSPITAL, NAWABSHAH, PAKISTAN

ABSTRACT

Objective: study will determine clinical presentation, risk, and type of seizures in epileptic patients. **Methodology:** This observational study was conducted in Medicine department People Medical College Hospital (PMCH) Nawabshah from January 2020 to December 2020. 110 patients were included for study after informed permission of the patient or their relative. Male and female were selected. **Known epileptic patients were included in the study, patients with false seizures were excluded from the study. Statically analysis was done by software SSPS 22 version.** **Results:** Age ranged 18 to 60 years. All the patients included in the study epilepsy was noted. The mean age of patients was 37.40 SD 8.71 years. The minimum age was 28 years while maximum 60 years. Pearson Chi-Square Value 105.000^a Asymp. Sig. (2-sided) .000, Likelihood Ratio Value 135.012 Asymp. Sig. (2-sided) .000, Linear-by-Linear Association Value 16.297 Asymp. Sig. (2-sided) .000 which were statistically significant. **Conclusion: Epilepsy is treatable common neurological disease in Pakistan. quality of life can be improved by Education of the patients and their relatives, without socioeconomically burden.**

Key Words: Seizure, Epilepsy, Clinical, Medicine

INTRODUCTION

In developed countries and developing countries epilepsy is most common neurological disorder. Recurrence of seizure is termed as epilepsy, not all convulsions are due to epilepsy. [1] Excessive discharge of neurons in brain alter neurological function termed as a seizure. Seizure occur when there is imbalance between excitation and inhibition in brain.[2] **Numerous causes are included in Epilepsy and brain dysfunction.[3] various causes of epilepsy include genetic predisposition, head injury, brain tumors, stroke and drug or alcohol withdrawal. Seizure can be due to hypoglycemia, fever, meningitis and psychogenic. Seizures due to alcohol withdrawal is not epilepsy, epilepsy is generation of seizures by cognitive, neurobiological, social and psychological consequences of this condition.[4] Epilepsy most common CNS disorder 50 per 100000 new cases per year of the population. Incidence of epilepsy about 1%, refractory epilepsy in 1/3 and epilepsy from childhood 75%.[5] Worldwide 50 million people are affected by this non communicable disease epilepsy. Epilepsy is leading neurological disease in the world. Every year 61.4 per 100000 incidence of epilepsy is reported in population.[6] 5% prevalence rate of epilepsy reported in Iran.[7] All age groups are affected by epilepsy but common in young children and older age group.[8] Frontal lobe epilepsy is second following temporal lobe**

epilepsy, this type of epilepsy originate from frontal lobe, occur during wake or sleep.[9] 20-30% patients of focal epilepsy are associated with frontal lobe epilepsy.[10] Epilepsy can be misdiagnosed as sleep disorder, non epileptic seizures and psychiatric disorder.[11] Seizures classified as generalized, partial(now focal) and epileptic spasm. Limited part of cerebral hemisphere is involved in focal or partial epilepsy. Bilateral distributed neuronal network is involved in generalized seizure. Initially seizure can be focal later become generalized.[12] Seizure event is common in 10% population.[13] Seizure control is important for Doctors, nursing staff and patients or their relatives with learning disabilities.[14] Communication is important for patient with Doctors, regarding drugs side effects and seizures.[15] Patients with learning disabilities are at risk of uncontrolled seizures and increased mortality rate.[16] More than twenty drugs are in use for the successful treatment of epilepsy. These drugs act by preventing neuronal depolarization, blocking calcium or sodium channels, decrease electrical activity of the brain, enhance potassium channel function, neurotransmitter excitation of glutamate is inhibited.[17]

METHODOLOGY

This study was conducted in Medicine department PMCH Nawabshah from January 2020 to December 2020. 105 patients were included for study after informed permission of the patient or their relative. Male and female were selected. Detailed proforma was used for the study, detailed history, clinical examination of the patient, compulsory investigations of the patient, RBS, Urea, creatinine, LFT, Blood CP, Serum Electrolyte, serum Calcium Level, Urine DR, X-ray Chest, EEG, CT Scan Brain and MRI Brain. All patients with true seizures were included for this study, patients with pseudo seizures and seizures due to metabolic disorders were excluded from the study. Statically analysis was done by software SSPS 22 version.

RESULTS

The mean age of patients was 37.40 SD 8.71 years. The minimum age was 28 years while maximum 60 years. As shown in table 1.

Table 1. Descriptive Statistics

	N	Range	Minimum	Maximum	Mean		Std. Deviation
	Statistic	Statistic	Statistic	Statistic	Statistic	Std. Error	Statistic
age in years	105	32.00	28.00	60.00	37.4000	.85030	8.71294
Valid N	105						

The details of different demographic variables like age group, gender, marital status, address, occupation, education, Address, addiction SE Class and Family History are shown in table 2.

Table 2. The details of different demographical data

Demographic variables		Frequency	Percent (%)
Age Group	20-40 Years Young Age	78	74.3
	41-60 Years Middle Age	27	25.7
Gender	Male	69	65.7
	Female	36	34.3
Marital Status	Married	90	85.7
	Un-Married	15	14.3
Occupation	No Occupation	24	22.9
	House Wife	27	25.7
	Manual Worker	44	41.9
	Office Worker	10	9.5
Education	Educated	65	61.9
	Un-Educated	40	38.1
Address	Rural	73	69.5
	Urban	32	30.5
Addiction	No	80	76.2
	Yes	25	23.8
SE Class	Poor Class	82	78.1
	Middle Class	16	15.2
	Upper Class	7	6.7
Family History	No	86	81.9
	Yes	19	18.1
Total		105	100.0




Fig. 1. The comorbidities in epileptic patients like head injury, SOL, MR, depression, DM, HTN and HCV.




Fig. 2. The different types of seizures noted in epileptic patients.




Fig. 3. Different types of EEG findings noted in epileptic patients.

Table 3 shows the gender * seizure type Cross tabulation Pearson Chi-Square Value 105.000^a Asymp. Sig. (2-sided) .000, Likelihood Ratio Value 135.012 Asymp. Sig. (2-sided) .000, Linear-by-Linear Association Value 16.297 Asymp. Sig. (2-sided) .000 which were statistically significant.

Table 3. Statistical results.

Cross tabulation							
Variable			seizure type				Total
			focal seizures	partial seizures	generalized tonic clonic	absence seizure	
gender	male	Count	15	0	44	10	69
		% of Total	14.3%	0.0%	41.9%	9.5%	65.7%
	female	Count	0	36	0	0	36
		% of Total	0.0%	34.3%	0.0%	0.0%	34.3%
Total	Count	15	36	44	10	105	
	% of Total	14.3%	34.3%	41.9%	9.5%	100.0 %	
Chi-Square Tests							
	Value	df	Asymp. Sig. (2-sided)				
Pearson Chi-Square	105.000 ^a	3	.000				
Likelihood Ratio	135.012	3	.000				
Linear-by-Linear Association	16.297	1	.000				

N of Valid Cases	105		
a. 1 cells (12.5%) have expected count less than 5. The minimum expected count is 3.43.			

DISCUSSION

Following factors stigmatization, poverty, attitude, lack of adequate knowledge and coping skills are important in epilepsy care. Majority of the patients and relatives lack the knowledge about precipitating factors and the cause of epilepsy mentioned in earlier studies.[18] Knowledge about epilepsy treatment is important, role of supportive care, information about the disease, precipitants of seizures, role of treatment and adverse effects of drugs.[19] Neuroimaging CT Scan and MRI are helpful for detection of demyelinating disorders, atrophic lesion of brain and structural lesions of brain. Neuroimaging is helpful in patients with partial epilepsy abnormal findings were reported, structural lesion were localized by radiological support in symptomatic epilepsy.[20] Role of Doctors, nurses and health assistant is important in the treatment of epilepsy, communication between patient and health care professional is encouraged.[21] Epilepsy not considered as major public health issue because treatment is readily available and cheap.[22] Epilepsy associated with depression, due to biological and sociological factors. Abnormalities in neurotransmitter 5HT and Glutamate leads to depression in epilepsy.[23] Better outcome is associated with surgery in epilepsy, when single region of brain responsible for epilepsy.[24] Mesial temporal sclerosis a structural lesion with intractable seizure, surgery is option.[25] Seizures associated with behavior, memory change, altered responsiveness and posturing in the medial temporal region. Seizures are intractable with co morbidities. Surgical option is considered when two drugs fail. For the pathophysiology of temporal lobe epilepsy investigations are performed, genetic factors are important in temporal lobe epilepsy.[26] Stimulation of nerve where surgical resection is contraindicated, Vagus nerve is safe for stimulation with low complications, like vocal cord paralysis, hematoma and infection.[27] Epilepsy associated with other comorbid conditions, these are psychiatric disorders anxiety, depression, learning disabilities, autism, intellectual disability and attention deficit hyperactivity disorder. These comorbid conditions considered to be integral part of the disease, previously these comorbid conditions considered to be due to side effects of antiepileptic drugs or uncontrolled seizures. [28] Epileptic circuits limbic and hippocampal dysfunction associated with common psychiatric comorbidity depression. Depression is more common in patients with history of epilepsy and epileptic patients develop depression. About 30% patients of epilepsy have depression and 10% patients have bipolar disorder. [29]

CONCLUSION

Epilepsy common neurological disease in Pakistan, is treatable with cost effective drugs and minimal side effects. Epilepsy is major health problem in our country, long term treatment is needed in majority of the patients. Awareness about seizure, precautionary measures are compulsory. Precautions from fire, water and sudden fall during seizure. Improvement in dietary

habits, sleep and early treatment of any infection. Education of the patients and their relatives' quality of life can be improved without socioeconomically burden. **Early treatment and education about disease, stress can be reduced with improved quality of life.**

REFERENCES

1. Berg AT, Bercovic SF, Brodie MJ, et al. revised terminology and concepts for organization of seizures and epilepsies: report of ILAE Commission on classification and terminology. 2005-2009. *Epilepsia* 2010;51: 676-685.
2. Stafstorm CE. 2010. Pathophysiological mechanism of seizures and epilepsy: A Primer In *Epilepsy: Mechanism, models and translational perspectives*(ed Rho JM, Sankar R, Stasform CE). pp 3-19. CRC, Boca Raton, FL.
3. Shorvon SD, Andermann F, Gueirrine R (ed) 2011. *The causes of Epilepsy*. Cambridge University Press, Cambridge.
4. Sander JW, The natural history of epilepsy in the era of new antiepileptic drugs and surgical treatment. *Epilepsia* . 2003; 44(Suppl 1): 17-20.
5. Hauser W, Hersdorffer D. 1990. *Epilepsy: Frequency, causes and consequences*. Demos, New York.
6. Fiest KM, Sauro KM, Wiebe S, Patten SB, Kwon C-S, Dykeman J, et al. prevalence and incidence of epilepsy: a systemic review and meta-analysis of International studies. *Neurology*. 2017;88(3): 296-303.
7. Sayehmiri K, Tavan H, Sayhimiri F, Mohammadi I, Carson KV. Prevalence of epilepsy in Iran: a meta-analysis and systemic review. *Iran J Child Neurol*. 2014;8(4)9.
8. m/L/2 Kenneth W, Lindsay, ian Bone. *Neurology and Neurosurgery Illusterated*; 4th edition 2004, 90-103.
9. Haut S, *Frontal lobe epilepsy*. Medscope 2009.
10. Doelken MT ,Mennecke A, Huppertz HJ. Multimodality approach in cryptogenic epilepsy with focus on morphometric 3 TMRI. *J Neuroradiol*. 2012;39(2): 87-96.
11. MayoClinic staff. Frontal lobe seizure. 2008.<http://www.mayoclinic.org/diseases-conditions/frontal-lobe-seizures/symptoms-causes/dxc-20246880> [Accessed 20 Nov 2016}.
12. Berg AT, Millichap JJ. 2013. The 2010 revised classification of seizures and epilepsy. *Contnuum*. (Minneapp Minn) 19: 571-597.
13. Sander JW. The natural history of epilepsy in the era of new antiepileptic drugs and surgical treatment. *Epilepsia*. 2003;44(Suppl1) 17-20.
14. Haut S. *Frontal lobe epilepsy*. Medscope, 2009.
15. Doelken MT, Mennecke A, Huppertz HJ. Multimodality approach in cryptogenic epilepsy with focus on morphometric3TMRI. *J Neuroradiol*. 2012;39(2): 87-96.
16. Selemon L, Goldman-Rakic P. Longitudinal topography andinterdigitation of corticospinal projections in the rhesus monkey. *J Neurosci*. 1985;5:776-94.

17. Bui A, Kim H, Moroso M, Soltez I. 2015. Microcircuits in epilepsy: Heterogeneity and hub cells in network synchronization. *Cold Spring Harb Perspect Med* 10. 1101/cshperspect a022855.
18. Ogunrin OA, Adeyekun A, Adudu P. Etiologies of epilepsy and health-seeking itinerary of patients with epilepsy in a resource poor setting: analysis of 243 Nigerian Africans. *Seizure* 2013;22(7): 572-6.
19. Nwani PO, Arinzichi EO, Asomugha AL, Enwereji KO, Nwosu MC, Ogunniyi AO. Illness concept among people with epilepsy and their caregivers and preferred treatment methods in a suburban community in South East Nigeria. *West Afr J Med*. 3013;32(1): 26-30.
20. manz. Lakh.15 Obajmi MO, Fotunde OJ, Oganseyinde. Computed Tomography in childhood seizure disorder in lodbor west Africa ; *J Med* 2004;232: 16772.
21. Rodgers J, Namaganda S. Making information easier for people with learning disabilities. *Br J Learn Disabil*. 2005; 33 (2) 52-8.
22. Bertolote JM. Epilepsy as a public health problem. Role of the World Health Organization and the cooperation between WHO and non Governmental organizations. *Top Geogr Med*. 1994; 46 (3Suppl): S 28-30.
23. Lothe A, Didilot A, Hammers A, et al. Comorbidity between temporal lobe epilepsy and depression. a [18F] MPPF PET Study *Brain*. 2008 :131: 2765-2782.
24. de Tisi J, Bell GS, Peacock JL, et al. The long term outcome of adult epilepsy surgery, pattern of seizure remission, a relapse: a cohort study *Lancet*. 2011 :378: 1388-1395.
25. Thom A, Mathern JW, Cross JH, Bertram EH. 2010. Mesial temporal lobe epilepsy: How do we improve surgical outcome ? *Ann Neurol* 68: 424-434.
26. Liu Z, Mikati M, Holmes GL. 1995. Mesial temporal sclerosis: Pathogenesis and significance. *Pediatr Neurol* 12: 5-16.
27. Revesez D, Rydenhag B, Ben-Menacham E. Complications and safety of vagus nerve stimulation: 25 years experience at a single center. *J Neurosurg Pediatr*. 2016. 18: 97-104.
28. Brookes-Kayal AR, Bath KG, Berg AT, Galanopoulou AS, Holemes GL, Gensen FE, Kanner AM, O' Brien TJ, Whittemore VH, Vinawer MR, et al 2013. Issues related to symptomatic and disease-modifying treatments affecting cognitive and neuropsychiatric comorbidities of epilepsy. *Epilepsia* 54: 44-60.
29. Kanner AM. 2013. The treatment of depressive disorders in epilepsy: What all Neurologists should know. *Epilepsia* 54: 3-12.