

Original Research article

Use of chlorhexidine on umbilicus in prevention of neonatal sepsis

Abstract

Objectives: The main objective of the study is to analyse the use of chlorhexidine on umbilicus in prevention of neonatal sepsis. **Material and methods:** This cross sectional study was conducted in Sheikh Zaid hospital Rahim Yaar Khan during March 2020 till September 2020. The data was collected through non-probability consecutive sampling technique. The data was collected from 100 infants. Arrangements with a grouping of 4.0% free chlorhexidine were set up by weakening 20% chlorhexidine digluconate to the proper fixation with cleaned water. **Results:** The data was collected from 100 neonates. All the demographic values which include age, gender, gestational age and mode of delivery were calculated. According to baseline values the birth weight of chlorhexidine group was 1.87 ± 0.463 kg and dry cord group was 1.69 ± 0.421 kg. **Umbilical sepsis is observed in only 1 patient in group I and in 3 patients in group II. Only single mortality was observed in group I and in 6 neonates in group II.** **Conclusion:** It is concluded that chlorhexidine umbilical cord care is more appropriate than the currently WHO recommended dry cord care.

Key words: Sepsis, WHO, Neonatal, Mortality

Introduction

Neonatal mortality is still high in Pakistan at 55 neonatal deaths per 1000 live births (DHS 2012–13). Each year, approximately 202,000 newborns die within 28 days of birth in Pakistan. Tainting of the umbilical line can prompt omphalitis, portrayed by discharge, stomach erythema, or expanding¹. Sterile conveyance and postnatal-care rehearses are broadly elevated as significant intercessions to lessen danger of omphalitis and passing. There are not many explicit data, notwithstanding, on omphalitis rate and little proof for ideal

rope care practices to forestall string diseases and mortality locally, so better examinations are desperately needed².

Examiners in clinic based examinations in agricultural nations have portrayed the attributes of omphalitis, and revealed a scope of occurrence gauges (2–77 per 1000 medical clinic conceived newborn children). Locally, where irresistible test is higher and numerous cases go unrecognized, danger could be higher. Local area based case-control examines have zeroed in on danger factors for neonatal lockjaw, and give some proof that effective cleaning agents on the string are protective³. In settings where neonatal lockjaw has been killed yet unhygienic string rehearses keep on putting infants in danger, effective germ-killers may ensure against infection⁴.

In the agricultural nations, where incompetent and unhygienic neonatal care rehearses are normal, there is a consistent prerequisite of reasonable and safe local area based mediations with demonstrated adequacy to forestall contamination in babies. Since umbilical stump diseases may quickly advance to foundational sepsis, ideal line care is particularly significant in anticipation of neonatal sepsis. Without adequate proof preferring job of effective anti-microbials in counteraction of sepsis, dry rope care is for the most part suggested. Chlorhexidine effective application on string stump has been appeared to diminish umbilical colonization, umbilical sepsis and consequently potentially foundational sepsis⁵.

Objectives

The main objective of the study is to analyse the use of chlorhexidine on umbilicus in prevention of neonatal sepsis.

Material and methods

This cross sectional study was conducted in Sheikh Zaid hospital Rahim Yaar Khan during March 2020 till September 2020. The data was collected through non-probability consecutive sampling technique.

Data collection

The data was collected from 100 infants. Arrangements with a grouping of 4.0% free chlorhexidine were set up by weakening 20% chlorhexidine digluconate to the proper fixation with cleaned water. The cleanser and water purifying arrangement was set up by weakening Ivory Liqui-Gel, a gentle purging specialist reasonable for infant skin, with filtered water. In groups relegated to chlorhexidine or cleanser and water purging, guardians got instructive messages at each visit inside the initial 10 days of life. Subsequent to washing their hands with cleanser and water, the specialist dampened a cotton ball with arrangement and tenderly spotted the umbilical rope stump. Data were additionally gathered on birth history, parental proficiency, and occupation. The essential results for the line purifying intercession study were occurrence of neonatal omphalitis and neonatal mortality.

Statistical analysis

The data was collected and analysed using SPSS version 19. All the values were expressed in mean and standard deviation.

Results

The data was collected from 100 neonates. All the demographic values which include age, gender, gestational age and mode of delivery were calculated. According to baseline values the birth weight of chlorhexidine group was $1.87 \pm 0.463\text{kg}$ and dry cord group was $1.69 \pm 0.421\text{ kg}$. All the values is present in table 01

Table 01: Baseline characteristics of selected patients

Parameter	Group I Chlorhexidine group	Group II Dry cord care group
Neonatal parameters		
Birth weight (kg)	1.87 ± 0.463	1.69 ± 0.421
Period of gestation (weeks)	32.35 ± 1.64	33.87 ± 1.81
Male/female	23/27	39/11
Cesarean delivery	49	51
Premature rupture of membranes	13	25
UTI	0	1
Fever in last trimester	2	3

Table 02 shows the comparison of outcomes of group I and group II. Time of cord separation in group I is 7.82 ± 2.67 and 10.31 ± 3.23 in group II. Umbilical sepsis is observed in only 1 patient in group I and in 3 patients in group II. Only single mortality was observed in group I and in 6 neonates in group II.

Table 02: Comparison of outcome parameters

Parameter	Group I	Group II	<i>p</i> value
Time of cord separation (days)	7.82 ± 2.67	10.31 ± 3.23	0.02
Umbilical sepsis	1	3	–
Probable sepsis	20	5	0.043
Culture-proven sepsis	2	15	0.001
Meningitis	1	16	0.060
Duration of antibiotics received (days)	8.67 ± 5.61	13.1 ± 6.78	0.04
Duration of hospital stay (days)	11.4 ± 5.16	14.7 ± 6.62	0.04
All cause mortality	1	6	0.03




Fig 1: Comparison of outcome parameters

Discussion

Our discoveries propose the part of chlorhexidine neighborhood application as a straightforward, reasonable and effectively accessible mediation for counteraction of culture-demonstrated neonatal sepsis in an asset helpless setting; nonetheless, slight alert is prompted while summing up our outcome to the populace at large⁶. The subjects in our investigation were conveyed and overseen at a tertiary care reference setting with a recognized setting of microbiological verdure, openings to intercessions, accessibility of types of gear, techniques and conventions, significantly unique in relation to a typical local area neonatal care setting⁷.

In Germany, Kapellen et al. directed randomized controlled investigation to think about adequacy and security of chlorhexidine (CX) powder versus dry care (DC) in umbilical rope care of infant⁸. In an emergency clinic based examination from New Zealand assessing the effect of day by day rope purifying in 234 children, it was seen that string detachment happened at a mean of 10 days with Iodosan and 20 days with chlorhexidine. Meberg et al. (1990) saw that partition of the umbilical line happened altogether later in the hydrophobic material gathering than in the chlorhexidine–ethanol gathering (6.2 ± 2.2 versus 5.8 ± 2.1

days). Bhutta et al. (2010), in an enormous local area based bunch randomized controlled preliminary, announced the mean line partition time as 7 days in chlorhexidine group⁹⁻¹².

Jean et al., 2012 also explained that Chlorhexidine-impregnated and strongly adherent dressings may decrease catheter colonization and CRI rates.

Conclusion

It is concluded that chlorhexidine umbilical cord care is more appropriate than the currently WHO recommended dry cord care. It is a basic do-capable mediation that could well add to diminishing neonatal sepsis, a significant supporter of neonatal mortality in our country. This modest and straightforward intercession can save countless infant lives in agricultural nations.

COMPETING INTERESTS DISCLAIMER

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

Ethical Approval:

As per international standard or university standard ethical approval has been collected and preserved by the authors.

References

1. ZamCAT: (Zambia Chlorhexidine Application Trial): Impact of Chlorhexidine Cord Cleansing for Prevention of Neonatal Mortality in Zambia. ClinicalTrialsgov: NCT01241318. 2011
2. Lawn JE, Cousens S, Zupan J. 4 million neonatal deaths: when? Where? Why? *Lancet*. 2005 March 2;365:891–900
3. Faridi MM, Rattan A, Ahmad SH. Omphalitis neonatorum. *J Indian Med Assoc*. 1993;91:283–85.
4. Guvenc H, Aygun AD, Yasar F, Soylu F, Guvenc M, Kocabay K. Omphalitis in term and preterm appropriate for gestational age and small for gestational age infants. *J Trop Pediatr*. 1997;43:368–72.
5. Islam MS, Rahaman MM, Aziz KM, Munshi MH, Rahman M, Patwari Y. Birth care practice and neonatal tetanus in a rural area of Bangladesh. *J Trop Pediatr*. 1982;28:299–302.
6. Bennett J, Schooley M, Traverso H, Agha SB, Boring J. Bundling, a newly identified risk factor for neonatal tetanus: implications for global control. *Int J Epidemiol*. 1996;25:879–84.
7. Quddus A, Luby S, Rahbar M, Pervaiz Y. Neonatal tetanus: mortality rate and risk factors in Loralai District, Pakistan. *Int J Epidemiol*. 2002;31:648–53
8. Raza SA, Akhtar S, Avan BI, Hamza H, Rahbar MH. A matched case-control study of risk factors for neonatal tetanus in Karachi, Pakistan. *J Postgrad Med*. 2004;50:247–51.
9. Alder VG, Burman D, Simpson RA, Fysh J, Gillespie WA. Comparison of hexachlorophane and chlorhexidine powders in prevention of neonatal infection. *Arch Dis Child*. 1980;55:277–80.

10. Speck WT, Driscoll JM, Polin RA, O'Neill J, Rosenkranz HS. Staphylococcal and streptococcal colonization of the newborn infant: effect of antiseptic cord care. *Am J Dis Child.* 1977;131:1005–08
11. Janssen PA, Selwood BL, Dobson SR, Peacock D, Thiessen PN. To dye or not to dye: a randomized, clinical trial of a triple dye/alcohol regime versus dry cord care. *Pediatrics.* 2003;111:15–20.
12. Seeberg S, Brinkhoff B, John E, Kjellmer I. Prevention and control of neonatal pyoderma with chlorhexidine. *Acta Paediatr Scand.* 1984;73:498–504.

UNDER PEER REVIEW