

## **Original Research Article**

### **Prevalence of xerostomia and associated systemic risk factors in Riyadh, Saudi Arabia: A cross-sectional study**

#### **ABSTRACT**

##### **Purpose**

Xerostomia is a feeling of dry mouth and may result in poor oral hygiene, dental caries, mucosal lesions and burning mouth syndrome. Medication, systemic diseases and stress are common etiological factors of xerostomia. The present study investigated the prevalence of xerostomia and explored the possible risk factors associated with this condition among Saudi population.

##### **Methods**

The present cross-sectional study was conducted on 418 participants visiting at Maxillofacial clinic, at secondary care hospital, Riyadh. Demographic data and complete medical history were obtained from participants. All participants were requested to fill the Xerostomia-Inventory (XI) questionnaire. Comparisons between mean scores of XI-inventory and categorical variables like gender, comorbid conditions were made with students t-test or Wilcoxon test. Chi-square or Fischer's Exact test were used for comparison between categorical variables like gender and responses to XI-items (No/Yes). P value of <0.05 was considered significant. All analysis was done on JMP-version-12.

##### **Results**

Total 418-participants with mean age ( $41.1 \pm 12.8$  years) were included in the study in which 302-participants (72.2%) were female and 116-participants (27.8%) were male. Xerostomia's prevalence was 24.7% and it was higher in males (34%) as compared to females (21%). A statistically-significant association was found between xerostomia and participant's age ( $P < 0.0001$ ). The participants with systemic illness showed a significantly higher mean Xerostomia Inventory (XI) score as compared to the healthy patients. In our study population common comorbid conditions were hypertension and diabetes.

## **Conclusion**

The prevalence of xerostomia increases with increasing age and it has strong positive co-relation with chronic systemic diseases.

**Keywords:** Dry mouth, Xerostomia, Systemic diseases, Prevalence

UNDER PEER REVIEW

## INTRODUCTION

Xerostomia is the subjective feeling of dry mouth<sup>1</sup>. Individuals having xerostomia do not have sufficient amount of saliva to keep their mouth moist and wet. Saliva is a clear, watery fluid secreted by major (parotid, submandibular, and sublingual) and minor salivary glands<sup>2</sup>. Saliva has many functions like maintenance of oral homeostasis, lubrication of oral mucosa, swallowing, speaking and digestion along with antimicrobial and antifungal properties<sup>3</sup>. Diminished salivary flow causes increase rate of dental caries, gingivitis, dysgeusia, oral infections resulting in mucosal lesions, oral halitosis and burning mouth syndrome<sup>4, 5</sup>. Salivary flow rates are commonly affected by systemic diseases, medications, radiotherapy, chemotherapy, stress, anxiety and depression<sup>6-9</sup>. Autoimmune disorders such as Sjogren syndrome, sarcoidosis and local factors like sialolithiasis, sialadenitis and has been strongly associated with xerostomia. As per recent literature the two variables such as age and salivary flow rate are independent of each other in healthy individuals<sup>10</sup>. Epidemiological studies have estimated the prevalence of xerostomia between 1% and 62% depending on the population age, health characteristics, study designs and xerostomia measurement methods<sup>11</sup>. The prevalence varies according to gender lying between 13–26% in men and approximately 20–46% in women<sup>12</sup>.

Hyper or hyposalivation has a negative impact on individual's quality of life causing numerous clinical and social problems<sup>13</sup>. Diagnosis for dry mouth is based on detailed patient's history, clinical examination and required investigations. As xerostomia is the subjective feeling of dry mouth, so it can be assessed simply by asking few simple questions<sup>14</sup>.

The xerostomia inventory (XI) is a validated and frequently used questionnaire, that can help in the diagnosis of dry mouth (Xerostomia) in the clinical setting<sup>15</sup>. This questionnaire consists of 11 items, each on a 5-point Likert scale.

Many studies in the past has mentioned, xerostomia as frequently encountered problem and has measured its prevalence globally and shows variation among different areas and age groups <sup>16</sup>. Moreover, studies have a disparity in the inclusion criteria and methodologies for assessment. To our knowledge few studies have been conducted in Saudi Arabia exploring different aspects of xerostomia. There is still much to discover about dry mouth and its related problems. Therefore, the present study was conducted with an aim to investigate the prevalence of xerostomia among patients presented to our oral medicine and oral maxillofacial surgery clinics and analyze the possible risk factors associated with this condition.

#### **MATERIALS AND METHODS:**

This cross-sectional study was conducted after approval from hospital ethical committee at King Salman Hospital, department of dentistry, Ministry of Health, Riyadh, Saudi Arabia. The sample size was calculated by using the Research Advisors- 2006; with a confidence interval of 95%, a margin of error of 5% and an estimated proportion of the population suffering from xerostomia to be ( $p=0.5$ ). By using Cochran's sample size formula, the calculated minimal sample size was 385. To allow for attrition and missing data we inflated the sample size by 10%, thus a total of 418 participants were included in the study. Data was collected from outdoor maxillofacial clinic between December 2016 to December 2020.

**Inclusion criteria:** All patients older than 21 years of age were included in the study.

**Exclusion criteria:** Patients under radiotherapy and chemotherapy were not included in the study.

Demographic information and complete medical history of the participants was also recorded. Participants were divided into four age groups based on the quantile distribution and is depicted in Figure 1.

Data was collected on eight possible comorbid conditions based on previous literature search <sup>17</sup>. These included hypertension, diabetes, renal disease, respiratory symptoms, joint pains, liver disease, psychiatric symptoms or oral pain during eating. All patients were requested to fill the Xerostomia Inventory (XI) questionnaire. Xerostomia Inventory (XI) is an 11-item summated rating scale and Individuals were asked to choose a response to Xerostomia Inventory based on their last six weeks experience (Table-1). Five-point Likert scale with a score of 1= Never, 2= Hardly ever, 3= Occasionally, 4=Fairly often, and 5=Very often was utilized to evaluate participants perception regarding dry mouth (Table-1). The combined total score was calculated from all responses, a high total score represented the severity of the underlying xerostomia. For the purpose of further comparison “Fairly often” and “Very often” were combined as “Yes” and “Never”, “Occasionally” and “hardly ever” were combined as “No”. Comparisons between mean scores of XI inventory and categorical variables like gender and comorbid conditions was made with students t-test or Wilcoxon test. Chi-square or Fischer’s Exact test were used for comparison between categorical variables like gender and responses to XI items (No/Yes). P value of <0.05 was considered significant. All analysis was done on JMP version 12, SAS Institute INC.

### **Results:**

A total of 467 participants were examined and 442 participants agreed to contribute in the study. Out of 442 participants, 24 were excluded from the study due to incomplete questionnaire. Total 418 participants were included in the study in which 302 participants (72.2%) were female and 116 participants (27.8%) were male.

The age range was from 21-74 and mean age of the participants was  $41.1 \pm 12.8$  years. Patients were divided into four age groups based on the quantile distribution, 112 participants (26.8%) were in group 1 (20-31 years); 103

participants (24.6%) were in group 2 (32-39 years); 108 participants (25.8%) were in group 3 (40-51 years); and 95 participants (22.7%) group 4 (>51 years).

The data on the distribution of responses to Xerostomia Inventory (XI) is presented in Table 1. For purpose of analysis, “fairly often” and “often” responses to the standard question “Does your mouth feel dry?” were combined together as “Yes” shown in Table 2. The prevalence of xerostomia (response “Yes”) was 24.7% (103 participants; 64 females and 39 males) and presented in Table-2. Prevalence of xerostomia was higher in males (34%) as compared to females (21%). The prevalence of xerostomia increased with increasing age and statistically significant association was found between xerostomia and participant’s age ( $P < 0.0001$ ). The relationship between xerostomia and participant’s age is depicted in Table-3 and Figure 1.

In the present study population, the distribution of systemic disease among participants with Xerostomia is presented in Figure-2. In the present study population, 34% of patients had hypertension, 27% of patients were diabetic, and 6% of patients had depression/stress/anxiety. Approximately 5% of the study population reported Sarcoidosis; 4 % renal disease; 1% hepatitis C and 1% Sjogren Syndrome respectively. Only 2% patient had demonstrated local causes of xerostomia. In our study population most common reason for multifactorial etiology is hypertension and diabetes. The participants with systemic illness showed a significantly higher mean Xerostomia Inventory (XI) score as compared to the healthy patients (Table-4).

## **Discussion.**

The present cross-sectional study is a descriptive epidemiological study of xerostomia. According to Thomson et al. 2005<sup>18</sup>, a cross-sectional study design (questionnaire) should be used for this type of study which specify a description of the occurrence of a condition. Many studies have used a single item (absence or presence) to evaluate the xerostomia among their study population. Fox et al.

(1987)<sup>14</sup> had described the four-question model to classify individuals with both xerostomia and salivary gland hypofunction. Thomson et al. (1999b)<sup>15</sup> established the 'Xerostomia Inventory' (XI), an 11-item assessment which combines the responses to 11 individual items into a single summated score which represents the severity of xerostomia in contrast to the battery of items which are eventually analysed separately. The XI scale was also further endorsed in another a study in which two groups of participants were evaluated over a time span of 6 months. The first group consisted of cancer participants who were receiving radiotherapy for head/neck region and the second group was a control group of elderly volunteers ( $\geq 60$  years of age)<sup>19</sup>. The major drawback of this method is that no cut-off values for xerostomia inventory are reported in the literature. Therefore, it was suggested that, if the XI score is to be employed, it should be used along with at least one of the other single- item measures to further validate the xerostomia inventory' (XI) score<sup>18</sup>. Therefore, in the present study, single item (answer to question 1) as well as xerostomia inventory' (XI) score were utilized for analysis. The strong co-relation was found between the positive answer to question one (Q1) and XI score. The individuals with a positive response to the Q1 exhibited significantly higher XI score ( $36.2 \pm 9.0$ ) as compared to the individual who answered no to the Q1 ( $17.0 \pm 4.7$ ).

The xerostomia prevalence in the current study population was 24.7%. A recent meta-analysis of population-based studies has estimated the overall prevalence of dry-mouth to be 23.0% (95%CI; 18.0-28.0%) with high heterogeneity among different studies<sup>11</sup>. Another systematic review reported the prevalence of xerostomia from epidemiological studies of older populations range from 12 to 39%<sup>18</sup>. The high variability in xerostomia estimates has been credited to differences in study design, assessment methodology, study population, and the age of the participants<sup>18</sup>. Different studies have utilized different tools to assess the prevalence of dry mouth e.g., self-reported questionnaires, stimulated or un-stimulated salivary flow rate<sup>20, 21</sup>. Even in the studies who have done the xerostomia assessment through self-report, different methods of measurements

are employed such as some studies report only presence or absence of xerostomia<sup>22-24</sup>, some studies evaluate the frequency of a sensation of dry mouth<sup>25-27</sup> and others have used Xerostomia Inventory (XI).

Increasing age has been stated as a one of risk indicator for xerostomia<sup>11, 15, 23, 27</sup>. In the present study, the prevalence of xerostomia was 0% in group 1 (20-31 years); 5% in group 2 (32-39 years); 29% group 3 (40-51 years); and 71% in group 4 (>51 years). In our study population ageing seems to be determinant of occurrence of xerostomia. Many studies have shown similar results and have demonstrated an increase in the prevalence of Xerostomia with increasing age<sup>12, 28</sup>. Similar observation was reported by Benn et al.<sup>25</sup> in a nationally based study population, in which the prevalence of xerostomia was 5% in the younger age group (18-24) and 26% in the individuals aged 75 years or older, however, no consistent age gradient was reported<sup>25</sup>.

In the present study, 30% of the patients with diabetes had xerostomia. The positive co-relation was found between Xerostomia and diabetes. Narhi et al.<sup>29</sup> also reported a relationship between xerostomia and diabetes. Shirzaiy et al. 2016<sup>30</sup> has observed 66.7% of the diabetic patients had xerostomia. In the present study the highest percentage (100%) of xerostomia was observed in patients suffering from depression/anxiety and using medication for this purpose. Similar to our finding, several studies had reported the higher occurrence of xerostomia in patients with neuro-psychological disorders<sup>31</sup>.

In the present study no association was found between gender and xerostomia. Our results are were consistent with the results of Orellana et al.<sup>12</sup>, and Murray Thomson et al.<sup>31</sup> studies who could not find any relationship between these two variants i.e. gender and xerostomia. Conversely, Shirzaiy et al. 2016<sup>30</sup>; Fox et al. study<sup>14</sup> and Nederfors et al.<sup>28</sup> demonstrated the prevalence of Xerostomia higher in Women as compared to men. This might be explained by the fact that in current study population, 35.3% of males and only 17.8% females were in group 4 (age>51 years).

## **Conclusion**

The xerostomia prevalence in the current study population was 24.7% and xerostomia increases with increasing age and it has strong positive co-relation with chronic systemic diseases. In addition, dental and medical practitioners should carefully take the patients' history and should ask xerostomia-related questions especially in elderly population and in patients with history of polypharmacy. Health care providers have a responsibility of early detection of xerostomia and providing appropriate prevention and treatment advice to their patients.

### **COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## References

1. Joanna NDY, Thomson WM. Dry mouth—an overview. *Singapur dent J.* 2015;36:12-7.
2. Berk LB, Shivnani AT, Small JW. Pathophysiology and management of radiation-induced xerostomia. *J support Oncol.* 2005;3(3):191-200.
3. Dawes C, Pedersen AML, Villa A, Ekström J, Proctor G, Vissink A, et al. The functions of human saliva: A review sponsored by the World Workshop on Oral Medicine VI. *Arch Oral Biol.* 2015;60(6):663-74.
4. Ikebe K, Matsuda K-i, Morii K, Wada M, Hazeyama T, Nokubi T, et al. Impact of dry mouth and hyposalivation on oral health-related quality of life of elderly Japanese. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2007;103(2):216-22.
5. Atkinson JC, Grisius M, Massey WJ. Salivary hypofunction and xerostomia: diagnosis and treatment. *Dent Clin North Am.* 2005;49(2):309-26.
6. Chiappin S, Antonelli G, Gatti R, Elio FJCa. Saliva specimen: a new laboratory tool for diagnostic and basic investigation. *Clin Chim Acta* 2007;383(1-2):30-40.
7. Dawes C. Salivary flow patterns and the health of hard and soft oral tissues. *J Am Dent Assoc.* 2008;139:185-245.
8. Gardner MS, Rowland MD, Siu AY, Bundy JL, Wagener DK, Stephenson Jr. Comprehensive defensin assay for saliva. *Anal Chem* 2009;81(2):557-66.
9. Ship J. Diagnosing, managing, and preventing salivary gland disorders. *Oral Dis.* 2002;8(2):77-89.
10. Gupta A, Epstein JB, Sroussi H. Hyposalivation in elderly patients. *J Can Dent Assoc.* 2006;72(9).
11. Agostini BA, Cericato GO, Silveira ER, Nascimento GG, Costa FS, Thomson WM, et al. How common is dry mouth? Systematic review and meta-regression analysis of prevalence estimates. *Braz Dent J.* 2018;29(6):606-18.
12. Orellana M, Lagravère M, Boychuk DG, Major P, Flores-Mir C, Ortho CJJophd. Prevalence of xerostomia in population-based samples: a systematic review. *J Public Health Dent* 2006;66(2):152-8.
13. Scully C, Felix DJBdj. Oral medicine—update for the dental practitioner: dry mouth and disorders of salivation. *Br Dent J* 2005;199(7):423-7.
14. Fox PC, Busch KA, Baum BJ. Subjective reports of xerostomia and objective measures of salivary gland performance. *J Am Dent Assoc* 1987;115(4):581-4.
15. Thomson WM, Chalmers JM, Spencer AJ, Williams SM. The Xerostomia Inventory: a multi-item approach to measuring dry mouth. *Community Dent Health* 1999;16(1):12-7.
16. Abdullah MJ, dentistry e. Prevalence of xerostomia in patients attending Shorish dental speciality in Sulaimani city. *J Clin Exp Dent.* 2015;7(1):e45.
17. Mortazavi H, Baharvand M, Movahhedian A, Mohammadi M, Khodadoust A. Xerostomia due to systemic disease: a review of 20 conditions and mechanisms. *Ann Med health Sci Res.* 2014;4(4):503-10.
18. Thomson WM. Issues in the epidemiological investigation of dry mouth. *Gerodontology.* 2005;22(2):65-76.
19. Thomson W, Williams SJ. Further testing of the xerostomia inventory. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2000;89(1):46-50.

20. Navazesh M, Kumar SK. Measuring salivary flow: challenges and opportunities. *J Am Dent Assoc* . 2008;139:35S-40S.
21. Dawes C, Macpherson LJC. Effects of nine different chewing-gums and lozenges on salivary flow rate and pH. *Caries Res* 1992;26(3):176-82.
22. Iwasaki M, Yoshihara A, Ito K, Sato M, Minagawa K, Muramatsu K, et al. Hyposalivation and dietary nutrient intake among community-based older Japanese. *Geriatr Gerontol Int*. 2016;16(4):500-7.
23. Hochberg MC, Tielsch J, Munoz B, Bandeen-Roche K, West SK, Schein OD. Prevalence of symptoms of dry mouth and their relationship to saliva production in community dwelling elderly: the SEE project. *Salisbury Eye Evaluation. J Rheumatol*. 1998;25(3):486-91.
24. Jacobsson LT, Axell TE, Hansen BU, Henricsson VJ, Larsson Å, Lieberkind K, et al. Dry eyes or mouth—An epidemiological study in Swedish adults, with special reference to primary Sjögren's syndrome. *J Autoimmun* . 1989;2(4):521-7.
25. Benn A, Broadbent J, Thomson WJ. Occurrence and impact of xerostomia among dentate adult New Zealanders: findings from a national survey. *Aust Dent J* .2015;60(3):362-7.
26. Anttila SS, Knuutila ML, Sakki TK. Depressive symptoms as an underlying factor of the sensation of dry mouth. *Psychosom Med* .1998;60(2):215-8.
27. Ekbäck G, Åström AN, Klock K, Ordell S, Unell LJA. Variation in subjective oral health indicators of 65-year-olds in Norway and Sweden. *Acta Odontol Scand*. 2009;67(4):222-32.
28. Nederfors T, Isaksson R, Mörnstad H, Dahlöf CJ. Prevalence of perceived symptoms of dry mouth in an adult Swedish population—relation to age, sex and pharmacotherapy. *Community Dent Oral Epidemiol*. 1997;25(3):211-6.
29. Närhi TO, Meurman JH, Ainamo AJ. Xerostomia and hyposalivation. *Drugs Aging*. 1999;15(2):103-16.
30. Shirzaiy M, Bagheri FJ. Prevalence of xerostomia and its related factors in patients referred to zahedan dental school in Iran. *J. Clin. Exp. Dent* 2016;2(1).
31. Murray Thomson W, Poulton R, Mark Broadbent J, Al-Kubaisy SJA. Xerostomia and medications among 32-year-olds. *Acta Odontol Scand*. 2006;64(4):249-54.

**Table 1: Distribution of responses to Xerostomia Inventory (XI). Score of Response to each Question; Never =1; Hardly ever = 2; Occasionally= 3; Fairly often= 4 and Very often=5**

Questions	Never		Hardly ever		Occasionally		Fairly often		Very often		Mean $\pm$ SD
	N	%	N	%	N	%	N	%	N	%	
<i>My mouth feels dry</i>	93	22.2	107	25.6	115	27.5	66	15.8	37	8.8	2.63 $\pm$ 1.24
<i>I have difficulty in eating dry foods</i>	133	31.8	108	25.8	106	25.4	51	12.2	20	4.7	2.32 $\pm$ 1.18
<i>I get up at night to drink</i>	226	54.1	82	19.6	75	17.9	23	5.5	12	2.8	1.83 $\pm$ 1.08
<i>My mouth feels dry when eating a meal</i>	166	39.7	95	22.7	82	19.6	60	14.3	15	3.5	2.19 $\pm$ 1.21
<i>I sip liquids to aid in swallowing food</i>	163	39.0	118	28.2	77	18.4	38	9.1	22	5.2	2.13 $\pm$ 1.18
<i>I suck sweets or cough lozies to relieve dry mouth</i>	237	56.7	83	19.9	61	14.6	30	7.2	7	1.6	1.77 $\pm$ 1.05
<i>I have difficulties swallowing certain foods</i>	166	39.7	96	23	92	22.0	47	11.2	17	4.0	2.17 $\pm$ 1.19
<i>The skin of my face feels dry</i>	247	59.1	102	24.4	38	9.1	23	5.5	8	1.9	1.67 $\pm$ 0.98
<i>My eyes feel dry</i>	253	60.5	108	25.8	44	10.5	6	1.4	7	1.6	1.58 $\pm$ 0.86
<i>My lips feel dry</i>	147	35.2	179	42.8	49	11.7	29	6.9	14	3.3	2.00 $\pm$ 1.02
<i>The inside of my nose feels dry</i>	319	76.3	50	12	34	8.1	10	2.4	5	1.2	1.40 $\pm$ 0.83

\*Wilcoxon test

#Yes= (Fairly often + very often)

**Table 2: Modified Responses to Standard Question for diagnosis of Xerostomia**

<b><i>My mouth feels dry</i></b>	<b>N</b>	<b>%</b>	<b>Mean XI score <math>\pm</math> SD*</b>
Yes <sup>#</sup>	103	24.7	36.2 $\pm$ 9.0
No	315	75.3	17.0 $\pm$ 4.7

\*Wilcoxon test

<sup>#</sup>Yes= (Fairly often + very often)

UNDER PEER REVIEW

**Table 3. Association of age with Xerostomia score**

<b>Age category Years</b>	<b>Number</b>	<b>Mean XI score</b>	<b>Std Dev</b>	<b>p-value*</b>
<32	112	13.2679	1.6107	<0.0001
32-39	103	16.9126	4.2127	
40-51	108	23.5648	7.2192	
>51	95	34.7789	10.0076	

\*Anova was done


UNDER PEER REVIEW

**Table 4: Relationship of Xerostomia with Systemic illness**


	<i>My mouth feels dry</i>		<i>p-value</i> <sup>®</sup>	<i>XI score</i>		<i>p-value</i> <sup>β</sup>
	No N (%)	Yes N (%)		Mean	Std Dev	
<b>Hypertension</b>						
No	238 (56.9)	39 (9.3)	0.03	19.45	9.7	<.0001
Yes	109 (26.1)	32 (7.7)		26.1	9.8	
<b>Hepatitis C</b>						
No	346 (82.8)	67 (16)	0.003*	21.51	10.1	0.003
Yes	1 (0.24)	4 (0.96)		38.4	9.2	
<b>Diabetes</b>						
No	268 (64.11)	37 (8.8)	<0.0001	19.3	9.05	<0.0001
Yes	79 (18.90)	34 (8.1)		28.23	10.5	
<b>Sjogren Syndrome</b>						
No	346 (82.8)	67 (16)	0.003*	21.44	10.0	0.0006
Yes	1 (0.24)	4 (0.96)		44	7.2	
<b>Sarcoidosis</b>						
No	332 (79.4)	66 (15.8)	0.03	21.4	10.2	0.002
Yes	15 (3.6)	5 (1.2)		27.4	10.3	
<b>Renal diseases</b>						
No	336 (80.4)	64 (15.3)	0.02*	21.3	10.1	0.001
Yes	11 (2.6)	7 (1.7)		28.77	11.6	
<b>Depression/ Anxiety/Stress</b>						
No	334 (79.9)	58 (13.9)	<0.0001	21.04	9.8	<0.0001
Yes	13 (3.1)	13 (3.1)		31.8	10.9	
<b>Local Causes</b>						
No	342 (81.8)	67 (16)	0.05	21.51	10.2	0.001
Yes	5 (1.2)	4 (0.96)		30.6	7.3	

®chi-square test    \*Fischer's exact test    β Wilcoxon test

**Figure-1:** Prevalence of Xerostomia in different age groups



**Figure 2:** Xerostomia and Systemic diseases



UNDER