

Saudi Population awareness regrading methods of colorectal cancer screening

ABSTRACT

Aims: To assess the level of knowledge about colorectal cancer screening methods among the population in Saudi Arabia.

Study design: A cross-sectional study

Place and Duration of Study: General Surgery Department, between January 2021 and November 2022.

Methodology: This was a cross-sectional, survey-based study conducted in Saudi Arabia. Data was collected using a predesigned self-administered online questionnaire. Inclusion criteria include Saudi male and female between 18-65 years in Saudi Arabia. Then data were analyzed by the SPSS program, version 25 at a level of significance of 0.05.

Results: A total of randomly taken 794 participants who filled our questionnaire according to the inclusion criteria. Most of our participants (67.1%) were females and 32.9% of them were males. Only 17.5% of the participants were less than 20 years old, 62.5% of them were between 20 and 30 years old and 20% were more than 30 years old. Most of our participants (58.3%) were university educated,

Conclusion: CRC screening were not well known in Saudi Arabia. Educational programs are compulsory and should target people at risk.

Keywords: colorectal Cancer, Screening, Saudi Population

1. INTRODUCTION

The third malignancy commonly diagnosed among all gender which also consider the fourth leading cause of cancer-associated mortality, despite its low incidence in Arab people is Colorectal cancer (CRC) [1]. in 2012 about 700,000 deaths were announced along with 1.4 million newly diagnosed cases internationally it should be noted that in 29% of the cases the rectum will be affected and about 71% of cases it affects the colon. Certainly, Countries in the west have high incidence rates, unlike the developing nations where rates are particularly low [2]. The application of colorectal screening programs lately decreases the incidence rate of colorectal cancer through the early discovery and treatment indeed, if the tumor discovered and surgically removed early the risk of CRC can be reduced by 95% [3]. The suggested age of screening for CRC by The American Cancer Society (ACS) is 45, they recommend two types of tests for the screening: visual/structural tests and stool tests, every 10 years the colonoscopy should be done, 5 years for the colon scan and the flexible sigmoidoscopy as they are the visual/structural tests, the fecal immunochemical test(FIT) and a guaiac-based fecal occult blood test (gFOBT) should be done yearly and every 3 years for multitarget DNA (MT-DNA or FIT DNA) test and they are the Fecal-based tests [2].

(Almadi et al., 2015) a population-based study found that people in Saudi Arabia have fewer backgrounds regarding the screening program, red flag symptoms, and the risk factors of colorectal cancer, however, the females and well-educated people had more knowledge [4].

The participants in large numbers of the previous researches that were done in KSA were from shopping centers that are not illustrative the general public, furthermore, the questionnaire was used to assess the knowledge about CRC is unsubstantiated [3]. Factors including social-class, individuals' beliefs, application of preventive care, healthcare access and background of cancer and screening tests and guidelines, considers as Barriers to Colorectal cancer screening [5].

In current work aimed to assess the knowledge about screening methods of colorectal cancer in Saudi Arabia population, due to the insignificant number of researches that done in Saudi Arabia regard this topic despite its importance. To Identify the level of knowledge among Saudi population concerning colorectal cancer screening methods and most common thoughts and believes that prevent them from agree with screening.

2. MATERIAL AND METHODS

This is an observational cross-sectional study in Saudi Arabia was done from January 2020 to November 2021. To assess the level of knowledge about colorectal cancer screening methods among the population in Saudi Arabia. The population of the study ranged in age from 18-65. The sample size was estimated using the Qualtrics calculator with a confidence level of 95%; a sample size of 384. A- Inclusion criteria: Inclusion criteria include Saudi male and female between 18-65 years in Saudi Arabia. B- Exclusion criteria Exclusion criteria include non-Saudi, younger than 20 and more than 65 years. Data collection was done in the form of the participants' responses to the questions. The survey tool was an anonymous self-administered questionnaire in Arabic, containing questions regarding colorectal cancer screening related knowledge

Data was entered on the computer using the "Microsoft Office Excel Software" program (2016) for windows. Data was then transferred to the Statistical Package of Social Science Software (SPSS) program, version 20 (IBM SPSS Statistics for Windows, Version 20.0. Armonk, NY: IBM Corp.) to be statistically analyzed.

3. RESULTS AND DISCUSSION

Table (1) showed the socio-demographic data of the participants. We included a total of randomly taken 794 participants who filled our questionnaire according to the inclusion criteria. Most of our participants (67.1%) were females and 32.9% of them were males. Only 17.5% of the participants were less than 20 years old, 62.5% of them were between 20 and 30 years old and 20% were more than 30 years old. Most of our participants (58.3%) were university educated, 6.8% were more than Bachelor's degree, 34.5% of them were secondary educated and only 0.4% were non educated.

Table (1): Sociodemographic characteristics of participants (n=794)

Parameter	No.	Percent	
Gender			
	Male	261	32.9
	Female	533	67.1

Age	Less than 20	139	17.5
	20 - 30years old	496	62.5
	31- 40years old	69	8.7
	41 – 50 years old	51	6.4
	51 - 60 years old	28	3.5
	More than 60	11	1.4
Residence area	Southern area	108	13.6
	Eastern Region	150	18.9
	The northern area	71	8.9
	Western Region	225	28.3
	Central Region	240	30.2
Education level	uneducated	3	4.
	secondary	274	34.5
	Bachelor's degree	463	58.3
	Master's	31	3.9
	PhD	23	2.9

Table (2) showed the Knowledge of participants of colorectal cancer. 74.1% of our participants had heard about colorectal tumors before, 55.9% of them think that colorectal tumors are present in abundance in Saudi Arabia, only 17.9% Has a member in their family had colorectal tumors, while 28.1% know someone with colorectal tumors. 84.1% of our participants think colorectal tumors are a preventable condition and 62.3% of them agreed that colon and rectal tumors are fatal.

Table (2): Knowledge of participants of colorectal cancer(n=794).

Questions	Yes	No
Have you ever heard of colorectal tumors?	588 74.1%	206 25.9%
Do you think that colorectal tumors are present in abundance in Saudi Arabia?	444 55.9%	350 44.1%
Has anyone in your family ever had colorectal tumors?	142 17.9%	652 82.1%
Do you know someone with colorectal tumors?	222 28.0%	572 72.0%

Do you think colorectal tumors are a preventable condition?	668 84.1%	126 15.9%
I think colon and rectal tumors are fatal	495 62.3%	299 37.7%

Table (3) showed the Knowledge of screening methods and associated variables of colorectal cancer. When we asked about what they think is the recommended age for screening for colorectal tumors, 28.8% of our participants thought that its 18-30 years old, 27.7 of them recommended 31-40 years old to be the beat age of CRC screening, 23% recommended 41-50 years old and only 20.4% recommended the age above 51 to be the best age of CRC screening.

Regarding the factors that they think may increase the risk of developing colorectal tumors, 44% thought that aging is the factor affecting CRC, 73% reported family history, 36% reported the obesity, 45% choosed the genetic factors, 37% choosed smoking, 35% thought it's the age above 50, 47% reported presence of polyps in the lining of the colon and rectum, 63% said its Inflammatory bowel diseases and only 5% thought that diet rich in fat and low in fiber may be a factor in CRC.

Anal bleeding, diarrhea or constipation, Presence of blood in stool, Abdominal pain, Sudden rapid weight loss, Increase in mucous secretions in stool and appearance without any symptoms were all reported to be symptoms or auxiliary factors for the emergence of colon and rectal tumors with percent of (61%, 59%, 62%, 58%, 49%, 30% and 42%) respectively. We reported that 50.6% of our participants previously heard of early detection of colorectal tumors. The early detection methods for large intestine tumors that the participants know were Endoscopy of the rectum and anal canal in addition to testing for blood in the stool sample (82%), Detection of blood in a stool sample (56%), Barium staining x-ray of the large intestine (42%), Abdominal CT scan (47%), Barium tincture for the large intestine (32%), Colon and rectal biopsy (66%), Clinical examination of the rectum (47%) and Blood tests (44%). 82.1% of our participants thought that early detection of colorectal tumors is very useful, 8.4% of them thought its useful, 9% of them thought its not useful and 8.6% didn't know if it's useful or not. Almost one quarter of our participants considered doing an early detection for themselves and they choosed different methods

Table (3): Knowledge of screening methods and associated variables of colorectal cancer

Parameter	No.	Percent
What do you think is the recommended age for screening for colorectal tumors?	• 18-30 years old	229 28.8
	• 31-40 years old	220 27.7
	• 41 - 50 years old	183 23.0
	• 51 - 60 years old	97 12.2
	• 61 - 65 years old	65 8.2
Which of the following is one of the factors that you think may increase the risk of developing colorectal tumors?	•aging	351 0.44
	Having a family history of colorectal tumors	585 0.73

	•obesity	291	0.36
	•Having some genes	365	0.45
	•to smoke	301	0.37
	•Age over 50 years old	278	0.35
	The presence of polyps in the lining of the colon and rectum	376	0.47
	Inflammatory bowel diseases (Crohn's and ulcerative colitis)	503	0.63
	•A diet rich in fat and low in fiber	397	0.5
	•exposure to radiation	177	0.22
Symptoms or auxiliary factors for the emergence of colon and rectal tumors	• anal bleeding	486	0.61
	• diarrhea or constipation	472	0.59
	• Presence of blood in stool	500	0.62
	• Abdominal pain	464	0.58
	• Sudden rapid weight loss	396	0.49
	• Increase in mucous secretions in stool	242	0.30
	• It may appear without any symptoms	338	0.42
Have you ever heard of early detection of colorectal tumors?	• Yes	402	50.6
	• No	392	49.4
If your answer is yes to the previous question above, please complete the research. The early detection methods for large intestine tumors that you know are:	• colonoscopy	548	1.36
	Endoscopy of the rectum and anal canal in addition to testing for blood in the stool sample	332	0.82
	• Detection of blood in a stool sample	229	0.56
	Barium staining x-ray of the large intestine	170	0.42
	• Abdominal CT scan	191	0.47
	• Barium tincture for the large intestine	130	0.32
	• Colon and rectal biopsy	268	0.66
	• Clinical examination of the rectum	190	0.47
	• Blood tests	180	0.44
How useful is it to detect colorectal tumors early in your opinion?	• very useful	652	82.1
	• useful	67	8.4
	• not very useful	7	9.
	• I don't know how useful it is	68	8.6

Have you ever considered doing an early detection for yourself?	• Yes	206	25.9
	• No	588	74.1
Would you prefer to make an early detection for you personally using any of the?	• colonoscopy	302	0.38
	• Blood tests	367	0.46
	• Detection of blood in a stool sample	195	0.24
	• Abdominal CT scan	236	0.29
	Proctoscopy and anal canal in addition to blood test in stool sample	139	0.17
	Barium staining x-ray of the large intestine	132	0.16
	• Barium tincture for the large intestine	72	0.09
	• Clinical examination of the rectum	105	0.13
	• Colon and rectal biopsy	82	0.10
	• Colon and rectal biopsy	7	0.01
Was your choice to answer the above question based on:	• Fear of working with binoculars	390	49.1
	• Not convinced of the issue	112	14.1
	•History of examination	292	36.8
If you have colorectal adenomas, do you want to know about it?	•Yes	562	70.8
	•No	73	9.2
	•I do not know	159	20.0
How worried are you about discovering that you have colorectal tumors?	•not worried	127	16.0
	•somewhat worried	227	28.6
	•Very worried	190	23.9
	•I do not know	158	19.9
	•I'm not worried at all	92	11.6
What are your sources of information about early screening for colorectal tumors?	•Internet and social media	587	0.73
	•Awareness Videos	193	0.24
	/olunteer campaigns in public places of the community	148	0.18
	•Consult the doctor and ask him directly	165	0.20
	•field of study	20	0.02
	•I do not know	12	0.01

Discussion

Personal attitude towards screening and availability of the appropriate screening methods affects highly screening strategies for colorectal cancer prevention. Limited knowledge of the population regarding CRC and its screening methods can strongly influence their decision, whether it is based on the experiences of family and friends, or as learned knowledge. Although Saudi Arabia has the available resources for diagnostic and therapeutic procedures, there is lack of a consistent organized screening program for CRC screening. Overall, the knowledge of the population is less than expected to encourage the spread of screening behavior, that's why we focused in assessing the level of awareness of the general population regarding such a critical subject.

We present a study that assesses the awareness of and attitudes towards undergoing CRC screening in KSA. In this study we included a sample of 794 participants who responded to the questionnaire according to our inclusion criteria. Most of our participants (67.1%) were females, 32.9% were males and more than half of them (62.5%) were between 20 and 30 years old, 17.5% were less than 20 years old and only 10.3% were more than 30 years old. The female, young age dominance in the participants may be because females and young aged individuals are more liable to use social media and respond to the questionnaire. The knowledge level of the population regarding CRC was 74.1% with a good level of awareness about its causes and risk factors. This result is less than the results found in Makkah by Ammar A Al-Maghrabi [6], who found that the majority of the participants (85.7%) heard of colorectal cancer. In another study, 37.4% of the total participants have heard about CRC, which is much lower results than ours [7].

In our study 50.6% of our participants were aware of the early detection of CRC using the CRC screening methods. This is higher than other studies done in Saudi Arabia. In Osamah Abdullah A. et al. [8], from all participants, only 136 (16.3%) were aware about screening test of CRC and 87 (10.5%) were not sure. In another study 32.7% of the participants knew the recommended beginning of colorectal cancer screening [9]. In similar cross-sectional studies, this rate was more favourable (47.9%; 83.0%) [9,10]. These respondents were likely to see their physician more frequently. Another Saudi study conducted on men in Alahsa showed that only 28.8% participants in the study knew about screening methods of CRC [11]. Another study from Lebanon reported higher percentage of knowledge about CRC screening as more than half of the participants knew the importance for screening [12], however a much lower results were reported in another study from East Iran, which showed that more than 90% did not have knowledge about CRC screening [13].

Our participant had average information regarding the risk factors and warning signs of CRC. The CRC risk factors that were most commonly recognized were Having a family history of colorectal tumors (73%), bowel disease (63%), A diet rich in fat and low in fiber (50%), family history (45%) and aging (44%). Participants were least aware of exposure to radiation (22%), obesity (36%) and smoking (37%) as risk factors. This results is a little higher than Mohamad Ali Tfaily's results which indicated that only 31.5% and 17.2% of the participants were aware of the warning signs and risk factors of CRC, respectively [14], and the reported percentages of CRC risk factors that were most commonly recognized were red meat (64%), bowel disease (62%), and smoking (60%). Participants were least aware of diabetes (26%) and no physical exercise (37%) as risk factors, and only 17.2% of the participants were aware of nine or more of the listed risk factors. Another previous study conducted in the MENA region reported similar studies. In these studies, it was and about 82%, 85% and 94% of participants from the United Arab Emirates had poor to no knowledge

on colorectal cancer risk factors, warning signs and screening methods respectively [13]. Participants were most commonly aware of presence of fecal blood anal bleeding and diarrhea or constipation (62%, 61% and 59%, respectively) as warning signs for CRC, and were least aware of Increase in mucous secretions in stool (30%).

Individuals were more likely to be aware of CRC risk factors and warning signs in relation with age and educational level, and our study found a significant association with these factors. This association was also reported in a number of other studies [15,16,17]. While different studies have demonstrated an increase in awareness of CRC screening with education and employment, our results showed no significant association with these factors [7,18,19].

In this study, when we asked the participants about the ability of applying screening on themselves, the majority 48% preferred blood test as a screening method, 38% choosed colonoscopy while Barium tincture for the large intestine, Colon and rectal biopsy, Clinical examination of the rectum were the least methods to be choosed, the decisions based on previous questions were further analyzed and showed that prior knowledge of the screening types of endoscopic and radiological methods as well as personal fear of doing the test in general, contributed to the decisions made.

In another study, after posing a direct question to the study respondents about their personal decision to undergo screening versus not undergoing screening, their responses were examined in relation to their choice of whether it was an invasive method or a less invasive method. It was shown significantly that among the people who agreed to be screened, colonoscopy or sigmoidoscopy with fecal occult blood testing was chosen as a screening method. Those who did not agree to be screened chose a less invasive method, which was barium enema or CT scan of the abdomen [20]. In Qumseya et al., they reported education below secondary school level was associated with a lower probability of accepting colonoscopy as a screening method and increased age was associated with decreased probability of accepting FOBT as a screening method [20], which is almost like our finding, as we reported a clear effect of educational level of the participants in knowledge of colonoscopy as a screening method in CRC and as well as the probability of the participant to accept it to be applied in himself.

4. CONCLUSION

There was a lack of awareness of CRC checks in Saudi Arabia. Educational programs are compulsory and should target people at risk because they are considered the most common sources of information in schools, the Internet and social media.

ACKNOWLEDGMENT

We would like to thank all the participants in this study for cooperating in the research by taking part actively and investing their time

FUNDING

This study has not received any external funding.

CONFLICT OF INTEREST

The authors declare that there are no conflicts of interests.

ETHICAL CONSIDERATION

Ethical approval was obtained from Research Ethics Committee at Prince Sattam Bin Abdulaziz University, Al-Kharj, Saudi Arabia. with the IRB approval number (REC-HSD-54-2021)

REFERENCES

- 1- Al-masoudi, T. H., Alghamdi, K. M., & Bokhari, G. A. (2021). Assessment of Knowledge about Colorectal Cancer Screening among Attendants of Primary Healthcare Centers. 9(4), 270–279.
- 2- Althobaiti, A., & Jradi, H. (2019). Knowledge, attitude, and perceived barriers regarding colorectal cancer screening practices and risk factors among medical students in Saudi Arabia. BMC Medical Education, 19(1), 1–8. <https://doi.org/10.1186/s12909-019-1857-7>
- 3- Arafa, M. A., & Farhat, K. (2015). Colorectal Cancer in the Arab World--Screening Practices and Future Prospects. Asian Pacific journal of cancer prevention : APJCP, 16(17), 7425–7430. <https://doi.org/10.7314/apjcp.2015.16.17.7425>
- 4- Carnahan, L. R., Jones, L., Brewer, K. C., Elizabeth, A., Peterson, C. E., Ferrans, C. E., Cipriano-steffens, T., Polite, B., Maker, A. V, Rauscher, G. H., Illinois, A., & Medical, M. (2021). HHS Public Access. 36(3), 567–575. <https://doi.org/10.1007/s13187-019-01666-4.Race>
- 5- Mahmoud, M., Parambil, J., Danjuma, M., Abubeker, I., Najim, M., Ghazouani, H., Al-Mohanadi, D., Al-Mohammed, A., Kartha, A., & Yassin, M. A. (2020). Knowledge, attitude and practice of physicians regarding screening of colorectal cancer in qatar: A cross-sectional survey. Advances in Medical Education and Practice, 11, 843–850. <https://doi.org/10.2147/AMEP.S268315>
- 6- Ammar A Al-Maghrabi. Assessment of the knowledge and awareness of colorectal cancer among general population of Makkah: a survey based study. International Journal of Academic Scientific Research ISSN: 2272-6446 Volume 4, Issue 4 (November - December 2016), PP 168-175 www.ijasjournal.org.

- 7- Khayyat YM, Ibrahim EM. Public awareness of colon cancer screening among the general population: A study from the Western Region of Saudi Arabia. *Qatar Med J*. 2014;2014(1):17-24. Published 2014 Jun 16. doi:10.5339/qmj.2014.3
- 8- Osamah Abdullah A. et al. Public awareness of colorectal cancer screening in Makkah, Saudi Arabia. *International Journal of Medicine in Developing Countries*. 2020;4(1):118–123.
- 9- Gede, N., Reményi Kiss, D. & Kiss, I. Colorectal cancer and screening awareness and sources of information in the Hungarian population. *BMC Fam Pract* 19, 106 (2018).
- 10- Tseng TS, Holt CL, Shipp M, Eloubeidi M, Britt K, Norena M, Fouad MN. Predictors of colorectal cancer knowledge and screening among church-attending African Americans and whites in the deep south. *J Community Health*. 2009;34(2):90–7.
- 11- Papanikolaou IS, Sioulas AD, Kalimeris S, et al. Awareness and attitudes of Greek medical students on colorectal cancer screening. *World J Gastrointest Endosc*. 2012;4(11):513–7.
- 12- Al Eid ZM, AlRadhi ZI, AlTuraiki ZA, AlSalem ES, AlAbbad ZM. Knowledge of Colorectal Cancer Screening among men in AlAhsa, Saudi Arabia. *Int J Sci Eng Res*. 2017;9.
- 13- Tfaily MA, Naamani D, Kassir A, Sleiman S, Ouattara M, Moacdieh MP, et al. Awareness of colorectal cancer and attitudes towards its screening guidelines in Lebanon. *Ann Glob Health*. 2019;85. <https://doi.org/10.5334/aogh.2437>
- 14- Bidouei F, Abdolhosseini S, Jafarzadeh N, Izanloo A, Ghaffarzadehgan K, Abdolhosseini A, et al. Knowledge and perception toward colorectal cancer screening in east of Iran. *Int J Health Policy Manage*. 2014;3:11–5. <https://doi.org/10.15171/ijhpm.2014.48>
- 15- Al-Sharbatti S, Muttappallymyalil J, Sreedharan J and Almosawy Y. Predictors of colorectal cancer knowledge among adults in the United Arab Emirates. *Asian Pac J Cancer Prev*. 2017; 18(9): 2355–2359. DOI: 10.1016/j.ijsu.2016.11.002
- 16- Nemer H, Hejase A, Hejase H, Othman M, Chawraba M and Trad M-A. Colorectal cancer: Exploring awareness in Lebanon. *J of Middle East and North Africa Sciences*. 2016; 2: 10–21. DOI: 10.12816/0032694
- 17- Bidouei F, Abdolhosseini S, Jafarzadeh N, et al. Knowledge and perception toward colorectal cancer screening in east of Iran. *Int J Health Policy Manag*. 2014; 3(1): 11–5. DOI: 10.15171/ijhpm.2014.48
- 18- Honein-AbouHaidar GN, Kastner M and Vuong V, et al. Systematic review and meta-study synthesis of qualitative studies evaluating facilitators and barriers to participation in colorectal cancer screening. *Cancer Epidemiol Biomarkers Prev*. 2016; 25(6): 907–17. DOI: 10.1158/1055-9965.EPI-15-0990
- 19- Lynes K, Kazmi SA, Robery JD, Wong S, Gilbert D and Thaha MA. Public appreciation of lifestyle risk factors for colorectal cancer and awareness of bowel cancer screening: A cross-sectional study. *Int J Surg*. 2016; 36(Pt A): 312–318.

20- Qumseya BJ, Tayem YI, Dasa OY, Nahhal KW, Abu-Limon IM, Hmidat AM, Al-Shareif AF, Hamadneh MK, Riegert-Johnson DL, Wallace MB. Barriers to colorectal cancer screening in palestine: a national study in a medically underserved population. Clin Gastroenterol Hepatol. 2014;12(3):463–469.

UNDER PEER REVIEW