

# **OVERVIEW OF EMULGEL AS EMERGENT TOPICAL DELIVERY: RECENT APPLICATIONS AND ADVANCEMENT**

## **ABSTRACT**

Emulgel is an emerging delivery system with added advantages over other topical delivery systems which is proven to be boon for cosmetics and dermatology. Emulgels are emulsion which is converted to Gel, which will make it a non greasy and enhances patient compliance. The amalgamation of the two systems would be advantages in term of high drug loading capacity and low skin irritation, site specific targeting avoiding first-pass metabolism and increase in bioavailability. Work carried out by researchers proved that, Emulgel/ Nanoemulgel loaded with drugs has been found effective in many topical disorders and it is emerging as potential drug delivery system in area of dermatology. In the present review current research works that carried out on Emulgel and nano emulgels are discussed and also highlighted the recent applications of Emulgel in transdermal as well as dental, ocular, vaginal, and other delivery system.

**KEYWORDS:** Emulgel, Nano Emulgel, Topical delivery system, Emulsion, Dermatology


## **INTRODUCTION**

Recent years witnessed the popularity of transdermal drug delivery systems such as ointments, creams, gels, lotions because they lacks drawbacks associated with oral, parenterals, nasal and other modes of drug delivery [1]. Many disadvantages associated with these traditional types of topical preparations such as stickiness, pourability, lack of spreadibility, stability issue leading to patient incompliance. 1980-90 was the era of popularity of Emulsion and Gel in pharmaceutical topical semisolid dosage form due to its wide applications in dermatology. Emulsions are commonly used for topical pharmaceutical and cosmetic products, such as lotions and creams. Emulsions can be designed to facilitate drug penetration into and/or through the skin. Both oil-in-water and water-in-oil emulsions have been extensively used to deliver drugs and cosmetic agents to the skin, depending on the property of active agents and the indications of the medicines. Emulsions possess certain degree of elegance and get easily washed off. Emulsions have a high ability to penetrate the skin. Regardless of lots of advantages of emulsion, topical application is limited due to its low viscosity and spreadibility.

Gels for topical use have several favorable properties such as **greaseless**, thixotropic, easily spreadable, easily removable, emollient, non-staining, compatible with many additives, and water-soluble or miscible. In spite, so advantageous gels show a major limitation in the delivery of hydrophobic drugs [2]. The majority of the newer drugs developed are lipophilic in nature resulting in unpredictable absorption, poor oral bioavailability, and also variation in pharmacokinetics. Emulgel, an emerging topical delivery system has proved to be boon for the delivery of lipophilic drugs over other formulations.

With the advancement and modernization in transdermal delivery over the years emulgels are gaining interest both in pharmaceutical and cosmetics industry due to greater patient compliance and improved efficacy.

Emulgels are matrices of polymeric gels into which oil phase is incorporated which increases its stability and makes it a dual control release system for hydrophilic and hydrophobic drugs [3], Nanoemulgels are oil-in-water nano emulsions gelled with the use of some gelling agent in it and possess the advantages of both emulsions and gels. Nano emulgels has many advantages such as, transparent, good appearance, greaseless, better spreadability, easily washable, thixotropic, emollient, nonstaining, longer shelf life and. [4], another merits are site specific targeting avoiding first-pass metabolism and ultimately increasing bioavailability as shown in figure 1.



**Figure 1. Advantages of Emulgel**

**Delivery system**

The amalgamation of the two systems makes emulgel formulation advantageous in many ways such as, high drug-loading capacity, low skin irritation, easy incorporation of lipophilic drugs and increase in skin permeability and ultimately, the pharmacokinetic and pharmacodynamic profiles of the lipophilic drugs are improved significantly [5].

The nanoemulsion component of the nanoemulgel helps in protection of drug by preventing the enzymatic degradation. The gel base by increasing the viscosity of the aqueous phase by decreasing the interfacial and surface tension provides thermodynamic stability. Nanoemulgels possess rheological characteristics which are important for transdermal as well as topical delivery such as buccal, dental etc. with the aid of better patient acceptance [6].

In the present scenario, emulgel and nanoemulgel formulation can be considered as potential and promising candidates for topical delivery of lipophilic drugs in the future.

This review aims to introduce the latest trends and applications in transdermal drug delivery via Emulgel for topical and transdermal drug delivery.

### **TOPICAL APPLICATIONS:**

Nanoemulgel preparations of many drugs are being used successfully in different areas of health care such as food supplement, cosmetics, pharmaceuticals and proved its significance in topical and other route of delivery.

Sungpud C, *et al.* [7], evaluated and compared techno-biofunctional characteristics of nanoemulsion and nanoemulgel loaded with mangostin extracts. They concluded that, nanoemulgels had a higher mangostin release than emulgels and presented the feasibility of nanoemulsions and nanoemulgels loaded with mangostin extracts as a promising delivery system for bioactive polyphenol in food supplements, pharmaceuticals and cosmetics. Nanoemulgel preparations of many approved drugs are used in different areas of health care and point out the significance of topical route of delivery as compared to other routes. Torregrosa A, *et al.* [8], developed an metronidazole emulgel for the treatment of rosacea (blushing or flushing and visible blood vessels), applying quality by design (QbD) and found that the emulgel releases metronidazole faster than the creams as it provides greater time of residence over the affected facial area. Anand K, *et al.* [9], reviewed extensively on nanoemulgel and found to be a promising system compared with nanolipoidal formulations. They stated that, Nanoemulgel can be applicable to various acute and chronic diseases through topical routes. Jagdale SC *et al.* [10], formulated emulgel with controlled delivery of mupirocin using Sepineo P 600 for the treatment of skin infection. Antimicrobial and anti-inflammatory study proved successful development of stably controlled release mupirocin emulgel. The optimized batch showed comparable results with marketed formulation against *Staphylococcus aureus*.

Ali Khan B, *et al.* [11], formulated and evaluated the efficacy of topical application of OB-based emulgel on wound healing in animal model. The wound healing effect was evaluated and compared with commercially available Silver Sulfadiazine cream Quench® in wound-induced rabbits by macroscopic and histopathological evidence. Histopathological assessment showed marked improvement in the skin histological architecture after 16 days of OB emulgel treatment. Nordholm-Carstensen A *et al.* [12], compare the standard treatment, diltiazem gel 2%, with Levorag® Emulgel for chronic anal fissures in a single-blinded, randomised, controlled, clinical trial with a non-inferiority design. Patients with a chronic anal fissure were randomised to treatment with diltiazem or Levorag® Emulgel twice daily for 8 weeks. The study demonstrated statistical non-inferiority of Levorag® Emulgel compared with diltiazem in the treatment of chronic anal fissure. Balata GF *et al.* [13], developed an alternative medicine, propolis, in emulgel formulation for burn and wound treatment. The optimized formulation found to be a promising formula for better management of wound and burn.

Mohamed MI, *et al.* [14], formulated zolmitriptan (Zol) niosomal vesicles to potentiate its transdermal effect. The optimized formulation was formulated in Emulgel (1:1 emulsion/gel ratio). Niosomal loaded emulgel showed thixotropic behavior of significant bioavailability and pharmacokinetic parameters as compared to the plain Zt-loaded Emulgel. Surini S, *et al.* [15], prepared Recombinant human epidermal growth factor (rhEGF) loaded transfersomal emulgel with enhanced skin penetration compared with that of non-transfersomal rhEGF emulgels. The skin penetration of rhEGF was enhanced by as much as 5.56 fold by transfersomal emulgel compared with that of non-transfersomal Emulgel.

Shanmugarajan TS, *et al.* [16], developed a squalene loaded emulgel-based system as a approach for potential skin regeneration. They concluded that the squalene-loaded emulgel scaffold could be an

effective formulation used in the treatment of the burnt skin tissue defects. The squalene-loaded emulgel scaffold showed outstanding contraction of wound as compared with the agar gel and negative control. Burki IK, *et al.* [17], designed and developed dexibuprofen-capsaicin emulgel (DC emulgels) for transdermal drug delivery with improved anti-inflammatory and analgesic effects. The DC emulgel showed good physical appearance and stability. The research study also concluded that synergistic potential of DC emulgel could be an alternative to the conventional topical dosage form. Ahmad J, *et al.* [18], designed a thymol loaded nanoemulgel preparation by exploiting low-energy emulsification method for topical application in acne. The study found that the prepared emulgels has improved moisturising properties.

Arshad W, *et al.* [19], developed phyto-cosmetic emulgel of *C. tamala* leaves extract and reported that, as compared to placebo, prepared emulgel was found to be effective in minimizing skin photo-damaging effects. Malavi S, Kumbhar *et al.* [20], developed tretinoin (TRT)-loaded topical emulgel formulation using 32 optimal response surface design (ORS) the optimized TE displayed significant *in vivo* anti-inflammatory activity when compared to marketed gel. Pal RR, *et al.* [21], prepared and evaluated levocetirizine loaded emulgel containing tamanu oil and sericin for atopic dermatitis. *In vivo* pharmacodynamic studies revealed enhanced therapeutic potential of emulgel when compared with conventional gel. Khiljee T, *et al.* [22], formulated a stable *Pyrus communis* fruit extract emulgel which, showed excellent antioxidant and anti-tyrosinase activities. From findings it is concluded that *Pyrus communis* fruit extract loaded emulgel possesses antiaging potential with improvement in skin tone and elasticity. Pinheiro IM *et al.* [23], has developed an emulgel of Amphotericin B and showed potential activity against the *in vitro* parasite with significant reduction of cytotoxicity on murine macrophages, indicating that emulgels formulation is promising for the treatment of cutaneous leishmaniasis.

Coêlho ES, *et al.* [24], developed and characterized a topical emulgel of amphotericin with bacuri butter (*Platonia insignis* Mart.) and evaluate its antileishmanial activity which showed reduction on cytotoxicity to host cells. Sparavigna A, *et al.* [25], evaluated the vasokinetic activity of a Visnadine Emulgel on mucosal genitalia of 15 healthy postmenopausal women clinically. Visnadine Emulgel single application showed significant increase of vulvar hyperemia, along with a significant increase of local turgor versus placebo. Gusai T, *et al.* [26], prepared acyclovir-loaded microsphere-based emulgel to improve its topical delivery. The results of the *ex vivo* permeation study proved significant improvement in drug permeation from optimized microsphere-loaded emulgel compared to the marketed formulation. Mohammadi S, *et al.* [27], compared physicochemical stability and clinical anti-wrinkle efficacy of transdermal emulgel preparations of sodium ascorbyl phosphate (SAP) and ascorbic acid (AA) on human volunteers. Formulations containing (AA) and or (SAP) both improved elasticity and wrinkles of the skin. Tasneem R, *et al.* [28], formulated a stable and effective *Albizia lebbek* bark extract loaded o/w emulsion based emulgel for antioxidant activity and concluded significant antioxidant effects on human skin of prepared emulgels.

Sainy J, *et al.* [29], has developed an Aloe vera-based emulgel formulation of Desoximetasone (DM) for effective treatment of plaque psoriasis and found to increase in *in vitro* permeation of drug from Aloe vera emulgel and could be employed for the topical treatment of skin ailments. de Lafuente Y, *et al.* [30], evaluated the influence of the incorporation of caffeine in bioadhesive emulgel formulation and

results suggested that the optimized emulgel could be an interesting topical bioadhesive delivery system for cosmetic applications. Morsy MA, *et al.* [31], formulated and evaluated the efficacy of topical application of ATR-based nanoemulgel on wound healing. Sithole MN, *et al.* [32], have developed nano-emulsions and nano-emulgels of statins via the transdermal route. The results indicated the benefits of nano-emulsions and nano-emulgels as potential alternative delivery systems for the transdermal delivery of statins. El-Salamouni NS, *et al.* [33], evaluated the efficacy of chamomile oil and nanoemulgel formulations for atopic dermatitis. Treatment with nanoemulgels showed significant decrease in duration of skin healing and no spongiosis compared to chamomile oil. Ijaz S, *et al.* [34], designed the polyphenolic contents of *Mimosa pudica* (MP) seed extract stable topical emulgel formulation and concluded that, the topical emulgel preparation loaded with MP seed extract could be a best option to deal with dematoheliosis.

Meer S, *et al.* [35], investigated the skin depigmenting and antierythemic effects of *Annona muricata* L. fruit extract emulgels and results revealed increase in the skin whitening effects. Hamed R, *et al.* [36], has developed nanoemulsion-based gel (NE gel) formulation of diclofenac diethylamine (DDEA) for topical administration. The release of DDEA from conventional gel, emulgel and NE gel showed a controlled release pattern over 12 h, which was consistent with the rheological properties of the gel. Khan Q, *et al.* [37], Developed dimenhydrinate (DM) emulgel formulation with enhanced permeation for transdermal delivery of DM in treatment of motion sickness. Abdallah MH, *et al.* [38], evaluated a nanoemulgel formulation of Brucine for enhanced topical anti-inflammatory and anti-nociceptive activities. Results suggested that myrrh oil-based nanoemulgel might represent a promising delivery vehicle for potentiating the anti-inflammatory and anti-nociceptive actions of brucine. Mwangi AN, *et al.* [39], formulated and evaluated meloxicam emulgels as a potential alternative topical treatment option for rheumatism. Soliman WE, *et al.* [40], has developed Curcumin (Cur) nanoemulgel for transdermal delivery. The *in vivo* anti-inflammatory activity was estimated using the carrageenan-induced rat hind paw edema method and the results concluded that myrrh oil and Cur-loaded nanoemulgel showed the lowest percent of swelling and thus confirmed the potential of the nanoemulgel dosage form. Sharma G *et al.* [41], prepared emulgel of probiotic, *Bacillus* for topical use. They found faster wound closure and reduction in oxidative stress in comparison to Soframycin®.

Siddiqui B, Rehman AU *et al.* [42], utilises the proficient function of diacerein (DCR) and anti-inflammatory polymers to develop sustained release nanoencapsulated emulgel for potential use in osteoarthritis (OA). Chitosan (CHS) and chondroitin sulphate (CS) were employed as natural anti-inflammatory polymers to encapsulate nanoformulation of DCR. nanoemulgel sustained release of drug having superior penetration properties with provision of enhanced therapeutic effect owing to the presence of CHS, CS, and Argan oil possessing indelible anti-inflammatory attributes. Tuncay Tanriverdi S, *et al.* [43], developed Melatonin-loaded hyaluronic acid (HA) and polyvinyl alcohol (PVA) Freeze-thaw hydrogels and emulgels. *In vitro* release studies showed that the release was improved by emulgel formulations. and concluded that, emulgel formulations could be promising systems for topical application of melatonin. Hariyadi DM, *et al.* [44], investigated and prepared emulgel mask of *C. nucifera* L. Extract (1%) from Kopyor coconut containing carbomer 940 in various

concentration (1% and 1.5%). They reported that, formulation had good spreadability, and also produced highest antibacterial activity against *P. acnes*.

## OTHER ROUTES:

Javed H, *et al.* [45], formulated diphenhydramine nasal nano-emulgels with better penetration for targeted controlled delivery to mucous membrane. The optimized formulation found to have excellent permeability and prolonged residence time on mucosal surface. Salem HF *et al.* [46], formulated nasal nano-emulgel of resveratrol, employing Tween 20, Capryol 90, and Transcutol in emulsion phase and carbopol 934 and poloxamer 407 as the gelling agents. They stated that nano-emulgel has the advantages of both gels and nano along with the intranasal safety and increase in the bioavailability and could be considered as delivery system to target the brain.

Sabry HS, *et al.* [47], formulated an ophthalmic emulgel of combination of betamethasone sodium phosphate and levofloxacin. The new combination in emulgels showed extended release of the two drugs which potentially improve resident time in ocular tissues, patient compliance, and adherence to treatment. Rawoath M, *et al.* [48], developed tamarind gum (TG) and rice bran oil (RBO)-based emulgels of ciprofloxacin HCl and suggested a significantly lower release from the emulgel matrices. The emulgels were able to improve the corneal permeation of drug and thus could be explored to deliver drugs to the internal structures of the eye.

Badawi NM, *et al.* [49], clinically compared the efficiency of metronidazole (MTD) loaded solid lipid nanoparticles (SLNs) vaginal emulgel with the marketed vaginal gel (Metron®) against Bacterial vaginosis (BV). Clinical studies recorded significant enhancement in therapeutic response of MTD from optimized SLNs vaginal emulgel formulation regarding the clinical treatment and low recurrence rate against the marketed product. De Oliveira Neto AS *et al.* [50], formulated atorvastatin loaded oral and vaginal emulgels and evaluated the antifungal activity of atorvastatin and concluded that atorvastatin emulgels could be utilized in treatment of fungal infection. Golshah A, *et al.* [51], investigated the gingivitis effectiveness of Resveratrol emulgel in orthodontic patients and found that the prepared Emulgel was effective in improving gingival health in orthodontic patients and in lowering gingival inflammation. Ferreira SBS, *et al.* [52], prepared curcumin emulgel for the treatment of oral cancer. They developed curcumin emulgels with poloxamer 407, acrylic acid derivatives along with sesame oil or isopropyl myristate as oil phase. They concluded that, the physicochemical properties, subsequent release and permeation of curcumin to selectively kill cancer cells could be improved by the incorporation into emulgel systems. Noronha LL, *et al.* [53], developed itraconazole and clotrimazole emulgels to improve therapeutic effectiveness against *Sporothrix brasiliensis*. They reported that the prepared emulgel found to be good *in vitro* inhibitory activity against *S. brasiliensis* yeasts and suggested that the emulgel containing itraconazole and clotrimazole might be an efficient therapy to oral administration in the treatment of sporotrichosis. Bera H *et al.* [54], developed FLU-loaded emulgel reliable drug carriers for furbiprofen intragastric delivery with a view to improve biopharmaceutical characteristics of drug and modulate its release in a controlled manner. All the formulations demonstrated excellent drug encapsulation efficiency and sustained drug release behavior along with the better *in vitro* gastro retention capabilities. Recent Applications of Emulgel with use and route of application are summarized in Table 1.

**Table 1: Applications of Emulgel with use and Route of application**

Drug	Route	Application	Reference
Mangostin Extracts	Topical Emulgel	Food Supplements, Pharmaceuticals and Cosmetics.	[7]
Metronidazole	Topical Emulgel	Rosacea	[8]
Propolis	Topical Emulgel	Burn and Wound	[13]
Dexibuprofen-Capsaicin	Topical Emulgel	Acne	[17]
C Tamala Leaves Extract	Topical Emulgel	Skin Photo-Damaging Effects	[19]
Tretinoin	Topical Emulgel	Anti-Inflammatory Activity	[20]
Tamanu Oil And Sericin	Topical Emulgel	Atopic Dermatitis	[21]
Pyrus Communis Fruit Extract	Topical Emulgel	Antioxidant and Anti-Tyrosinase Activities	[22]
Amphotericin B	Topical Emulgel	Cutaneous Leishmaniasis.	[23]
Amphotericin With Bacuri Butter	Topical Emulgel	Antileishmanial Activity	[24]
Visnadine	Topical Emulgel	Postmenopausal Superficial Dyspareunia:	[25]
Acyclovir	Topical Emulgel	Cold Sores	[26]
Oidium Ascorbyl Phosphate & Ascorbic Acid	Topical Emulgel	Antioxidant Activity	[27]
Desoximetasone	Topical Emulgel	Plaque Psoriasis	[29]
Caffeine	Topical Emulgel	Cosmetic Applications	[30]
Atorvastatin	Topical Emulgel	Wound Healing.	[31]
Chamomile Oil	Topical Emulgel	Atopic Dermatitis	[33]
Mimosa Pudica (Mp) Seed Extract	Topical Emulgel	Dematoheliosis	[34]
Annona Muricata L. Fruit Extract	Topical Emulgel	Cosmetic	[35]
Diclofenac Diethylamine	Topical Emulgel	Analgesic	[36]
Brucine	Topical Emulgel	Anti-Inflammatory and Anti-Nociceptive Activities	[38]
Meloxicam	Topical Emulgel	Rheumatism	[39]
Curcumin	Topical Emulgel	Anti-Inflammatory Activity	[40]
Probiotic , ( <i>Bacillus</i> )	Topical Emulgel	Faster Wound Closure and Reduction in Oxidative Stress	[41]
Diacerein	Topical Emulgel	Osteoarthritis	[42]
Melatonin	Topical Emulgel	An Anti-Aging and Skin Protective Agent.	[43]
<i>C. Nucifera</i> L. Extract	Topical Emulgel	Antibacterial Activity Against <i>P. Acnes</i> .	[44]
Diphenhydramine	intranasal	Nasal Congestion Due to Allergies.	[45]

Resveratrol	intranasal	Brain Targeted Delivery	[46]
Betamethasone Sodium Phosphate and Levofloxacin	Ophthalmic	Ocular Drug Delivery	[47]
Ciprofloxacin Hcl	Ophthalmic	Corneal Permeation	[48]
Metronidazole	vaginal emulgel	Enhancement in Therapeutic Response	[49]
Atorvastatin	vaginal emulgel	Antifungal Activity	[50]
Resveratrol	Oral	Gingival Inflammation	[51]
Curcumin Emulgel	Oral	Oral Cancer	[52]
Itraconazole And Clotrimazole Emulgels	Topical	Sporothrix Brasiliensis	[53]
Furbiprofen	Topical	Arthritis	[54]
Ofloxacin	Topical Emulgel	Site Specific Delivery	[55]

## CONCLUSIONS

Emulgels are greaseless, easy spreadable, removable, thyrotrophic, good appearance, and transparent. In current scenario the emulgels are being used for the delivery of many drugs like analgesics, anti-inflammatory, anti-acne and anti-fungal. Hence, it is of great pharmacological importance and is relatively free of side effects. In the coming years, it will be most commonly used because it is easy to use and enhances patient compliance. After a thorough literature survey, we can conclude that emulgels has proven to be one of the most effective, better and convenient drug delivery systems as emulgels has the ability to deliver hydrophobic as well as hydrophilic drugs. Therefore, this novel transdermal dosage form is open for research to target specific dermatological disorders as well as in the improvement of various systemic ailments.

## REFERENCES

- [1] Rehman K, Zulfakar MH. Recent advances in gel technologies for topical and transdermal drug delivery. *Drug Dev Ind Pharm*. 2014;40(4):433-440. doi:10.3109/03639045.2013.828219
- [2] Panwar A, Upadhyay N, Bairagi, Gujar, S et al. Emulgel: a review. *Asian J Pharm Life Sci*, 2011; 2231, 4423.
- [3] Ajazuddin, Alexander A, Khichariya A, et al.. Recent expansions in an emergent novel drug delivery technology: Emulgel. *J Control Release*. 2013;171(2):122-132. doi:10.1016/j.jconrel.2013.06.030
- [4] Ashara KC, Paun JS, Soniwala MM, Chavda JR, Mendapara VP, Mori NM. Microemulgel: an overwhelming approach to improve therapeutic action of drug moiety. *Saudi Pharm J*. 2016;24(4):452-457. doi:10.1016/j.jsps.2014.08.002
- [5] Sengupta, P., & Chatterjee, B. (2017). Potential and future scope of nanoemulgel formulation for topical delivery of lipophilic drugs. *International journal of pharmaceutics*, 526(1-2), 353-365.
- [6] Aithal GC, Narayan R, Nayak UY. Nanoemulgel: A Promising Phase in Drug Delivery. *Curr Pharm Des*. 2020;26(2):279-291. doi:10.2174/1381612826666191226100241

- [7] Sungpud C, Panpipat W, Chaijan M, Sae Yoon A. Techno-biofunctionality of mangostin extract-loaded virgin coconut oil nanoemulsion and nanoemulgel. *PLoS One*. 2020;15(1):e0227979. Published 2020 Jan 29. doi:10.1371/journal.pone.0227979
- [8] Torregrosa A, Ochoa-Andrade AT, Parente ME, Vidarte A, Guarinoni G, Savio E. Development of an emulgel for the treatment of rosacea using quality by design approach. *Drug Dev Ind Pharm*. 2020;46(2):296-308. doi:10.1080/03639045.2020.1717515
- [9] Anand K, Ray S, Rahman M, et al.. Nano-emulgel: Emerging as a Smarter Topical Lipidic Emulsion-based Nanocarrier for Skin Healthcare Applications. *Recent Pat Antiinfect Drug Discov*. 2019;14(1):16-35. doi:10.2174/1574891X14666190717111531
- [10] Jagdale SC, Kothekar PV. Development of Emulgel Delivery of Mupirocin for Treatment of Skin Infection. *Recent Pat Antiinfect Drug Discov*. 2020;15(2):137-156. doi:10.2174/1386207323999200819153404
- [11] Ali Khan B, Ullah S, Khan MK, Alshahrani SM, Braga VA. Formulation and evaluation of *Ocimum basilicum*-based emulgel for wound healing using animal model. *Saudi Pharm J*. 2020;28(12):1842-1850. doi:10.1016/j.jsps.2020.11.011
- [12] Nordholm-Carstensen A, Perregaard H, Wahlstrøm KL, Hagen KB, Hougaard HT, Krarup PM. Treatment of chronic anal fissure: a feasibility study on Levorag® Emulgel versus Diltiazem gel 2. *Int J Colorectal Dis*. 2020;35(4):615-621. doi:10.1007/s00384-020-03515-z
- [13] Balata GF, Shamardl HE, Abd Elmoneim HM, Hakami AA, Almodhwahi MA. Propolis emulgel: a natural remedy for burn and wound. *Drug Dev Ind Pharm*. 2018;44(11):1797-1808. doi:10.1080/03639045.2018.1496449
- [14] Mohamed MI, Abdelbary AA, Kandil SM, Mahmoud TM. Preparation and evaluation of optimized zolmitriptan niosomal emulgel. *Drug Dev Ind Pharm*. 2019;45(7):1157-1167. doi:10.1080/03639045.2019.1601737
- [15] Surini S, Leonyza A, Suh CW. Formulation and *In Vitro* Penetration Study of Recombinant Human Epidermal Growth Factor-Loaded Transfersomal Emulgel. *Adv Pharm Bull*. 2020;10(4):586-594. doi:10.34172/apb.2020.070
- [16] Shanmugarajan TS, Selvan NK, Uppuluri VNVA. Development and Characterization of Squalene-Loaded Topical Agar-Based Emulgel Scaffold: Wound Healing Potential in Full-Thickness Burn Model. *Int J Low Extrem Wounds*. 2021;20(4):364-373. doi:10.1177/1534734620921629
- [17] Burki IK, Khan MK, Khan BA, Uzair B, Braga VA, Jamil QA. Formulation Development, Characterization, and Evaluation of a Novel Dexibuprofen-Capsaicin Skin Emulgel with Improved *In Vivo* Anti-inflammatory and Analgesic Effects. *AAPS PharmSciTech*. 2020;21(6):211. Published 2020 Jul 31. doi:10.1208/s12249-020-01760-7
- [18] Ahmad J, Gautam A, Komath S, Bano M, Garg A, Jain K. Topical Nano-emulgel for Skin Disorders: Formulation Approach and Characterization. *Recent Pat Antiinfect Drug Discov*. 2019;14(1):36-48. doi:10.2174/1574891X14666181129115213
- [19] Arshad W, Khan HMS, Akhtar N, Nawaz M. Assessment of changes in biophysical parameters by dermocosmetic emulgel loaded with Cinnamomum tamala extract: A split-faced and placebo-controlled study. *J Cosmet Dermatol*. 2020;19(7):1667-1675. doi:10.1111/jocd.13198

- [20] Malavi S, Kumbhar P, Manjappa A, Disouza J, Dwivedi J. Emulgel for improved topical delivery of Tretinoin: Formulation design and characterization [published online ahead of print, 2021 May 21]. *Ann Pharm Fr.* 2021;S0003-4509(21)00073-0. doi:10.1016/j.pharma.2021.05.004
- [21] Pal RR, Parashar P, Singh I, Saraf SA. Tamanu oil potentiated novel sericin emulgel of levocetirizine: repurposing for topical delivery against DNCB-induced atopic dermatitis, QbD based development and *in vivo* evaluation. *J Microencapsul.* 2019;36(5):432-446. doi:10.1080/02652048.2019.1637474
- [22] Khiljee T, Akhtar N. Investigation of antiaging and skin rejuvenation potential of phyto-constituents from *Pyrus communis* loaded topical emulgel. *Pak J Pharm Sci.* 2019;32(1(Supplementary)):293-300.
- [23] Pinheiro IM, Carvalho IPS, Neto JAT, *et al.*. Amphotericin B-Loaded Emulgel: Effect of Chemical Enhancers on the Release Profile and Antileishmanial Activity In Vitro. *AAPS PharmSciTech.* 2019;20(3):122. Published 2019 Feb 25. doi:10.1208/s12249-019-1323-1
- [24] Coêlho ES, Lopes GLN, Pinheiro IM, *et al.*. Emulgel based on amphotericin B and bacuri butter (*Platonia insignis* Mart.) for the treatment of cutaneous leishmaniasis: characterization and in vitro assays. *Drug Dev Ind Pharm.* 2018;44(10):1713-1723. doi:10.1080/03639045.2018.1492610
- [25] Sparavigna A, Caputo A, Natoli A, Mandelli C. A randomized single-blind placebo-controlled study of a Visnadine Emulgel formulation on healthy postmenopausal women. *Minerva Ginecol.* 2019;71(5):353-358. doi:10.23736/S0026-4784.19.04398-3
- [26] Gusai T, Dhavalkumar M, Soniwala M, Dudhat K, Vasoya J, Chavda J. Formulation and optimization of microsphere-loaded emulgel to improve the transdermal application of acyclovir- a DOE based approach. *Drug Deliv Transl Res.* 2021;11(5):2009-2029. doi:10.1007/s13346-020-00862-w
- [27] Mohammadi S, Shokri J, Ranjkesh M, Akbari Hamed S, Monajjemzadeh F. Comparative physicochemical stability and clinical anti-wrinkle efficacy of transdermal emulgel preparations of 5% sodium ascorbyl phosphate and or ascorbic acid on human volunteers. *J Cosmet Dermatol.* 2021;20(1):174-180. doi:10.1111/jocd.13471
- [28] Tasneem R, Khan HMS, Zaka HS, Khan P. Development and cosmeceutical evaluation of topical emulgel containing *Albizia lebeck* bark extract [published online ahead of print, 2021 May 22]. *J Cosmet Dermatol.* 2021;10.1111/jocd.14244. doi:10.1111/jocd.14244
- [29] Sainy J, Atheriya U, Kori JL, Maheshwari R. Development of an *Aloe vera*-based Emulgel for the Topical Delivery of Desoximetasone. *Turk J Pharm Sci.* 2021;18(4):465-475. doi:10.4274/tjps.galenos.2020.33239
- [30] de Lafuente Y, Ochoa-Andrade A, Parente ME, Palena MC, Jimenez-Kairuz AF. Preparation and evaluation of caffeine bioadhesive emulgels for cosmetic applications based on formulation design using QbD tools. *Int J Cosmet Sci.* 2020;42(6):548-556. doi:10.1111/ics.12638
- [31] Morsy MA, Abdel-Latif RG, Nair AB, *et al.*. Preparation and Evaluation of Atorvastatin-Loaded Nanoemulgel on Wound-Healing Efficacy. *Pharmaceutics.* 2019;11(11):609. Published 2019 Nov 13. doi:10.3390/pharmaceutics11110609

- [32] Sithole MN, Marais S, Maree SM, Du Plessis LH, Du Plessis J, Gerber M. Development and characterization of nano-emulsions and nano-emulgels for transdermal delivery of statins. *Expert Opin Drug Deliv.* 2021;18(6):789-801. doi:10.1080/17425247.2021.1867533
- [33] El-Salamouni NS, Ali MM, Abdelhady SA, Kandil LS, Elbatouti GA, Farid RM. Evaluation of chamomile oil and nanoemulgels as a promising treatment option for atopic dermatitis induced in rats. *Expert Opin Drug Deliv.* 2020;17(1):111-122. doi:10.1080/17425247.2020.1699054
- [34] Ijaz S, Shoaib Khan HM, Anwar Z, Talbot B, Walsh JJ. HPLC profiling of Mimosa pudica polyphenols and their non-invasive biophysical investigations for anti-dermatoheliotic and skin reinstating potential. *Biomed Pharmacother.* 2019;109:865-875. doi:10.1016/j.biopha.2018.10.089
- [35] Meer S, Akhtar N. Annona muricata extract containing pharmaceutical emulgels with and without penetration enhancer for depigmenting and antierythmic effects. *Pak J Pharm Sci.* 2018;31(6 (Supplementary)):2683-2688.
- [36] Hamed R, Basil M, AlBaraghthi T, Sunoqrot S, Tarawneh O. Nanoemulsion-based gel formulation of diclofenac diethylamine: design, optimization, rheological behavior and in vitro diffusion studies. *Pharm Dev Technol.* 2016;21(8):980-989. doi:10.3109/10837450.2015.1086372
- [37] Khan Q, Shah SNH, Arshad MS, *et al.*. Formulation and optimization of dimenhydrinate emulgels for topical delivery using response surface methodology. *Pak J Pharm Sci.* 2021;34(1(Supplementary)):245-255.
- [38] Abdallah MH, Abu Lila AS, Unissa R, Elsewedy HS, Elghamry HA, Soliman MS. Preparation, characterization and evaluation of anti-inflammatory and anti-nociceptive effects of brucine-loaded nanoemulgel. *Colloids Surf B Biointerfaces.* 2021;205:111868. doi:10.1016/j.colsurfb.2021.111868
- [39] Mwangi AN, Njogu PM, Maru SM, *et al.*. Meloxicam emulgels for topical management of rheumatism: Formulation development, *in vitro* and *in vivo* characterization. *Saudi Pharm J.* 2021;29(4):351-360. doi:10.1016/j.jsps.2021.03.005
- [40] Soliman WE, Shehata TM, Mohamed ME, Younis NS, Elsewedy HS. Enhancement of Curcumin Anti-Inflammatory Effect via Formulation into Myrrh Oil-Based Nanoemulgel. *Polymers (Basel).* 2021;13(4):577. Published 2021 Feb 14. doi:10.3390/polym13040577
- [41] Sharma G, Sharma M, Sood R, *et al.*. Self-preserving gelatin emulgel containing whole cell probiotic for topical use: preclinical safety, efficacy, and germination studies [published online ahead of print, 2021 Jul 26]. *Expert Opin Drug Deliv.* 2021;1-13. doi:10.1080/17425247.2021.1947239
- [42] Siddiqui B, Rehman AU, Haq IU, Ahmad NM, Ahmed N. Development, optimisation, and evaluation of nanoencapsulated diacerein emulgel for potential use in osteoarthritis. *J Microencapsul.* 2020;37(8):595-608. doi:10.1080/02652048.2020.1829140
- [43] Tuncay Tanrıverdi S, Cheaburu-Yılmaz CN, Carbone S, Özer Ö. Preparation and in vitro evaluation of melatonin-loaded HA/PVA gel formulations. *Pharm Dev Technol.* 2018;23(8):815-825. doi:10.1080/10837450.2016.1268158

- [44] Hariyadi DM, Isnaeni I, Sudarma S, Suciati S, Rosita N. Peel-off emulgel mask of *Cocos nucifera* L. Extract using gelling agent carbomer 940 as antiacne against *Propionibacterium acnes* ATCC 11827. *J Adv Pharm Technol Res.* 2020;11(4):220-225. doi:10.4103/japtr.JAPTR\_51\_20
- [45] Javed H, Shah SNH, Iqbal FM. Formulation Development and Evaluation of Diphenhydramine Nasal Nano-Emulgel. *AAPS PharmSciTech.* 2018;19(4):1730-1743. doi:10.1208/s12249-018-0985-4
- [46] Salem HF, Kharshoum RM, Abou-Taleb HA, Naguib DM. Nanosized nasal emulgel of resveratrol: preparation, optimization, *in vitro* evaluation and *in vivo* pharmacokinetic study. *Drug Dev Ind Pharm.* 2019;45(10):1624-1634. doi:10.1080/03639045.2019.1648500
- [47] Sabry HS, Al-Shohani ADH, Mahmood SZ. Formulation and Evaluation of Levofloxacin and Betamethasone Ophthalmic Emulgel. *J Pharm Bioallied Sci.* 2021;13(2):205-211. doi:10.4103/jpbs.JPBS\_338\_20
- [48] Rawoath M, Qureshi D, Hoque M, *et al.*. Synthesis and characterization of novel tamarind gum and rice bran oil-based emulgels for the ocular delivery of antibiotics. *Int J Biol Macromol.* 2020;164:1608-1620. doi:10.1016/j.ijbiomac.2020.07.231
- [49] Badawi NM, Elkafrawy MA, Yehia RM, Attia DA. Clinical comparative study of optimized metronidazole loaded lipid nanocarrier vaginal emulgel for management of bacterial vaginosis and its recurrence. *Drug Deliv.* 2021;28(1):814-825. doi:10.1080/10717544.2021.1912211
- [50] de Oliveira Neto AS, Souza ILA, Amorim MES, *et al.*. Antifungal efficacy of atorvastatin-containing emulgel in the treatment of oral and vulvovaginal candidiasis. *Med Mycol.* 2021;59(5):476-485. doi:10.1093/mmy/myaa071
- [51] Golshah A, Mirzaeei S, Nikkerdar N, Ghorbani F. Gingivitis Effectiveness of Emulgel Containing 2% Resveratrol in Orthodontic Patients: An 8-Week Randomized Clinical Trial. *Int J Dent.* 2021;2021:6615900. Published 2021 Mar 29. doi:10.1155/2021/6615900
- [52] Ferreira SBS, Slowik KM, Castro Hoshino LV, *et al.*. Mucoadhesive emulgel systems containing curcumin for oral squamous cell carcinoma treatment: From pre-formulation to cytotoxicity in tissue-engineering oral mucosa. *Eur J Pharm Sci.* 2020;151:105372. doi:10.1016/j.ejps.2020.105372
- [53] Noronha LL, Ferreira PG, G S Lima C, *et al.*. Formulation and Evaluation of a Novel Itraconazole-Clotrimazole Topical Emulgel for the Treatment of Sporotrichosis. *Curr Pharm Des.* 2020;26(14):1566-1570. doi:10.2174/1381612826666200406081249
- [54] Bera H, Nadimpalli J, Kumar S, Vengala P. Kondogogu gum-Zn<sup>+2</sup>-pectinate emulgel matrices reinforced with mesoporous silica for intragastric furbiprofen delivery. *Int J Biol Macromol.* 2017;104(Pt A):1229-1237. doi:10.1016/j.ijbiomac.2017.07.027
- [55] Jagdale S, Pawar S. Gellified Emulsion of Ofloxacin for Transdermal Drug Delivery System. *Adv Pharm Bull.* 2017;7(2):229-239. doi:10.15171/apb.2017.028