

**Protocol on Comparative efficacy of *Bruhatyadi Kwath* as compared to Furosemide for improving e –GFR and Albuminuria in Chronic Kidney Disease-A Randomized controlled study**

**ABSTRACT:**

**Background:** The steady decrease of kidney function is referred to as chronic kidney disease. The kidney function is measured by **Glomerular** Filtration Rate (GFR). According to Ayurveda the CKD can be correlated to Mutraghata because of similarity of symptoms. In Ayurveda Mutraghata is described under Mutrarogawhich comes under Mutravahastrotas (urinary system). **Objective** – To **assess** and compare the efficacy of Bruhatyadikwath, Furosemide and Bruhatyadikwath along with Furosemide on symptoms, eGFR, and Albuminuria in various stages of Chronic Kidney Disease. **Methodology** – Total 90 patients will be divided in 3 equal groups. Patients in group A will be treated with Bruhatyadikwatha, group B patients will be treated with Furosemide and patients in group C will be treated with Bruhatyadikwatha and Furosemide for 90 days. Follow up will be taken after every 30 days. **Expected results** – Furosemide along with Bruhatyadikwath will show better improvement in e-GFR and Albuminurea as compared to only treated with Bruhatyadikwath and Furosemide. Assessment of subjective criteria like edema, anorexia, weakness and vomiting will be done on day 0, 30, 60 and 90 whereas assessment of Serum creatinine, Blood Urea, Sr. Sodium, Sr. Potassium, eGFR (Cockcroft formula) and Albuminuria will be done before and after treatment (on 0 and 90th day). Result: Subjective and objective outcomes will be assessed by statistical analysis. **Conclusion:** It will be drawn from the result obtained.

**Keywords** – Albuminuria; *Bruhtyadikwath*; e- GFR; CKD; Furosemide; *Mutraghat*.

**INTRODUCTION–**

Chronic Kidney Disease (CKD) is a progressive, permanent decline in kidney function that usually occurs over months to years. It begins as a biochemical aberration, but when the kidney's excretory, metabolic, and endocrine functions deteriorate, clinical signs and symptoms of renal failure, commonly known as uraemia, develop. End Stage Renal Disease (ESRD) is a term used to describe a condition in which mortality is likely without RRT (Renal Replacement Therapy) (CKD stage 5)<sup>1</sup>.

Chronic Kidney Disease (CKD) is becoming a major chronic disease worldwide. One reason is that the global incidence of diabetes and hypertension is quickly growing. Given India's population of over one billion people, the increased frequency of CKD is projected to cause severe challenges in the future for both healthcare and the economy. Indeed, the age-adjusted incidence rate of end-stage renal illness in India has recently been estimated to be 229 per million people, with more than 100,000 additional patients entering renal replacement programmes each year<sup>2</sup>. Only 10% of Indian patients with end-stage renal disease receive any renal replacement therapy due to a lack of funding.

Therefore, exploration of a safe and alternative cost-effective therapy is highly required, which proves to be helpful in reducing requirement or frequency of dialysis and in postponing the renal transplantation.

Chronic kidney disease (CKD) is a kind of kidney disease in which kidney function gradually deteriorates over months to years.

Kidney function is a measure of the kidney's health and its contribution to renal physiology.

The Glomerular Filtration Rate (GFR) is a metric for measuring kidney function.

The Glomerular Filtration Rate (GFR) is the rate at which filtered fluid flows through the kidneys. Without Chronic Kidney Disease, a Glomerular Filtration Rate (GFR) of 60 ml/min/1.73 m<sup>2</sup> is considered normal<sup>3</sup>. Chronic Kidney Disease is characterised as having a GFR of less than 60 ml/min/1.73 m<sup>2</sup> for three months. In CKD we get changes in blood, urine and imaging studies. In blood there is raised Serum Creatinine and Urea even than normal. Albuminuria in urine is the oldest and widely used marker for kidney dysfunction, Albumin is the most prevalent plasma protein, and its urine excretion is determined by the combined effects of glomerular filtration and renal tubular processing. It's also used to track CKD progression. On the basis of GFR, Chronic Kidney Disease (CKD) is divided into five stages.

The majority of patients with slowly progressive illness are asymptomatic until their GFR drops below 30 ml/min/1.73 m<sup>2</sup>, and others can be asymptomatic even with significantly lower GFR values.

Symptoms and indicators are typical when GFR goes below 15-20 ml/min/1.73m<sup>2</sup>, and they can influence practically all physiological systems<sup>4</sup>. Tiredness or shortness of breath, as well as lower limb swelling, can all be signs of renal anaemia or fluid overload.

Patients with worsening renal function may have pruritus, anorexia, weight loss, nausea, vomiting, and hiccups.

Due to significant metabolic acidosis, the patient's respiration may be exceptionally deep (Kussmaul breathing) in very severe renal failure and muscle twitching, fits, sleepiness, and coma are all possible side effects<sup>5</sup>.

Ayurvedic management has proved its potential as an alternative medicine for the treatment of a variety of ailments in recent years, and it continues to be a key source for the discovery of new medications, which has gotten more attention recently. Ayurvedic therapy is also becoming more popular for enhancing healthcare and preventing Chronic Kidney Disease, according to data (CKD). Chronic Kidney Disease is not described in *Ayurveda* but due to similarity of symptoms it can be correlated with *Mutraghata*, which is one of the most important *Mutraroga* as described in ancient Samhitas. There are a variety of formulations available that target urinary system problems and have a variety of activities. *Brihatyadikwatha* is one of them described in Sushruta Samhita<sup>8</sup>. It contains *Gokshur*, *Brihati*, *Patha*, *Indrayava*, *Kantakari* and *Yashtimadhu* as shown in table no. 1. All these herbs have *Mutral* (Diuretic) property. In Sushruta Samhita *BruhatyadiGana* is mainly described for the management of *Mutrakruccha* where as in Sahastryogam it is indicated in the management of all *Mutravikara*<sup>6</sup>. It possesses *Rasayana* property which is helpful in regeneration of damaged kidneys. *Deepan-Pachan* property of these drugs reduces production of *Aam* as well as *Kleda*. It corrects *Mansa* and *Medadhatwagni* by its *Katu*, *Tikta Rasa* and *Ushna Veerya* thus reduces production of *Kha-Mala*.

### **Rationale of the study:**

Chronic Kidney Disease (CKD) is a global health problem that costs health systems a lot of money, and it's also a risk factor for cardiovascular disease (CVD).

CKD is linked to an increased risk of cardiovascular morbidity, premature mortality, and/or a lower quality of life at all stages. In modern science management of Chronic Kidney Disease mainly includes supportive treatment, use of diuretics and in severe cases dialysis or renal transplant. But all these treatments have their own limitations like high cost, adverse effects as well as complications. In *Ayurveda* many herbal formulations having *Mutral* (diuretic) and *Rasayana* (rejuvenating) properties are recommended in the management of *Mutravikar* as like *Mutraghata* (Chronic Kidney Disease). Various research studies conducted on herbal drugs in *Mutravikaras* are available showing their efficacy<sup>7,8,9</sup>. *BruhtyadiKwath* possesses diuretic, rejuvenating, antibacterial and anti-inflammatory properties. Research studies are conducted on *Bruhatyadikwath* in the management of *Mutrakruchra* but no study is conducted on *Mutraghata*<sup>10</sup>. So for early stage prevention and to check further progression this study is planned along with Furosemide to evaluate the efficacy of *Bruhtyadikwath* for improving e-GFR and Albuminuria in Chronic Kidney Disease.

### **Aim and Objectives:**

#### **Aim**

Comparative efficacy of *Bruhatyadikwath* as compared to Furosemide for improving e –GFR and Albuminuria in Chronic Kidney Disease.

#### **Objectives:**

##### **(a) Primary objective:**

- To assess the efficacy of *Bruhatyadikwath* on symptoms, eGFR, and Albuminuria in Chronic Kidney Disease.
- To assess the efficacy of *Bruhatyadikwath* with Furosemide on symptoms, eGFR, and Albuminuria Chronic Kidney Disease.
- To assess the efficacy of Furosemide on symptoms, eGFR, and Albuminuria in Chronic Kidney Disease.
- To compare the effect of above therapy in various stages of Chronic Kidney Disease.

#### **Case definition-**

Patients having age  $\geq 20$  to 70 years of either sex having eGFR  $\geq 90$  to 30 (Cockcroft formula) that is Chronic Kidney Disease of stage 1 to 3 with Albuminuria will be included in the study.

#### **Research Question-**

Whether *Brihatyadikwath* along with Furosemide is effective as compared to *Brihatyadikwath* and Furosemide given separately in improving e-GFR and Albuminuria in Chronic kidney Disease?

#### **Hypothesis:**

##### **(a) Null Hypothesis (H0):**

*Bruhatyadikwath* along with Furosemide may not be more effective than *Bruhatyadikwath* and Furosemide given separately in improving the e-GFR and Albuminuria in Chronic Kidney Disease (CKD)

##### **(b) Alternative Hypothesis (H1):**

*BruhatyadiKwath* along with Furosemide may be more effective than *Bruhatyadikwath* and Furosemide given separately in improving the e-GFR and Albuminuria in Chronic Kidney Disease (CKD)

**(c)Component of Hypothesis**

- 1] Estimated Glomerular filtration Rate by Chronic Kidney Disease Epidemiology Equation (eGFR-EPI) and Albuminuria is a useful marker to assess the progress in Chronic Kidney Disease and efficacy of *Bruhatyadikwath*<sup>11</sup>.
  - 2]. Duration of 3 months is sufficient to prove the efficacy of *Bruhatyadikwath*.
  - 3]. Dose of 40 ml twice a day is adequate for the desired effect.
- Trial design: randomized reference controlled clinical trial

**METHODOLOGY:**

**Study setting:**

Patients of Chronic Kidney Disease (CKD) will be selected from OPD and IPD of Mahatma Gandhi Ayurved College, Hospital and Research centre (MGACH & RC), Salod (H), Wardha, and Jawaharlal Nehru Medical College (JNMC), Sawangi (Meghe).

**CTRI Ref. Number:** The clinical study is registered to Clinical Trial Registry and reference no is REF/2021/05/043802

**Study Formulation (*Bruhtyadikwatha*) contents-**

*Bruhtyadikwatha* will be freshly prepared each time by Standard operating procedure as per the reference of *Sharangadhar Samhita (Madhyam Khanda 2/1 and 8/1)*. In coarse powder of raw material 16 parts of water will be added and it will be heated to reduce to 1/8 part. Mixture will be filtered and decoction will be obtained.

**Table no.1 *BruhatyadiKwath* ingredients-**

Sr.	Drug name	Botanical Name	Part Used	Relative Quantity
1.	<i>Bruhati</i>	<i>Solanum indicum</i> Linn.	<i>Mula</i> (Root)	Equal
2.	<i>Kantakari</i>	<i>Solanumsurrattense</i> .Burm.	<i>Phal</i> (Fruit)	Equal
3.	<i>KutajbeejaIndrayava</i>	<i>Holarrhenaantidysentrica</i> .Linn	<i>Beej</i> (seed)	Equal
4.	<i>Patha</i>	<i>Cissampelospariera</i> .Linn	<i>Mula</i> (Root)	Equal
5.	<i>Yastimadhu</i>	<i>Glycyrrhizaglabra</i> Linn.	<i>Kand</i> (Stem)	Equal
6.	<i>Gokshur</i>	<i>Tribulusterrestris</i> .Linn	<i>Phal</i> (Fruit)	Twice of above all

**Eligibility criteria:**

### Inclusion Criteria

1. Age  $\geq$  20 to 70 years
2. Diagnosed patients of Chronic Kidney Disease ( Best on eGFR, and Albuminuria)  
Clinically stable patients of stage 1 to 3 will be included  
Stage 1- GFR  $\geq$ 90 (Cockcroft formula)  
Stage 2- GFR between 60 to 89.  
Stage 3 3a: e GFR (EPI) 45 to  $<$  60  
3b: e GFR (EPI) 30 to  $<$  45
3. Albuminuria.

### Exclusion Criteria

1. Postrenal transplant.
2. Known case of HIV patients or subjects on immune suppressive drugs.
3. Known cases of uncontrolled diabetes mellitus, hypertension, heart diseases, and cancer
3. Pregnant and lactating females.

### 3.4] Interventions of groups: Table no.2

Group	Sample size	Intervention	Dose and frequency	Duration	Follow up
Group-A	30	<i>BruhtyadiKwath</i>	40 ml (BD) Morning & Evening After meal	3 months	Monthly
Group-B	30	Furosemide+ <i>BruhtyadiKwath</i>	40mg OD 10am + 40 ml (BD) Morning & Evening After meal	3 months	Monthly
Group- C	30	Furosemide	40 mg OD 10 am	3months	Monthly

### Criteria for discontinuing or modifying allocated interventions:

1. Discontinuation of Drug during trial.
2. Participants developing life threatening complications.
3. Development of any non related ailments which may require other medications.

**Follow up:** on 30th, 60th and 90th day.

### Primary Outcomes: Outcome of Both the Treatment will be seen in –

- 1.Changes in albuminuria from baseline
- 2.Changes in eGFR from baseline
3. Fraction of patients with improved eGFR of more than 25%
4. Proportion of participants achieving up to 50% reduction from baseline albuminuria.
5. Fraction of patients reverted to stage 2[eGFR(EPI)-60 to 89] from stage 3 [eGFR (EPI)-45 TO 60] or from stage 3b[eGFR(EPI)-30 to  $<$ 45] to stage 3a [eGFR(EPI)-45 to  $<$ 60]

6. Increase in time interval for progression to grade 4 [eGFR(EPI)-15 to 29] from grade 3 or from grade 3a to grade 3b.

**Sample Size :** For calculating sample size with desired error of margin-

$$n = \frac{Z_{\alpha/2}^2 \times P \times (1-P)}{d^2}$$

Where,

$Z_{\alpha/2}$  is the level of significance at 5% i.e 95% confidence interval=1.96

P = Expected Response Rate = 5%= 0.05

D=Derived error of margin=7%=0.06

$$n = \frac{1.96^2 \times 0.05 \times (1-0.05)}{0.06^2}$$

= 25.86  $n = 30$  patients needed in each groups.

**Statistical analysis:** The observations will be analyzed by using chi square test and student unpaired t test.

**Time duration till follow up:** The patient will be treated for a total 90 days and will be followed up on day 30, 60, 90

**Time schedule of enrolment, interventions:** Drug will be given from 0 to 30 days

**Recruitment:** Patients will be recruited by Computerised algorithm for random allocation into three groups of 90 (30 patients in each group).

**5]Methods:**

**5.1] Data collection methods: Assesment criteria:**

**SUBJECTIVE CRITERIA:**

**Gradation of Subjective Criteria Table no.3**

Assessment of subjective criteria like Edema, Anorexia, weakness and vomiting will be done on day 0, 30, 60 and 90.

Sr.No	Symptom	Grade 0	Grade 1	Grade 2	Grade 3
1	Edema	Absent	Mild both ankle feet	Moderate both feet, hands, lower arms and legs	Early morning puffiness
2	Anorexia	Normal instinct for food	Seek for food but refuses	Does not seek for food	Intense havoc n taste dislikes even sight and smell of food

3	Weakness	No	Weakness on strenuous work	Weakness on slight work but relieved soon	Weakness on slight work relieving over period of time
4	Vomiting	Absent	Sensation only for food of dislike	Vomiting few episodes <5	Vomiting episodes >5

**OBJECTIVE CRITERIA:**

1. Serum creatinine.
2. Blood Urea
3. Sr. Sodium
4. Sr. Potassium
5. eGFR (Cockcroft formula)
6. Albuminuria

} will be done before and after treatment

**Data management:** The data entry coding will be done by PI

**Statistical methods:** The observations will be analyzed by using chi square test and student unpaired t test.

**Ethics and dissemination:** Research ethics approval; approval from the research ethics committee has taken. Ref. No.-MGACHRC/IEC/February-2021/188

**Consent or assent:** The written consent will be taken from the patient before starting the study. During the study the confidentiality of each patient will be maintained.

**Dissemination policy:** The data will be disseminated by paper publication.

Guidelines for authorship eligibility, as well as any planned employment of professional writers  
**Informed consent materials:** With all the information consent form and other related documentation will be given to participants.

**Strengths:** Furosemide is a known loop diuretic drug used in CKD, if *Bruhatyadi Kwath* is given along with it may be found effective in alleviating the subjective and objective parameters. So the use of two drugs in combination will have a significant effect on preventing progression of CKD and correcting albuminuria in early stages.

**Limitations:** This study will not be conducted on major systemic diseases like uncontrolled diabetes mellitus, hypertension, malignance and immunocompromised disorders and also in advanced stages of stage 4 and 5.

**Chart 1. Gantt chart (in quarterly based)**

	Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10	Q11
--	----	----	----	----	----	----	----	----	----	-----	-----

Literature search	■											
preparing study material	■	■										
Validation of study material			■	■								
Study intervention					■	■						
Data collection & Analysis							■	■	■	■		
Thesis writing										■	■	
Thesis submission										■	■	

### Discussion-

In Chronic kidney disease, the kidney functions gradually go on decreasing due to various pathologies. It reduces excretion of waste products formed during metabolism, increases leakage of proteins and electrolytes due to damage of nephrons. This loss of proteins, electrolytes depletion and accumulation of nitrogenous waste products produces symptoms. As per Ayurveda, *Mutraghata* means retention of urine along with pain in supra pubic region are observed due to obstructive pathology, there is deranged function of *Vatadosha*, particularly *apanavata* which is the prime causative factor also vitiating *Kaphadosha*, affecting to *mutravahastrotas* and derangements occurs in *Basti* which results in *Mutraghata*. In this study patients will be divided in three equal groups and will be given interventions as shown in table no.2. The contents of *BruhatyadiKwatha* having *Vata-Kapha* pacifying, *Strotoshodhana*, *Shothahara*, *Mutral*, and *Deepana Pachana* (appetizer and digestive) properties which help in breaking *samprapti* of *Mutraghata*. The combination of these drugs possesses the *Rasayana* (rejuvenating) property as it contains *Gokshura* and *Yashtimadhu*<sup>12</sup>. Due to *Rasayana* property it will promote and boost cell growth and their function. It acts as a *Kaphaghna* because of *Tiktakatu rasa*. It also acts as *Kledashoshak* because of the *Grahiguna* of *Indrayava* and *Brihati*. *Srotovishodhana* as it contains *Patha* and *Kantakari* and lastly it has *Mutral* property because of *Gokshur*. In this way *Brihatyadikwath* helps in breaking pathogenesis.

The herbal drugs having *Mutral* property are useful in *Mutraghata* is also proved by various research studies<sup>13,14,15</sup>. A number of studies on different aspects of Chronic Kidney Diseases were reported<sup>16-20</sup>. Few of the related studies were reviewed<sup>21-25</sup>.

Furosemide is a loop diuretic that is used to treat hypertension and edema in people with congestive heart failure, liver cirrhosis, renal illness, and high blood pressure. Diuretics enhance urine salt and water excretion by inhibiting sodium reabsorption in particular renal tubules. It is a drug used in CKD. The subjective criteria as shown in table no. 3 and objective criteria will be assessed before and after treatment.

**Funding** –No Funding

**Conflict of interest**-No any conflict of interest

### References:

1. Davidson's principles practice of medicine, Churchill livinstone Elsevier publications - 23<sup>rd</sup> edition; 2018, page no.384

2. Modi GK, Jha V. The incidence of end-stage renal disease in India: a population-based study. *Kidney international*. 2006 Dec 2;70(12):2131-3.
3. Walker BR, Colledge NR. *Davidson's principles and practice of medicine e-book*. Elsevier Health Sciences; 2013 Dec 6.
4. Sidharth N. Shah *Textbook of API Medicine*, 8<sup>th</sup> edition, volume 1, editor in, published by the association of physician of india, 2008.
5. *Davidson's principles practice of medicine*, Churchill livingstone Elsevier publications - 23<sup>rd</sup> edition; 2018, page no.385
6. ShastriAmbikadatta, "Ayurveda-tattva-sandipika" hindi commentary on *SushrutaSamhita*, Chaukhambha Sanskrit Sansthan Varanasi; reprint edition 2005 p.no.65. *Sushrutasutrasthana* 38/31,32
7. Sadhana Misar (Wajpeyi), Management of Renal Parenchymal Disease in Ayurveda - A Case Study, *International Journal of Science and Research (IJSR)*, February 2017, Volume 6 Issue 2, p.492-495
8. Bhokardankar PS, Rathi B. Indigenous wisdom of Ayurvedic drugs to treat Urinary tract infections. *International Journal of Ayurvedic Medicine*. 2020;11(3):370-7.
9. Rathi B, Khobragade P, Rathi R, Gupta R. Ethno-botanical Survey on Medicinal plants used by Tribes of Karanja (Ghadge) Tahsil of Wardha District, Maharashtra, India. *International Journal of Ayurvedic Medicine*.;12(1):43-52.
10. R. vidyanath, sahastrayogam, mutrakrucchaharkashay chaukhmba Sanskrit pratishthan. 39/6 page no.29
11. C.A. Peralta, M.G. Shlipak, S. Judd, *et al.* Detection of chronic kidney disease with creatinine, cystatin C, and urine albumin-to-creatinine ratio and association with progression to end-stage renal disease and mortality *JAMA*, 305 (2011), pp. 1545-1552
12. Ali et al. [ al-ali m, wahbi s, twaij h, al-badr a. *Tribulusterrestris*: preliminary study of its diuretic and contractile effects and comparison with *zea mays*. *J ethnopharmacol*. 2003;85:257–60. [pubmed] [google scholar].
13. Bhende SV, Parwe S. Ayurveda management of Mutrashmari with special respect to urolithiasis: A case study. *Journal of Indian System of Medicine*. 2019 Jul 1;7(3):189.
14. Hagone PA, Kuchewar V. Management of Glomerulonephritis through Ayurveda – A Case report. *International Journal of Ayurvedic Medicine*, 2019;10(4);345-347
15. Sadavarte A R, Wajpeyi SM , Patil M , Patil A. Study the effect of *Yavakshar Yoga* in the management of *Mutrakrucchra*. *Wutan Huatan Jisuan Jishu*, 2020;16(12),481-496.
16. Agrawal, Amit, Sankalp Dwivedi, Brij Raj Singh, and Pankaj J. Banode. "Simultaneous Hemorrhage in Intracranial Aneurysms and in Renal Cyst in a Case of Polycystic Kidney Disease." *SAUDI JOURNAL OF KIDNEY DISEASES AND TRANSPLANTATION* 23, no. 4 (August 2012): 794–98. <https://doi.org/10.4103/1319-2442.98163>.
17. Aradhey, Parag, Anand Bakre, Sunil Kumar, and Sourya Acharya. "Clinical Profile of Uremic Polyneuropathy in Chronic Kidney Disease Patients." *MEDICAL SCIENCE* 24, no. 102 (April 2020): 945–51.
18. Balwani, Manish, Amit Pasari, Feroz Aziz, Dhananjay Ookalkar, Mohan Patel, Vivek Kute, and Pankaj Shah. "Attitude of the Chronic Kidney Disease Patients Toward Xenotransplantation of Organs." *TRANSPLANTATION* 102, no. 7 (July 2018): S740. <https://doi.org/10.1097/01.tp.0000543729.26735.74>.
19. Balwani, Manish R., Charulata Bawankule, Prakash Khetan, Vishal Ramteke, Priyanka Tolani, and Vivek Kute. "An Uncommon Cause of Rapidly Progressive Renal Failure in

- a Lupus Patient: Pauci-Immune Crescentic Glomerulonephritis.” SAUDI JOURNAL OF KIDNEY DISEASES AND TRANSPLANTATION 29, no. 4 (August 2018): 989–92. <https://doi.org/10.4103/1319-2442.239632>.
20. Balwani, Manish R., Charulata P. Bawankule, Prakash Khetan, and Amit Pasari. “Awareness about Kidney and Its Related Function/Dysfunction in School Going Children: A Survey from the Central India.” SAUDI JOURNAL OF KIDNEY DISEASES AND TRANSPLANTATION 30, no. 1 (February 2019): 202–7.
  21. Balwani, Manish R., Charulata P. Bawankule, Amit Pasari, Priyanka Tolani, Swati Vakil, and Rajashri Yadav. “Minimal Change Disease and Kimura’s Disease Responding to Tacrolimus Therapy.” SAUDI JOURNAL OF KIDNEY DISEASES AND TRANSPLANTATION 30, no. 1 (February 2019): 254–57.
  22. Balwani, Manish R., Amit Pasari, Amol Meshram, Anil Jawahirani, Priyanka Tolani, Hansini Laharwani, and Charulata Bawankule. “An Initial Evaluation of Hypokalemia Turned out Distal Renal Tubular Acidosis Secondary to Parathyroid Adenoma.” SAUDI JOURNAL OF KIDNEY DISEASES AND TRANSPLANTATION 29, no. 5 (October 2018): 1216–19.
  23. Gadewar, Pradyumna, Sourya Acharya, Praveen Khairkar, Samarthi Shukla, and S. N. Mahajan. “Dynamics of Electroencephalogram (EEG) in Different Stages of Chronic Kidney Disease.” JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH 9, no. 3 (March 2015): OC25–27. <https://doi.org/10.7860/JCDR/2015/11257.5705>.
  24. Goswami, Jitendra, Manish R. Balwani, Vivek Kute, Manoj Gumber, Mohan Patel, and Umesh Godhani. “Scoring Systems and Outcome of Chronic Kidney Disease Patients Admitted In Intensive Care Units.” SAUDI JOURNAL OF KIDNEY DISEASES AND TRANSPLANTATION 29, no. 2 (April 2018): 310–17. <https://doi.org/10.4103/1319-2442.229268>.
  25. Lahoti, Sashiraj, Sunil Kumar, and Sachin Agrawal. “Study of Carotid Intimal Medial Thickness in Chronic Kidney Disease at Rural Teaching Hospital.” ANNALS OF MEDICAL AND HEALTH SCIENCES RESEARCH 7, no. 1 (September 2017): 76–80.