

## Original Research Article

### **Estimating the correlation of Ankle Brachial Index with vascular risk factors in Pre-diabetics**

#### **Abstract:**

**Background:** Prediabetes is characterized by a long duration of insulin resistance and increased fasting plasma glucose. Among the various risk factors for cardiovascular atherosclerosis and prediabetes, insulin resistance is one of them. Glycemia has been recognized to affect microvascular damage as compared to macrovascular damage. The ankle-brachial index (ABI) is a simple, affordable and non-invasive test as compared to angiography and has been found to have a high sensitivity and accuracy for the diagnosis of PAD, the gold-standard diagnostic method. This index is based on the fact that in healthy people, systolic blood pressure in the legs is typically equal to or marginally greater than in the upper limbs. This study aims to assess the correlation of ABI with vascular risk factors among subjects with Prediabetes like anthropometric measurement like BMI, Waist hip ratio and neck circumference.

**Methods:** This will be a observational cross sectional study. The present research will involve a cluster of prediabetic subjects. ABI will be determined for all patients. Anthropometric measurements will be taken and blood investigations will be conducted. All data will be compiled and analyzed with appropriate statistical tests.

**Expected results:** We expect a significant association between prevalence of low ABI in prediabetics and heart problems.

**Conclusion:** This research will highlight the adequate clinical and sociological assessment of pre-diabetes patients, including those who have been diagnosed for a long period of time.

**Keywords:** Prediabetics, Ankle Brachial Index, , Cardiovascular risk, Peripheral Arterial Disease.

#### **Introduction:**

Prediabetes is characterized by a long duration of insulin resistance during which at the cost of fasting and postprandial hyperinsulinemia, a compensatory rise in the activity of pancreatic cells preserves normal glycemic that later progresses to increase the state of fasting plasma glucose.<sup>1</sup> Among the various risk factors for cardiovascular atherosclerosis and prediabetes, insulin resistance is one of them. Glycemia has been recognized to microvascular damage as compared to macrovascular damage,<sup>2-3</sup> but the latter was also seen to be associated as shown in current literature.<sup>3</sup> Heart problems among diabetic subjects is found more among subjects having diabetes for a long period of time, low ABI value as well as poor metabolic control.<sup>4-7</sup> Subjects with low ABI value must be given the

required importance so that diabetes can be diagnosed as early as possible along with its prevention<sup>8-9</sup>.

For the elevated cardiovascular risk, experiments include glucose per se and an abnormality in glucose control rarely occurs in isolation. Instead it appears to cluster with other recognized variables of cardiovascular risk. This hypertension, abdominal obesity, and atherogenic dyslipidemia (low high-density lipoprotein [HDL] cholesterol or high triglycerides [TGs]) clustering of impaired fasting glucose is defined as metabolic syndrome. In metabolic syndrome, pro-inflammatory and prothrombotic factors are usually elevated and are often associated with impaired glucose tolerance<sup>10</sup>.

The ankle-brachial index (ABI) is a simple, affordable and non-invasive test<sup>11-13</sup> as compared to angiography and has been found to have a high sensitivity and accuracy for the diagnosis of PAD, the gold-standard diagnostic method.<sup>10</sup> This index is based on the fact that in healthy people, systolic blood pressure in the legs is typically equal to or marginally greater than in the upper limbs. In the case of arterial stenosis, the pressure drop happens distally from the lesion<sup>14</sup>. Furthermore, low ABI in other sites has a strong association with arterial disease and has been shown to be a good indicator of mortality in the general population<sup>15,16</sup>.

ABI determine the occurrence of asymptomatic peripheral arterial disease (PAD), suggested by ABI values of <0.9, if conducted by qualified health practitioners. Most patients with PAD are asymptomatic and their related cardiovascular risk is as high as in symptomatic situations, which is especially troubling. This is also a quiet disease not only associated with worsening of leg symptoms but also with risk of other arterial territories<sup>17</sup>.

Low ABI was linked with death and cardiovascular outcomes among subjects having pre-diabetes, but only limited consideration was paid to its association with other complications. In addition, several studies have investigated the possibility of both high and low ABI values, but it has been difficult to resolve the correlation between the degree of arterial damage (obstruction or pathological constriction) and complications,<sup>18</sup> in particular with respect to microvascular complications. Moreover, recent studies of patients with asymptomatic PAD involved patients with previous coronary events,<sup>19</sup> but primary prevention is of special significance in prediabetes/diabetes patients, as the incidence of a first cardiovascular accident is linked with a rise in morbidity and mortality relative to the diabetes-free population.<sup>20</sup>

A review was carried out in 2017 by Liliana Chevtchouk et al<sup>21</sup> on the ankle-brachial index and diabetic neuropathy. The authors concluded that in patients with type 1 diabetes and type 2 diabetes, particularly those who are asymptomatic, the ABI test is important. More than 10 years ago, ABI was reformed by a diagnosis, regardless of the presence of neuropathic pain or ischemic signs.

### **Rationale**

Previous studies have indicated that individuals with asymptomatic peripheral vascular disease have involved patients having past heart problems, still primordial cure is of special significance among people having prediabetes/diabetes. This might be due to the fact that incidence of 1st cardiovascular occurrence is correlated with a rise in poor quality of life and death relative to the population with a pre-diabetes/diabetes event. Our research aims to determine the relationship of categorized ABI in patients with pre-diabetes with death, cardiovascular outcomes, and microvascular complications.

### **Objectives**

1. To evaluate the association among ABI & demographics profile and anthropometric measurement like BMI, Waist hip ratio and neck circumference.
2. To evaluate the association among ABI and cardiovascular menace variables viz lipid profile.

### **Methods**

This study will be done in AVBH, Sawangi (Meghe) & it will be an observational cross sectional study. The duration of study will be from 2020 to 2022.

**Study design:** Cross sectional study.

### **Selection of subjects:**

All cases will be selected from the university students, staff and patients attending OPD and inpatients admitted. The present research will involve a cluster of prediabetic subjects. Those subjects having age >18 years and who were diagnosed as having prediabetic condition by the expert doctor.

### **Exclusion criteria:**

Patients with bilateral filariasis, cellulitis, malignancy as well as amputation of lower limb and who not willing to give consent will be excluded from the study.

### **Methods:**

Detailed history and examination of the patients will be undertaken. Family history and previous history of diabetes, H/O smoking and Alcohol intake, medication history will be taken. Assessment of pre-existing comorbidities in patient if any will be enquired by H/O raised blood pressure, malignancy, CKD and CLD.

- a. **Measurement of the ankle-brachial index:** Measurements will be carried out on patients lying down with fully balanced head and heels, i.e. not leaning over the end of the test table and after a 5-10 minute break. Patients would be prohibited from deleterious habits like cigarette smoke, alcohol consumption and severe physical effort two hrs prior to the test. Two Doppler systolic pressure readings, two auscultatory systolic pressure readings and two diastolic pressure readings at the right posterior tibial artery, and the final two auscultatory systolic pressure readings and two diastolic pressure readings at the right posterior tibial artery were obtained consecutively. A handheld 8 MHz Aloka Prosound Alpha-7 (20259721) in Bangalore, India, and a normal mercury sphygmomanometer will be used by the observer for the Doppler measurements. For Doppler calculation, an 8 MHz Doppler probe would be used. Dopplergel from system would be spread over the sensor. The probe will be mounted in the region of the pulse at forty-sixty degree angle to the surface of the skin after the Doppler unit is turned on. The probe then jumped about until it heard a strong signal. The cuff is then steadily inflated up to 20 mmHg above the level of the loss of the flow signal and then slowly deflated to detect the reappearance of the flow signal pressure level. There would be a maximum inflation of 300 mmHg. As noted by the maximum amplitude of the waveform seen on the Doppler system display, the systolic pressure would be reported in both the lower limbs. The tubing position of the cuff should be away from the probe, as this may conflict with the

orientation of the probe. By dividing the average systolic BP of the index ankle artery by the average systolic BP of the index limb, the ABI is determined. We found an ABI score of <0.9 and more than 1.4 as possibly sign of PAD & incompressible arteries respectively, in accordance with existing guidelines.

#### **Laboratory Investigation**

- Complete blood count
- Fasting blood glucose
- HbA1c
- Platelet indices
- Blood lipids: HDL, LDL, Total Triglycerides, Total Cholesterol.

#### **Anthropometric measurements:**

Anthropometric features include BMI, waist hip ratio and neck circumference measurement by standard method.

**ABI:** By dividing the average SBP of the index ankle artery by the average SBP of the index arm, the ABI will be determined. ABI value of more than 1.3, more than 0.90, 0.70-0.89, 0.50-0.69 and <0.05 is considered as non-compressible severely calcified vessel as is the case in old age, normal, mild, moderate and severe disease status respectively.

#### **Sample size**

In testing terminology, a sample is a group of entities, objects or things for evaluation taken from a wider population. To ensure that we can generalize the results from the test study to the population as a whole the survey should be representative of the population.

According to the formula i.e.  $n = z^2 pq / d^2$ ,

P= Prevalence of pre-diabetes=14%.

So minimum sample size required will 184 patients

In this study we will recruit two hundred subjects.

.

**Expected outcome:** We expect this study reveals high prevalence of low ABI in prediabetics which might lead to heart problems. We anticipate our research that ABI lower value i.e.  $\leq 1.00$  might give an indication of diabetes i.e. whether it will happen in future or not. However to validate this findings, further studies will be required in the upcoming time. Also from our findings we recommend that subjects with low ABI value must be given the required importance so that diabetes can be diagnosed as early as possible along with its prevention.

#### **Discussion:**

Prediabetes is a condition where the level of blood sugar is greater than average but not yet elevated enough to be graded as Type 2 DM, and Type 2 diabetes is expected to evolve in 10 years or less without intervention. Tests for all people of any age having obesity along with any other associated risk variables responsible for diabetes must be conducted. Macrovascular disorders and type 2 diabetes, leading causes to microvascular disorder, are the most adverse effects in people with prediabetes. In India, over the last three decades, the incidence of CAD has doubled. ABI is non-invasive and can prove to be a valuable tool both in the detection of possible issues and in the tracking of therapy and its effects,

especially in prediabetes. Using the idea of prediabetes in a term may be a valuable intervention technique for avoiding both macrovascular and microvascular illness.

A review was carried out in 2017 by Liliana Chevtchouk et al<sup>21</sup> on the ankle-brachial index and diabetic neuropathy. The authors concluded that in patients with type 1 diabetes and type 2 diabetes, particularly those who are asymptomatic, the ABI test is important. More than 10 years ago, ABI was reformed by a diagnosis, regardless of the presence of neuropathic pain or ischemic signs.

Hirday Pal Singh Bhinder et al<sup>22</sup> tested 100 prediabetes patients in 2018 and participated in similar numbers of age- and sex-matched monitors. In the common carotid artery, a bilateral IMT examination was performed. The findings showed that about 56 percent of cases were male and 44 percent were female, with 58 percent being male and 42 percent being female in controls. The mean age of the cases was  $45.06 \pm 13.08$  and  $44.15 \pm 13.64$  for the controls. The mean CIMT value was higher for the cases ( $0.79 \pm 0.06$  mm) than for the controls ( $0.72 \pm 0.02$  mm). The discrepancy was found to be statistically important between the two groups ( $P < 0.05$ , S). The authors concluded that the mean CIMT was higher relative to controls in the prediabetes group; however, CIMT was not in the elevated category in both categories. Compared to monitors, body mass index and waist-hip ratio is substantially greater in prediabetes. In contrast to controls, systolic and diastolic blood pressures were considerably higher in prediabetes. In prediabetes, overall cholesterol, low-density lipoprotein (LDL), triglyceride (TG) and rather LDL were substantially higher, and high-density lipoprotein (HDL) in prediabetes was significantly lower relative to controls. According to this model, the highest variability of CIMT was found to be responsible for serum Tg and age, among other cases.

Lia Alves-Cabratosa et al<sup>23</sup> measured values of the ankle-brachial index and the possibility of complications of diabetes mellitus in 2019. The investigators concluded that various chances of type 2 diabetes complications were correlated with the analysed types of ABI in asymptomatic PAD patients in primary cardiovascular prevention. In this group, these results may be useful in improving preventive strategies according to the ABI definition.

T.K Kamble<sup>24</sup> researched the myocardial output index and its association with other cardiovascular risk factors for prediabetes. The findings revealed that body mass index, waist hip ratio, total cholesterol, triglycerides, lipoprotein of low density and lipoprotein of very low density were significantly higher in prediabetes and lipoprotein of high density was significantly lower in prediabetes than regulation. Few of the prediabetes related studies were reported<sup>25-33</sup>.

**Conclusion:** The current research will highlights the adequate clinical and sociological assessment of pre-diabetes patients, including those who have been diagnosed for a long period of time. Our results would indicate that in all patients with pre-diabetes type, ABI calculation is important.

#### References:

1. Standards of Medical Care in Diabetes-2016: summary of revisions. Diabetes Care. 2016;39 Suppl 1:S4-5.
2. Ismail-Beigi F, Craven T, Banerji MA, et al. Effect of intensive treatment of hyperglycaemia on microvascular outcomes in type 2 diabetes: an analysis of the Accord randomised trial. Lancet 2010;376:419–30.

3. Sarwar N, Gao P, Seshasai SRK, et al. Diabetes mellitus, fasting blood glucose concentration, and risk of vascular disease: a collaborative meta-analysis of 102 prospective studies. *Lancet* 2010;375:2215–22.
4. Schramm TK, Gislason GH, Køber L, et al. Diabetes patients requiring glucose-lowering therapy and nondiabetics with a prior myocardial infarction carry the same cardiovascular risk: a population study of 3.3 million people. *Circulation* 2008;117:1945–54.
5. Mondesir FL, Brown TM, Muntner P, et al. Diabetes, diabetes severity, and coronary heart disease risk equivalence: reasons for geographic and racial differences in stroke (REGARDS). *Am Heart J* 2016;181:43–51.
6. Hu FB, Stampfer MJ, Solomon CG, et al. The impact of diabetes mellitus on mortality from all causes and coronary heart disease in women: 20 years of follow-up. *Arch Intern Med* 2001;161:161.
7. Alves-cabrato L, Garcia-gil M, Comas-cuf M, et al. Role of low ankle – brachial index in cardiovascular and mortality risk compared with major risk conditions. *J Clin Med* 2019;8:870.
8. Fowkes FGR, Murray GD, Butcher I, et al. Ankle brachial index combined with Framingham risk score to predict cardiovascular events and mortality: a meta-analysis. *JAMA* 2008;300:197–208.
9. Forés R, Alzamora MT, Pera G, et al. Contribution of the Ankle-brachial index to improve the prediction of coronary risk: the ARTPER cohort. *PLoS One* 2018;13:1–12.
10. Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Executive summary of the third report of the national cholesterol education program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult treatment panel III). *JAMA* 2001;285:2486–97.
11. Fishbane S, Youn S, Flaster E, Adam G, Maesaka JK. Ankle-arm blood pressure index as a predictor of mortality in hemodialysis patients. *Am J Kidney Dis.* 1996;27(5):668–72.
12. Ono K, Tsuchida A, Kawai H, Matsuo H, et al. Ankle-brachial blood pressure index predicts all-cause and cardiovascular mortality in hemodialysis patients. *J Am Soc Nephrol.* 2003;14(6):1591–8.
13. Jimenez ZN, Pereira BJ, Romão JE Jr, et al. Ankle-brachial index: a simple way to predict mortality among patients on hemodialysis—a prospective study. *PLoS One.* 2012;7(7):e42290.
14. Donnelly R, Hinwood D, London NJ. Noninvasive methods of arterial and venous assessment. *BMJ.* 2000;320(7236):698–701.
15. Newman AB, Shemanski L, Manolio TA, et al. Ankle-arm index as a predictor of cardiovascular disease and mortality in the cardiovascular health study. The Cardiovascular Health Study Group. *Arterioscler Thromb Vasc Biol.* 1999;19(3):538–45.
16. Vogt MT, Cauley JA, Newman AB, Kuller LH, Hulley SB. Decreased ankle/arm blood pressure index and mortality in elderly women. *JAMA.* 1993;270(4):465–9.
17. Sigvant B, Lundin F, Wahlberg E. The risk of disease progression in peripheral arterial disease is higher than expected: a meta-analysis of mortality and disease progression in peripheral arterial disease. *Eur J Vasc Endovasc Surg* 2016;51:395–403.

18. Meves SH, Diehm C, Berger K, et al. Peripheral arterial disease as an independent predictor for excess stroke morbidity and mortality in primary-care patients: 5-year results of the get ABI study. *Cerebrovasc Dis* 2010;29:546–54.
19. Cardoso CRL, Melo JV, Salles GC, et al. Prognostic impact of the Ankle-brachial index on the development of micro- and macrovascular complications in individuals with type 2 diabetes: the Rio de Janeiro type 2 diabetes cohort study. *Diabetologia* 2018;61:2266–76.
20. Bundó M, Muñoz L, Pérez C, et al. Asymptomatic peripheral arterial disease in type 2 diabetes patients: a 10-year follow-up study of the utility of the ankle brachial index as a prognostic marker of cardiovascular disease. *Ann Vasc Surg* 2010;24:985–93.
21. Chevtchouk L, Silva MH, Nascimento OJ. Ankle-brachial index and diabetic neuropathy: study of 225 patients. *Arquivos de neuro-psiquiatria*. 2017;75(8):533-8.
22. Bhinder HP, Kamble TK. The study of carotid intima-media thickness in prediabetes and its correlation with cardiovascular risk factors. *J Datta Meghe Inst Med Sci Univ* 2018;13:79-82.
23. Alves-Cabratosa L, Comas-Cufí M, Ponjoan A, et al. Levels of ankle-brachial index and the risk of diabetes mellitus complications. *BMJ Open Diab Res Care* 2020;8:e000977.
24. Kamble TK, Kapse A, Kumar S, et al. Study of myocardial performance index in prediabetes and its correlation with other cardiovascular risk factors. *J. Evolution Med. Dent. Sci.* 2020;9(10):721-725.
25. Gupte, Y., S. Acharya, S. Kumar, and P.S. Patil. "Assessment of Endothelial Function by FMD (Flow Mediated Dilatation) in Prediabetes." *International Journal of Pharmaceutical Research* 11, no. 2 (2019): 1808–12. <https://doi.org/10.31838/ijpr/2019.11.02.202>.
26. Raja, K.K., A.H. Inamdar, S. Lahole, and P. Palsodkar. "Prevalence of Non-Alcoholic Fatty Liver Disease in Prediabetes and Diabetes." *International Journal of Pharmaceutical Research* 11, no. 3 (2019): 1424–27. <https://doi.org/10.31838/ijpr/2019.11.03.166>.
27. Rathi, N., B. Taksande, and S. Kumar. "Nerve Conduction Studies of Peripheral Motor and Sensory Nerves in the Subjects with Prediabetes." *Journal of Endocrinology and Metabolism* 9, no. 5 (2019): 147–50. <https://doi.org/10.14740/jem602>.
28. Rai A, Datarkar A, Borle RM. Are maxillomandibular fixation screws a better option than Erich arch bars in achieving maxillomandibular fixation? A randomized clinical study. *Journal of oral and maxillofacial surgery*. 2011 Dec 1;69(12):3015-8.
29. Behere PB, Das A, Yadav R, Behere AP. Religion and mental health. *Indian journal of psychiatry*. 2013 Jan;55(Suppl 2):S187.
30. Agrawal A, Timothy J, Cincu R, Agarwal T, Waghmare LB. Bradycardia in neurosurgery. *Clinical neurology and neurosurgery*. 2008 Apr 1;110(4):321-7.
31. Bourne R, Steinmetz JD, Flaxman S, Briant PS, Taylor HR, Resnikoff S, Casson RJ, Abdoli A, Abu-Gharbieh E, Afshin A, Ahmadi H. Trends in prevalence of blindness and distance and near vision impairment over 30 years: an analysis for the Global Burden of Disease Study. *The Lancet Global Health*. 2021 Feb 1;9(2):e130-43.
32. Borle RM, Nimonkar PV, Rajan R. Extended nasolabial flaps in the management of oral submucous fibrosis. *British Journal of Oral and Maxillofacial Surgery*. 2009 Jul 1;47(5):382-5.

33. Franklin RC, Peden AE, Hamilton EB, Bisignano C, Castle CD, Dingels ZV, Hay SI, Liu Z, Mokdad AH, Roberts NL, Sylte DO. The burden of unintentional drowning: global, regional and national estimates of mortality from the Global Burden of Disease 2017 Study. *Injury prevention*. 2020 Oct 1;26(Supp 1):i83-95.

UNDER PEER REVIEW