

1,3,4-Thiadiazole Derivatives as an Antimicrobial: An Update

Review Article

ABSTRACT

Persistent and uncontrolled use of antibiotics results in development of bacterial resistant. The situation is getting worsen day by day and scientists are investigating thousands of potentially active drugs like molecule in laboratories around the world every day in search of effective antibiotics. During last decade considerable attention was given to five-member heterocyclic moieties while designing new antimicrobial agents. One of important heterocycle is five-membered 1,3,4-thiadiazole with unique bioisosteric properties displaying unusually wide spectrum of biological activities. This comprehensive review represent the recent 1,3,4-thiadiazole and its derivatives, which can be considered as potential antimicrobial agents in the period of 2015 and onwards. This review may help the medicinal chemists to develop new leads possessing 1,3,4-thiadiazole nucleus with higher efficacy and reduced side effects.

Keywords: 1,3,4-thiadiazole; anti-tubercular activity; antimicrobial activity; thiadiazole.

1. INTRODUCTION

Among organic compounds, heterocyclic compounds stood at front for the use of drugs for different biological activities in human and veterinary medicine or as insecticides and pesticides in agriculture [1]. During the past decade, five-membered heterocyclic moieties remain an important target in search of new lead for different therapeutic areas [2]. Among all the heterocycles, five-membered thiadiazole ring system is one of the important heterocycles which comprises different isomers of thiadiazole 1,2,3-thiadiazole, 1,2,4-thiadiazole, 1,2,5-thiadiazole, and 1,3,4-thiadiazole [3]. The hybrid technique is an innovative and strong synthetic tool for the synthesis of two or more different entities in one molecule with novel biological functions, according to the review study. Because of their tremendous potential for displaying notable biological features, -lactams have long been a key component of hybrid molecules. Lactams with four members have been identified as a component of penicillin. The synthesis of -lactams can be done in a variety of ways. For the production of these compounds, the Staudinger reaction of Schiff bases with diphenylketenes is an effective and well-known approach [4]. Several studies have been

published on the topic of hybrid molecules, such as the synthesis of The reaction of readily accessible starting materials such as 4-oxo-4H-chromene-3-carbaldehyde, 1-phenyl-2-(1,1,1-triphenyl-5-phosphanylidene)ethan-1-one, and dibromoformaldoxime under mild conditions in the presence of KHCO_3 yielded a novel, metal-free, and chemo selective approach for 4,5-dihydroisoxazole derivatives [5-6]. The authors described a one-pot sequential four-component reaction of 4-oxo-4H-chromene-3-carbaldehyde/2-chloroquinoline-3-carbaldehyde, 1-phenyl-2-(1,1,1-triphenyl-5-phosphanylidene)ethan-1-one, ninhydrin or isatin, and L-proline in EtOH to obtain a library of diverse polycyclic pyrrolizid Excellent chemical yields, great diastereoselectivity, and operational simplicity are all hallmarks of this procedure [7]. An efficient, practical, and generic approach for the synthesis of [1,8] was explored and discussed by a group of researchers. It is described a one-pot four-component reaction of 2-chloroquinoline-3-carbaldehyde and 1-phenyl-2-(1,1,1-triphenyl-5-phosphanylidene)ethan-1-one, 1,1-bis(methylthio)-2-nitroethylene or ketene N,S-acetals, aromatic/aliphatic amine or diamines, and aromatic/aliphatic amine or diamines under mild and High yields, gentle and catalyst-free conditions, fast reaction durations,

and the use of green solvent are all advantages of this protocol [8].




Fig. 1. Isomers of thiadiazole

Literature survey revealed that among all the isomer, 1,3,4-thiadiazole have received considerable attention and has been because of their broad spectrum of pharmacological activity. 1,3,4-thiadiazole having high aromatic property is a very weak base because of inductive effect imparted by ring sulfur. 1,3,4-thiadiazole is stable in aqueous acid solution and undergoes ring cleavage in aqueous basic solution. Nucleophilic attack is prevalent as ring is electron deficient because of electronegative ring nitrogen nonreactive towards electrophilic attack. Nucleophilic substitution is favored at 2' or 5' position of the ring. 1,3,4-thiadiazole [9,10].




Fig. 2. Numbering of 1,3,4-thiadiazole ring system

Most of the authors assumed that presence of =N-C-S- moiety is responsible for biological potential of 1,3,4-thiadiazole derivatives [11]. While some other authors supposed that great in vivo stability of 1,3,4-thiadiazole is because of its strong aromatic is responsible for biological activity and low toxicity [12].

1,3,4-thiadiazole is the bioisosteres of pyridazine through the substitution of -CH=CH- by -S- (**Fig. 2**) [13, 14]. The thiadiazole derivatives show good lipid solubility attributed to ring sulfur and thereby show oral absorption and good cell permeability leading to a good bioavailability [14-16]. Discovery of heterocyclic sulfonamides established the biological importance of 1,3,4-thiadiazole derivatives [17]. Presently only sulfathiazole is used in clinical practice for the management of *Haemophilus vaginalis* vaginitis [18]. Importance of 1,3,4 thiadiazole highlighted among researchers with report of acetazolamide synthesis as carbonic anhydrase inhibitor used in glaucoma [19], epileptic seizures [20], hemiplegic migraine [21] etc. Its methylated derivative methazolamide is a more potent

carbonic anhydrase inhibitor and displays diuretic, antiglaucoma and potential antineoplastic activity [22], 1,3,4-thiadiazole derivatives displays diverse pharmacological activities including antimicrobial [23], anti-cancer [24], antifungal [25], antituberculosis [26], local anaesthetic [27], antiglaucoma [28], anticonvulsant [29], anti-inflammatory [30], antidepressant and anxiolytic [31], antihypertensive [32], antiviral [33] and antioxidant activity [34].

This review discusses the antimicrobial potential of 1,3,4-thiadiazole which appeared very recently in literature. Special attention is given to anti-tubercular activity and discussed under separate heading.

2. ANTIMICROBIAL ACTIVITIES ASSOCIATED WITH 1,3,4-THIADIAZOLE SYSTEM



Infective diseases caused by pathogenic microorganisms such as bacteria, fungi, viruses, protozoa and helminthes affects millions of people worldwide and results in considerable deaths Interestingly among 1400 different species of microorganisms which are reported in literature only 20 of them (mostly bacteria) accounts for around two thirds of the casualties [35]. Although highly developed countries experiencing fall in death from 16 million in 1990 to approximately 15 million and forecasting 13 million in 2050, death rate is still high in developing countries because of tuberculosis, pneumonia, malaria, HIV/AIDS, diarrhea and many other diseases [36,37]. Powerful immunosuppressive drugs are occasionally prescribed for the management of cancer therapy, organ transplant and spread of HIV infection resulting in increased incidence of fungal infections among immunocompromised patients. Occurrence of systemic fungal infections witnessed sharp rise recently [38,39] Considering the recent pandemic because of COVID-19 and similar pandemic threat in future and issue of dramatic increase in antibiotics resistance, discovering new effective antibacterial/antiviral drugs and the development of modern therapies are two challenges of top importance.

2.1 Antibacterial and Antifungal Activities

Being bioisosteres of the thiazole ring, thiadiazole ring acts as a pharmacophore and is the part of the third- and fourth-generation

cephalosporins, and this observation makes it possible to use it in the synthesis of antimicrobial agents [40] Since the discovery of penicillin in 1942) the race of finding new antibiotics continued and became most intense with time [41]. With the passing time microorganisms are becoming more resistant and invasive which resulted in dramatically increased bacterial infections. On other side systemic fungal infection is now more evident with the use of powerful immunosuppressive drugs for cancer therapy and organ transplants [30,31].

As per reported work, synthesized a new series of 1,3,4-thiadiazol-4,5-dihydropyridazin-3(2H)-ones derivatives and evaluated the compounds








As per reported work, the synthesis of Schiff bases clubbed with 1,3,4-thiadiazole moiety and evaluated them for antimicrobial activity. Antimicrobial assay results indicated that most of the synthesized thiadiazole derivatives exhibited good to excellent antimicrobial activity. Schiff bases **4**, **5**, **6** exhibited excellent bactericidal activities against all the strains with inhibition at MIC 4–16 µg/mL. Compound **7a** and **7b** comprising fluorine atom displayed remarkable inhibition at MIC 4–8 µg/mL against Gram positive bacteria. The entire above compound displayed good antifungal activity against all examined fungal strains at MIC 16–31.5µg/mL [43].

Author have reported the 1,3,4-thiadiazole derivatives of 2-(4-formyl-2-methoxyphenoxy) acetic acid (6a–s) and tested them against bacteria *S. aureus*, *S. enterica*, *V. cholera*, *B. subtilis*, *P. mirabilli*, *E. coli* V517, *M. smegmaticus*,


for antimicrobial potential. Most of the synthesized derivatives were found to be bactericidal. Compounds **1a–e**, **2a–c** and **3** exhibited moderate activity against *B. subtilis*. Compounds **1a**, **1c–e** and **2a–c** were potent against *S. pneumoniae*. Compounds **1a–e** exhibited moderate to potent activity against fungal strains *A. fumigates*, *S. racemosum* and *G. candidum*. Authors observed that compounds bearing only thiadiazole moiety at C-4 of pyridazinone ring were found more active towards all microbial strains except for *P. aeruginosa* and *C. albicans*, and the presence of either carbamoyl or thiocarbamoyl group at N-2 improved the activity [42].

P. aeruginosa and antifungal activity against *C. albicans*. Compound **8b** exhibited excellent activity against all the strains. It showed maximum activity (97%) against *S. enterica* (95%), against *V. cholera* and (87.9%) inhibition of *E. coli* V517 when compared with standard drug ampicillin. Compound **8a**, **8c**, **8d** and **8e** also displayed very good activity. Compound **8b** showed maximum inhibition (87.8%) whereas, compound **8a** showed (83.3%) inhibition against fungal strain *C. albicans* [44].

As per reported in literature, synthesized novel pyridine imidazo [2,1b]-1,3,4-thiadiazole derivatives. When investigated for antimicrobial potential it was revealed that compounds, 4(a), 4(b), 4(f), 4(h) and 4(k) exhibited promising candidates against *B. pumillus*, *S. aureus*, *V. cholera*, *E. coli*, *P. mirabilis*, *P. aeruginosa* and *C. albicans* with reference to standard drugs ampicillin and amphotericin B [45].




8a, R = 3-NO₂; 8b, R = 4-NO₂; 8c, R = 3-OH;
8d, R = 2,4-diOH; 8e, R = 4-NH₂




9a, R = 2,4-di-OH; 9b, R = 2-OH; 9c, R = 4-Cl;
9d, R = 4-NO₂; 9e, R = 3,4-di-CH₃

Substituted benzimidazoles containing 1,3,4-thiadiazole were synthesized and tested for antimicrobial activity. Compound **10** displayed moderate antibacterial activity [46].




Synthesized (1,3,4-thiadiazole)-methylthio-derivatives of 2-mercapto-quinazolin-4-one analogues and investigated them for in vitro DHFR inhibition, antitumor and antimicrobial activity. Compounds **11** exhibited promising activity against the Gram-positive bacteria *Staphylococcus aureus* and also found against

Bacillus subtilis with a comparable potency to the broad spectrum antibiotic Ciprofloxacin [47].



Muglia et al. reported synthesis and antimicrobial screening of disubstituted 1,3,4-thiadiazole derivatives. Some of the synthesized compounds were found active against *Staphylococcus aureus*, *S. typhimurium*, *C. albicans*, *Enterobacter aerogenes*, and *S. Kentucky*. Compound **12a** displayed very good activity against *Staphylococcus aureus* (gram-positive). Compounds **12b**, **12c** and **12d** also


exhibited significant activity against *Staphylococcus aureus* [48].



12a-d


Ar = a = 3-Cl-C₆H₅, b = 4-F-C₆H₅
c = C₆H₅CH₂, d = 4-OCH₃-C₆H₅

In vitro antimicrobial activity of Novel 4-(5-(10-(3-N, N-dimethylamino) propyl)-10H-phenothiazine3-yl)-1, 3, 4-thiadiazole-2-yl) Azo dye/Schiff base derivative was evaluated against several strains. Compound **13** showed promising anti-microbial activity against various pathogenic microorganisms as compared to the antibiotics Ciprofloxacin and Fluconazole [49].




13


New 1,3,4-thiadiazole derivatives based on the key precursor 2,5-bis(mercaptoacetichydrazide)-1,3,4-thiadiazole were reported and tested against *E. coli* and *E. faecalis* strains. The tested compounds exhibited higher to moderate activity in comparison with ceftriaxone at concentration 30 mg/discs. Compounds **14**, **15**, and **17** showed higher activity against bacteria *E. coli*, and compounds **15**, **16**, **17**, and **18** showed higher activity against *Enterococcus bacteria* [50].




14




15



16




18



17

1, 3, 4-thiadiazole-based thioglycosides were synthesized and evaluated for antimicrobial activity against several strains. Among the entire compounds lauric acid and myristic acid derivatives showed good and moderate antimicrobial activity. Compound **19a** and **19b** were found most potent against Gram-negative bacteria strain *Klebsiella pneumonia* with MIC values 12.5, 25 g/mL respectively when compared with penicillin and streptomycin [51].



19a-b

19a, R = -(CH₂)₁₂-CH₃; 19b, R = -(CH₂)₁₄-CH₃

Novel 5-(3,5-dinitrophenyl)-1,3,4-thiadiazole derivatives screened for *in-vitro* antimicrobial activity. Compounds **20**, **21**, **22**, and **23** showed

excellent and broad-spectrum antimicrobial activity comparable to Amoxicillin and Fluconazole as positive antibiotic and antifungal controls, respectively [52].





Synthesized new a series of novel myricetin derivatives containing 1,3,4-thiadiazole and tested for antibacterial activities against Xoo and Rs and their antiviral activity against TMV. Bioassay results indicated that some target compounds exhibited potential antibacterial and antiviral activities.



Among them, compounds **24**, 5,7-Dimethoxy-2-(3,4,5-trimethoxyphenyl)-3-((5-((4-chlorobenzyl)thio)-1,3,4-thiadiazol-2-yl)methoxy)-4H-chromen-4-one exhibited highest activity [53].

Synthesized new 1,2,4-Triazolo [3,4-b][1,3,4]thiadiazole derivatives and evaluated for their in vitro antibacterial and antifungal activities against pathogenic microorganisms using two fold serial dilution method. Compound **25a** was most active

while compounds **25b** and **25c** displayed remarkable antimicrobial activity against all the tested microorganisms comparable to reference drugs gentamicin and miconazole. Docking study revealed that test compound **25e** had lesser estimated binding free energy and predicted inhibitory constant values when compared with fluconazole. A SAR study indicated that the activity was the highest when halogen groups were substituted at the ortho and Meta positions of the first phenyl ring attached to the 3rd position of the triazolothiadiazole nucleus [54].


1,3,4-thiadiazoles derived from 4-phenoxybutyric acid were synthesized and screened tested against gram negative bacteria, gram positive bacteria and for antifungal potential. However, all the tested compounds showed good antimicrobial activities against *S. aureus* only. The highest antimicrobial activity was exhibited by compound **26a**. Compound **26b**, **27** and **28** also exhibited significant antimicrobial activity [55].






25a, $R_1 = H$, $R_2 = Cl$;
 25b, $R_1 = Br$, $R_2 = H$;
 25c, $R_1 = Cl$, $R_2 = H$

25a-c




As per published work, synthesized new series of 1,3,4-thiadiazolyl-sulfanyl-4,5-dihydropyridazin-3(2H)-ones and evaluated them for antimicrobial potential. Compounds **29d**, **30** and **31** exhibited potent activity against *B. subtilis* and found more potent against *S. pneumonia* compared to control drug. Compounds **29a**, **29d**, **30**, and **31** displayed remarkable activity against *E. coli* and their bacteriostatic effect has exceeded over control drug. Compounds **29a-c**, **5**, and **29d** exhibited potent antifungal activity against *A. fumigates*, *S. racemosum*, *G. candidum*. Only compound **29d** was found potent against *C. albicans* strain. From the obtained results it was noted that; pyridazinone rings **29a-d** and **30** resulted from thia-Michael adduct were generally very potent against all bacterial and fungal strains except for *P. aeruginosa* and *C. albicans*. The presence of either carbamoyl or thiocarbamoyl moiety at N-2 significantly increased the activity to precede the control drug [56].




29a-d

29a, R = H; 29b, R = COCH₃ ;
29c, R = CHO, 29d, R = CSNH₂




30



31


A new class of methylthio linked pyrimidinyl 1,3,4-thiadiazoles **32a-f** were synthesized and evaluated for antimicrobial activity. The compounds **32c** and **32f** displayed strong antibacterial activity against *P. aeruginosa* at all tested concentrations. It was observed that electron withdrawing groups on the aromatic ring increased the activity. Results indicated that tested compounds were more susceptible towards the Gram negative bacteria than Gram positive ones. All the compounds exhibited very good antifungal activity against *A.niger* than *P. chrysogenum* [57].




32a-f

R = a) H, b) 4-Me, c) 4-Cl,
d) 4-Br, e) 4-OMe, f) 4-NO₂

Synthesized 2-aryl,5-substituted 1,3,4-thiadiazoles and tested some of the compounds for antibacterial potential. Compound **33a** and **33b** displayed good activity against Gram-positive bacteria *Bacillus megaterium*, *Enterococcus faecalis* and *Streptococcus mutans* MTCC 497) and Gram-negative bacteria *Escherichia coli*, *Proteus vulgaris* and *Pseudomonas aeruginosa* [58].



33a



33b


1,3,4-thiadiazole derivatives containing pyrazole oximes were synthesized and tested against acaricidal and insecticidal activity. Some of the target compounds showed promising acaricidal and insecticidal properties. Importantly, compound **34a** showed 80% acaricidal activity against *Tetranychus cinnabarinus* at the concentration of 50µg/mL and compound **34b** exhibited 100% insecticidal activities against *Aphis craccivora* at the concentration of 50 µg/mL. Moreover 100% insecticidal activities against *Plutella xylostella* was exhibited by compound **34c** and **34d** at the concentration of 50µg/mL. Furthermore, compounds **34c** (LC₅₀ = 19.61 µg/mL) and **34d** (LC₅₀ = 9.78 µg/mL) possessed comparable or even better insecticidal activities than the control Pyridalyl (LC₅₀ = 17.40 µg/mL) against *Plutella xylostella* [59].

Synthesized new 1,3,4-thiadiazole derivatives and screened for antimicrobial activity. Levofloxacin. Compounds **35**, **36a** and **36b** showed very good antimicrobial activity towards Gram-positive (*Rhodopseudomonas* sp., *Bacillus cereus* and *Micrococcus luteus*) and Gram negative (*E. coli* and *Salmonella typhi*) bacteria with MIC values (1- 8) mol mL⁻¹ when compared to Levofloxacin (MIC= 2-5 mol mL⁻¹). Compounds 8, 10a and 10b also exhibited higher antifungal activity against *Alternaria alternate*,



Aspergillus flavus, *Candida albicans* and *Curvularia lunata* comparable to Nystatin [60].

Vanillin derivatives of 1,3,4-thiadiazole were synthesized and screened for antibacterial


activity. Compound **37** displayed strong antibacterial activities against *Xanthomonas oryzae pv. oryzae* (Xoo) and *Xanthomonas oryzae pv. oryzicola* (Xoc) in vitro, with the EC50 values of 3.14 and 8.83 $\mu\text{g/mL}$, respectively [61].




34a; $R_1 = \text{Et}$, $R_2 = \text{Me}$, $R_3 = 4\text{-OCH}_3$,
 34b; $R_1 = \text{Me}$, $R_2 = \text{Me}$, $R_3 = 4\text{-Cl}$,
 34c; $R_1 = \text{Et}$, $R_2 = \text{Me}$, $R_3 = 3\text{-Cl}$,
 34d; $R_1 = \text{Et}$, $R_2 = \text{Me}$, $R_3 = 2,4\text{-diCl}$






36a, $R = \text{C}_2\text{H}_5$;
 36b, $R = \text{C}_6\text{H}_5$;




Reported synthesized several 1,3,4-thiadiazole derivatives of 4-phenoxybutyric acid which displayed strong antimicrobial properties against *S. aureus*. The highest antimicrobial activity was exhibited by compound **38**. It has been observed that the tested compounds exhibited increased potential antimicrobial activities against *S. aureus* [62].



1,3,4-thiadiazole derivatives from N-aminobenzyl or N-arylhydrazone series were synthesized and tested against the *trypomastigote* form of *Trypanosoma cruzi*. Compounds **39**, **40** and **41** exhibited outstanding activity and excellent selectivity indexes when compared to reference drug posaconazole. Compound **39** was also active against the intracellular *amastigote* form of *T. cruzi*. Furthermore, its corresponding hydrochloride, compound **40**, showed the most promising profile, producing phenotypic changes similar to those caused by posaconazole, a well-known inhibitor of sterol biosynthesis. Findings of the study indicated compound 118 as a possible prototype for the development of new drug candidates for Chagas disease therapy [63].






Novel 1,3,4-thiadiazole derivatives **42a-h** were prepared via cyclization reaction of 2-phenylbutyric acid with N-phenylthiosemicarbazide and POCl₃. All the synthesized derivatives were screened against several microbial strains using a disk diffusion method. From the biological finding it was observed that the synthesized 1,3,4-thiadiazole derivatives exhibited antibacterial activity against *S. aureus*, *E. coli*, and *C. albicans* [64].




42a, R = 2-F; 42b, R = 2-OCH₃; 42c, R = 2-CH₃,
 42d, R = 2-F; 42e, R = 2-Cl; 42f, R = 3-OCH₃;
 42g, R = 2-NO₂; 42h, R = H


Synthesis of 1,3,4-thiadiazole derivatives containing N-aryl 2-aryloxyhydrazone-propanehydrazonoyl chlorides. Among them **43**, **44**, **45**, **46**, **47a-b**, **48a-b**, and **49a-b**, were tested for their *in-vitro* antibacterial activity against Gram-positive bacteria (*Staphylococcus pneumoniae* and *Bacillus subtilis*) and Gram-negative bacteria (*Pseudomonas aeruginosa* and *Escherichia coli*). Compounds were also tested for their *in vitro* antifungal activity against fungi species (*Aspergillus fumigatus*, *Geotrichum candidum*, *Candida albicans* and *Syncephalastrum racemosum*). Ampicillin, Gentamicin and Amphotericin B were used as reference drugs. From the data obtained from experiment it was observed that tested compound exhibited moderate to strong activity against *S. pneumoniae*, of *B. subtilis* while have no inhibitory effect toward *P. aeruginosa*. Compounds **45**, **46**, **47b** and **49b** exhibited high inhibitory effects against *E. coli*. Moreover all the derivatives exhibited good activity against all fungi strains [65].



Novel quinazolin-4(3H)-one derivatives containing a 1,2,4-triazolo [3,4-b][1,3,4] thiadiazole were synthesized and tested for antibacterial potential. The experimental data indicated that the compounds **50a**, **50b**, **50c** and **50d** had the EC₅₀ values of 34.8, 28.2, 41.5 and 42.5µg/mL against the phytopathogenic bacterium *Xanthomonas oryzae pv. oryzae* (*Xoo*), respectively, which were significantly better than commercial bactericide Bismethiazol (EC₅₀ =95.8µg/mL) [66].




Highly potent antimicrobial imidazo [2,1-b][1,3,4] thiadiazole derivatives were reported. Compounds **51a-f** were found ten time more potent with MIC values as low as 0.03 mg/ml when compared to control chloramphenicol against Gram-positive (*Staphylococcus aureus* and *Bacillus subtilis*) bacteria and Gram negative (*Escherichia coli*) bacteria. All the compounds fail to show significant antifungal activity. According to the electronic structure calculations, almost all active compounds obey the drug likeness properties [67].




As reported data, synthesis of N-(5-[(1-Phenyl-5-(thiophen-2-yl)-1H-1,2,4-triazol-3-yl)thio]methyl)-1,3,4- thiadiazol-2-yl)thiophene-2-carboxamide **52** and its evaluation for antimicrobial potential. The compound was found

active against gram positive bacteria (*Staphylococcus aureus* and *Bacillus subtilis*) and Gram negative bacteria (*Escherichia coli* and *Pseudomonas aeruginosa*) [68].




Synthesis of novel 1,3,4 thiadiazole derivatives with potent antifungal activity. Compounds **53a** and **53b** showed remarkable antifungal activity against all eight *Candida* species. Compound **53b** was the most effective derivative against *C. albicans* ATCC 10231. It was observed that presence of fluoro and chloro groups at the second position of the phenyl moiety in compounds **53a** and **53b** was responsible for their potent activity. Moreover compound **53a** and **53b** exhibited a good predicted pharmacokinetics profile. Furthermore, when investigate for primary mechanism of action it was revealed that inhibition of ergosterol biosynthesis in *C. Albicans* was the reason behind activity of **53a** and **53b**. In the docking study, significant interactions were observed


between compounds **53a** and **53b** and 14- α -sterol demethylase, which is a key enzyme in ergosterol biosynthesis [69].



New imidazo[2,1-b][1,3,4]thiadiazole derivatives containing benzothiazole moiety were tested against anti-leishmanial and antibacterial activity. Compound **54a** exhibited most potent antileishmanial activity (MIC=10000 $\mu\text{g/mL}$) whereas compound **54b** was found to be effective at the highest concentration studied (MIC=20 000 $\mu\text{g/mL}$). In terms of antibacterial activity, compounds **54b** were found to be the most effective compounds against *Escherichia coli* (MIC = 625 $\mu\text{g/mL}$) and against *Yersinia enterocolitica* (MIC=1 250 $\mu\text{g/mL}$). The docking study revealed that compounds **55**, and **54b** could be new potential inhibitor compounds for the 2eg7 protein structure [70].




New derivatives of 5-(nitroheteroaryl-2-yl)-1,3,4-thiadiazole were tested for leishmanicidal activity. Entire compounds exhibited potent activity against both promastigote and amastigote forms of *Leishmania major* (*L. major*). Compounds, **56** and **57** displayed highest activity. The analysis of redox-related factors indicated that exposure of *L. major* cells to **56** and **57** led to an increase in reactive oxygen species (ROS). Authors concluded that the anti-leishmanial potential of **56** and **57** is mediated by apoptosis through an imbalance in the redox system resulting in the elevation of ROS [71].




Synthesis of N-(5-(pyridin-4-yl)-1,3,4-thiadiazol-2-yl) cinnamamide **58**. The compound was found

moderately active against *Salmonella typhimurium* ATCC14028 [72].




58

Synthesis of new 2-(6-phenylimidazo[2,-1-b][1,3,4]thiadiazol-2-yl)-1H-indoles derivatives and evaluated for their anti-biofilm properties against the Gram-positive and the Gram-negative bacterial strains. Compounds **59a** and **59b** displayed excellent anti-biofilm activity against *S. aureus* ATCC 25923 with BIC50 values of 0.5 and 0.8 mg/ml, respectively, whereas compound **59c** was the most potent against *S. aureus* ATCC 6538, with a BIC50 of 0.3 mg/ml. Notably, these compounds showed effects in the early stages of the biofilm formation without affecting the mature biofilm of the same strains and the viability of the planktonic form. Author proposed the derivatives as novel valuable anti-virulence agents because of their ability in counteracting virulence factor (biofilm formation) without interfering with the bacterial growth in the free life form makes them [73].



59a, R = H, R₁ = H, R₂ = 2,5-OCH₃;
 59b, R = H, R₁ = CH₃, R₂ = 3-OCH₃;
 59c, R = H, R₁ = Cl, R₂ = 2,5-OCH₃;


Novel [1,2,4]triazolo[3,4-b][1,3,4]thiadiazole derivatives from aryl acetic acids and tested all the compounds for antimicrobial activity using ciprofloxacin and fluconazole as reference drugs. Compounds **60a** and **60b** exhibited highest activity against *S. aureus* and were more potent than standard; **60c** and **60d** showed remarkable activity against *M. luteus*. **60d** displayed strong antibacterial activity than the reference drug against *E. coli*. The compound **60e** showed significant activity against *P. aeruginosa*. Among fungal strains, compounds **60c** and **60f** displayed potent activity against *A. niger*, and **60d** exhibited moderate potency against *C. albicans* [74].



60a-f

60a, R₁ = C₆H₅, R₂ = n-C₉H₁₉;
 60b, R₁ = 2,4-Cl₂C₆H₃, R₂ = n-C₉H₁₉;
 60c, R₁ = C₆H₅, R₂ = 2,4-Cl₂C₆H₃CH₂;
 60d, R₁ = 4-Cl-C₆H₄, R₂ = 2,4-Cl₂C₆H₃CH₂;
 60e, R₁ = 2,4-Cl₂-C₆H₃, R₂ = 2,4-Cl₂C₆H₃CH₂;
 60f, R₁ = 4-CH₃OC₆H₄, R₂ = n-C₉H₁₉;


Some 1,3,4-thiadiazole derivatives derived from azo dye. All the compounds were tested for antimicrobial activity. Compound **61a** displayed antimicrobial activity against *Candida albicans*, *Enterobacter aerogenes* and *Salmonella enteritidis*. Compound **61b** was found active against *Salmonella typhimurium* and *Enterococcus durans*. Compound **61c** exhibited activity against *Staphylococcus epidermidis* and *Candida albicans*. Compound **61d** had antimicrobial activity against *Enterobacter aerogenes* and *Candida albicans* [75].




61a, R = CH₃; 61b, R = H;
 61c, R = NO₂; 61d, R = Cl;

61a-d

The series of 2-amino-5(2,4-hydroxyphenyl)-1,3,4-thiadiazole-derived homologues and examined their ability to form metal complexes with Zn(II) and Cu(II) ions. Authors observed strong synergistic antibacterial effect against *Staphylococcus aureus*, using concomitant treatment of thiadiazole derivatives with the commercial antibiotic kanamycin. Compounds **62** and **63** revealed a promising synergistic interaction with kanamycin resulting in a considerably enhanced activity against *S. aureus*. The MIC value of 0.5 µg/mL calculated for kanamycin coupled with relatively inactive compound **62**, found 8-fold lower compared to that of separately tested kanamycin (3.9 µg/mL) and few orders of magnitude lower compared to that of thiadiazole **62** alone. Interestingly, an identical result was received from the mixture of kanamycin with complex **63**, suggesting that the interactions between kanamycin may occur via moieties which are not involved in the formation of metal complex [76].




62



63

New 1,3,4-thiadiazole-1H-pyrazol-4-ylthiazolidin-4-one derivatives. All the derivatives were subjected for their antibacterial activity. Compounds **64a** and **64b** bearing nitro substituent exhibited fourfold (MBC = 156.3 µg/cm³) and twofold (MBC = 312.5 µg/cm³)

stronger activity respectively against *P. aeruginosa* when compared to the reference drug ciprofloxacin (MBC = 625 µg/cm³) [77].




64a, R = 4-NO₂;
64b, R = 3-NO₂;

64a-b


Synthesis of bis-1,3,4-thiadiazoles derivatives and screened them for antimicrobial activity. Activity data indicated that compound **65** exhibited stronger activity against (*Aspergillus flavus*) compared to standard drug used (Ketoconazol) and exhibited equipotent activity against (*Candida albicans*) when compared with Ketoconazol. Compound **66** displayed moderate

activity against all tested fungi and bacteria except (*Aspergillus flavus*). Molecular docking study revealed strong binding pattern of compound **65** at active site of target enzymes, Aspartic Proteinase (SAP2) from *Candida albicans* and enoyl-ACP reductase enzyme [78].


New dihydro-1,3,4-thiadiazole derivatives. The antimicrobial screening of newly synthesized compounds revealed that compounds **67** and **68** are the most potent against the Gram positive (*S. aureus*) and the Gram-negative (*E. coli*) bacteria compared to ciprofloxacin. The docking study was carried out using the bacterial DNA gyrase structure. Docking results revealed that compound **67**, nicely bound to the substrate binding pocket of 4URO via incorporation of four hydrogen bonds with Arg84, Gly85, Arg144 and Thr173. In addition to five hydrophobic interactions with Glu58, Arg84, Pro87 and Ile102 and this may explain the high antibacterial activity of compound. Compound **68** engaged in two hydrogen bonds, one with Arg84 and Arg144 [79].




65




66



Synthesis and antibacterial activity of 2,6-disubstituted and 2,5,6-trisubstituted imidazo [2,1-b][1,3,4] thiadiazole derivatives was reported. The investigation data indicated that compounds bearing morpholine and piperidine exhibited highest activity. Compound **69** containing the piperidine group exhibited highest activity against all species of bacteria and fungi. Also, it was observed that compounds **70** and **69**, which displayed higher antibacterial, have the highest docking score (-9.5 and 9.3 kcal/mol, respectively) as well [80].



Synthesis of imidazo[2,1-b][1,3,4]thiadiazoles from carbohydrates with D-ribo and D-xylo configuration was reported. The compounds were tested for the antiviral activity against Junin virus (the etiological agent of Argentine hemorrhagic fever) by a virus yield inhibition assay. The study indicated that only the p-chloro derivatives (**71**, **72** and **73**) displayed moderate and selective antiviral activity with EC50 close to 200 micro molar although they were less effective than the reference compound rivabirin with EC50 19.2 [81].



3. ANTITUBERCULAR ACTIVITY

Increasing bacterial resistance to commonly prescribed antitubercular drugs is one of the major issue while treatment of infections caused by *Mycobacterium tuberculosis* strains. Additionally increase in overlap of the AIDS and tuberculosis pandemics coupled with the multidrug-resistant tuberculosis (MDR-TB) worsens the situation further Moreover, coupled


with the increasing have brought tuberculosis among the major worldwide. The development of new classes of antitubercular drugs containing a core of 1,3,4-thiadiazole moiety is a very challenging task to many scientists.

The discovery and structure-activity relationships of 5-substituted-2-[(3,5- dinitrobenzyl)sulfanyl]-1,3,4-thiadiazoles (**74a-l**) derivatives with remarkable in vitro activity against

Mycobacterium tuberculosis CNCTC My 331/88 and six multidrug-resistant clinically isolated strains of *M. tuberculosis*, MIC values as low as 0.03 μM (0.011-0.026 $\mu\text{g/mL}$) have been reported.



5-substituted 2-[(3,5-dinitrobenzyl)sulfanyl]-1,3,4-thiadiazoles (**74k**) exhibited outstanding activity against drug-susceptible and multi drug resistant *M. tuberculosis*, with no cross resistance with first- and second-line anti-TB drugs. Moreover, these compounds exhibited excellent activity against the non-replicating *M. tuberculosis* strain SS18b-Lux. SAR study revealed that 3,5-dinitro substitution plays important role in

antimycobacterial activity: any changes to the positions or numbers of nitro groups led to a major decrease in antimycobacterial activity. The antimycobacterial effects of the investigated compounds were selective, as they showed no growth inhibitory activity against other bacteria or against fungi and had low toxicity against proliferating cell lines and isolated human hepatocytes. Moreover, several genotoxicity and mutagenicity assays indicated that these nitro group-containing compounds have low mutagenicity. These results indicate that the reported compounds affect some specific mycobacterial system [82].




- 74a, R = CH₃; 74b, R = 2-Cl-Ph;
 74c, R = 3-Cl-Ph; 74d, R = 4-Cl-Ph;
 74e, R = 4-CH₃-Ph; 74f, R = 3,5-(CH₃)₂-Ph;
 74g, R = 3-CH₃O-Ph; 74h, R = 4-CH₃O-Ph;
 74i, R = 3-NO₂-Ph; 74j, R = 4-NO₂-Ph;
 74k, R = 3,4-di-NO₂-Ph; 74l, R = 4-Pyridyl;

Synthesis of novel triazole-imidazo[2,1-b][1,3,4]thiadiazole hybrids derivatives have been reported by Ramprasad et al. exhibiting activity against *Mycobacterium tuberculosis* H37Rv strain. Compound **75** and **76** displayed highest activity with a MIC of 3.125 $\mu\text{g/mL}$ which is comparable with reference drug ethambutol. The active molecules exhibited positive drug-likeness score and their Clog *P* values are in the range 2.2–2.9. The active derivatives do not shown any kind of cellular toxicity [83].




Synthesis of Schiff bases by reacting a variety of carbonyl compounds with 5-amino-1,3,4-thiadiazole-2-thiol have been reported. All these compounds were evaluated for their antibacterial, antifungal and antitubercular activities. The compounds **77e** and **77d** substituted with the electron withdrawing fluorine and nitro groups exhibited remarkable inhibitory activity against *Staphylococcus aureus*, *Aspergillus niger* and *Candida tropicalis* with an MIC of 8 $\mu\text{g/mL}$ whereas **77a** containing the electron releasing dimethylamine group displayed strong activity against *Proteus vulgaris*. Moreover compounds **77e**, **77b**, **77c** and **77d** also displayed outstanding antimycobacterial activity than the standard pyrazinamide [84].




- 77a, R₁ = CH₃; R₂ = *p*-C₆H₄-N-(CH₃)₂;
 77b, R₁ = CH₃; R₂ = *p*-C₆H₄-Cl;
 77c, R₁ = CH₃; R₂ = *p*-C₆H₄-Br;
 77d, R₁ = CH₃; R₂ = *p*-C₆H₄-NO₂;
 77e, R₁ = CH₃; R₂ = *p*-C₆H₄-F;

Synthesis of 2,5-disubstituted-1,3,4-thiadiazole derivatives and in vitro antimycobacterial activity against *Mycobacterium smegmatis* MC-155. Compounds **78** and **79** displayed significant


antitubercular activity with MIC value 65.74 and 40.86 respectively. Compound **78** was found to be safe and potent antimycobacterial agent when tested on human normal cells HEK293T [85].



Synthesis of substituted derivatives of imidazo[2,1-b][1,3,4]thiadiazole. All the synthesized derivatives were screened for anti-TB and anti-fungal activity. The compounds **80a**, **80b**, **80c**, **80f**, **80g** and **80h** with MIC 1.6-6.25 μ g/ml exhibited strong antitubercular activity. Compounds **80a**, **80d**, **80e**, and **80h** showed strong antifungal activity with MIC 5 μ g/ml owing to presence of electron withdrawing groups at 4th position to both phenyl rings which are attached to the thiazole of the imidazo thiadiazole and imidazo thiadiazole ring [86].




Imidazo[2,1-b][1,3,4]thiadiazole derivatives (**81a–j**) were tested for antitubercular activity against *M. tuberculosis* strain H37Rv by using the MABA method.. Compounds **81a**, **81b**, **81c**, **81d**, **81f** and **81i** showed excellent antitubercular activities. Compound **81f** containing the nitro phenyl substituent exhibited highest activity with MIC of 3.14 μ g/mL. Author noticed considerable variation in activity with different substituents at the 6th position of imidazo(2,1-b)-1,3,4-thiadiazole nucleus [87].




R = a = 3-NO₂, b = 4-Br, c = 4-Cl, d = 4-F, e = 2-OH, f = 4-NO₂,
g = 4-CH₃, h = 3-OH, i = 2,4-Cl, j = 2,4-OH

Novel triazole–imidazo[2,1-b][1,3,4]thiadiazole hybrids designed by a molecular hybridization approach were synthesized and tested against *Mycobacterium tuberculosis* H37Rv strain. Compounds **82a** and **82b** exhibited excellent growth inhibitory activity against the bacterial strain with a MIC of 3.125 μ g/mL. It was noticed that the presence of chloro substituent on the imidazo[2,1-b][1,3,4]thiadiazole ring and ethyl, benzyl or cyanomethylene groups on the 1,2,3-triazole ring

increase the inhibition activity of the molecules. The active compounds are devoid of any toxicity to a normal cell line makes these compounds safe [88].




Novel substituted 1,3,4-thiadiazole derivatives and tested them for in vitro anti-mycobacterial activity against the *Mycobacterium tuberculosis* H37Rv and resistance MDR-TB strain. Compound N-(5-(4-nitrophenyl)-1,3,4-thiadiazol-2-yl)furan-2-carboxamide (**83c**) exhibited highest activity with MIC of 9.87 μ M against the MDR-TB strain as compared to the standard isoniazid (> 200 μ M). The tested compounds **83a**, **83b**, **83c**, **83d**, **83e**, **83f**, **84a**, **84b**, **84c** and **85** also displayed significant MDR inhibitory activity. Compounds were found safe at non-cytotoxic concentrations when assessed for cyto-toxicity to a mammalian Vero cell line using the MTT assay. SAR study revealed that activity is significantly influenced by various 5 substituents at the 2nd position of 1,3,4-thiadiazole and electron withdrawing group on aliphatic side chain at 2nd position of 1,3,4-thiadiazole has diminishing effect on anti-mycobacterial and MDR inhibitory activity [89].




83a-f

R = a = CH₂-CH₂-CH₂-CH₃, b = CH₂-CH₂-CH₂-CH₂COCl,
c = CH₂-CH₂-CH₃, d = C(Cl)₃, e = 4-NO₂-C₆H₄, f = 4-CH₃-C₆H₄,




84a-c



85


R = a = CH₂-CH₂-CH₂-CH₃, b = -furoyl, c = 4-NO₂-C₆H₄

New cyrhetrenyl and ferrocenyl 1,3,4-thiadiazoles were designed, synthesized, characterized and evaluated against *M. Tuberculosis* using Isoniazid as the reference drug in this study. Taking into account the similar MIC values found for the ferrocenyl (**86a-c**) and cyrhetrenyl (**87a-c**) TZDs (MIC N 100 μ g ml⁻¹), author concluded that the opposite electronic effects of the organometallic fragments are not an important factor in the antitubercular activities of these types of compounds. The MIC values are far higher than those of isoniazid but are comparable [90].



86a, R = -CH₃;
86b, R = -
C₂H₅;
86c, R = -C₆H₅;


86a-c




87a, R = -CH₃;
87b, R = -C₂H₅;
87c, R = -C₆H₅;

87a-c


Pyrrolyl-1,3,4-thiadiazoles derivatives having activity against *M. tuberculosis* H37Rv strain were reported. Compounds **88** and **89** showed potent activities (3.125 and 6.25 lg/mL) against *M. tuberculosis* H37Rv strain. Designed structures have shown interactions with the substrate binding site of InhA, confirming their high inhibitory potency, depending on the type of aryl ring modification. The CoMFA models displayed high correlative and predictive abilities. Compounds **90**, **91**, **92** and **89** having halogen 4-F at the phenyl moiety, displayed higher activity compared to other derivatives in the series. Compounds showed moderate cytotoxicity compared to standard INH [91].




88




89



90




91



92

Novel imidazo[2,1-b][1,3,4]thiadiazole (ITD) derivatives with for antimicrobial properties were reported. Among all the microorganisms *Mycobacterium smegmatis* was found most sensitive microorganism with the MIC value between <0.24 and 0.49 µg/mL concentration compared to reference drug Streptomycin (4 g/mL). It was noticed that Mannich bases **93a**, **93b**, **93c**, **93d**, 1,3-thiazolidin-4-one derivatives **94a**, **94b** and **95** exhibited outstanding antimycobacterium activity and also exhibited activity against bacteria with high inhibition at low concentrations (< 0.24 µg/mL) compared with standard drug Ampicillin. Compound **96** including piperidin-2,6-dione and compound **95** showed potent activity against entire test microorganisms than that of standard drugs [92].

New compound, N-{3-(methylsulfanyl)-1-[5-(phenylamino)-1,3,4-thiadiazole-2-yl]propyl}benzamide (**97**) which found active against *Influenza A H3N2* virus with an EC₅₀ value of 20-40 µM [93].




97

New thiazol-imidazo[2,1-b]-1,3,4-thiadiazole hybrids with antitubercular activity was reported. Compound **98a**, substituted with trifluoromethyl and p-chlorophenyl at 2 and 6 positions of the ITD ring, respectively, exhibited highest activity with an MIC of 6.03 µM comparable with MIC values of standard drugs ethambutol (15.3 µM) and ciprofloxacin (9.4 µM). Compounds **98b**, **98c**, **98d** and **98e** also displayed promising inhibitory activity with MIC values in the range of 11.7–13.9 µM. SAR study revealed that the trifluoromethyl substitution at position-2 and p-chlorophenyl substitution at position-6 of the imidazo[2,1-b][1,3,4]thiadiazole ring enhanced


the inhibitory activity. Also, the methyl, methoxy, fluoro or nitro substituents on the thiazole ring enhanced the activity of the compounds. All the compounds are devoid of general cellular. In silico molecular docking studies revealed the favorable interaction of the potent compounds with the target enzymes InhA and CYP121 [94].

Phenothiazine and 1,3,4-thiadiazole hybrid derivatives were evaluated for their in vitro inhibition activity against *Mycobacterium tuberculosis* H37Rv (MTB). Among the series compounds **99a** and **99b** came up with most potent derivative with MIC of 0.8µg/mL (~1.9

µM). Moreover, compounds **99c**, **99d**, **99e**, **99f**, **99g** and **99h** (MIC = 1.6 µg/mL), and compounds **99i**, **99j** and **99k** (MIC = 3.125 µg/ml) displayed strong inhibition activity. The SAR study revealed that an alkyl (methyl/n-propyl) or substituted (4-methyl/4-Cl/4-F) phenyl groups on the 1,3,4-thiadiazole ring enhance the inhibition activity of the compounds. The cytotoxicity study demonstrated all the compounds lacks in cellular toxicity. The molecular docking study revealed strong pi-pi stacking interaction of the active molecules with the target enzymes InhA and CYP121 [95].




- 98a, R₁ = -CF₃; R₂ = -Cl; R₃ = -CH₃;
 98b, R₁ = -CH₃; R₂ = -CH₃; R₃ = -F;
 98c, R₁ = -CH₃; R₂ = -OCH₃; R₃ = -NO₂;
 98d, R₁ = -CF₃; R₂ = -Cl; R₃ = -OCH₃;
 98e, R₁ = -CF₃; R₂ = -Cl; R₃ = -F;




- 99a, R₁ = H; R₂ = -*p*-CH₃-C₆H₄; 99b, R₁ = Cl; R₂ = --CH₃(CH₂)₂;
 99c, R₁ = H; R₂ = -CH₃; 99d, R₁ = Cl; R₂ = -CH₃;
 99e, R₁ = CF₃; R₂ = -CH₃; 99f, R₁ = Cl; R₂ = -*p*-Cl-C₆H₄;
 99g, R₁ = CF₃; R₂ = -*p*-F-C₆H₄; 99h, R₁ = H; R₂ = --CH₃(CH₂)₂;
 99i, R₁ = Cl; R₂ = -*p*-F-C₆H₄; 99j, R₁ = CF₃; R₂ = -*p*-F-C₆H₄;
 99k, R₁ = CF₃; R₂ = --CH₃(CH₂)₂;

2,5-substituted-1,3,4-thiadiazoles were synthesized and evaluated for antituberculosis activity. Compound 5-phenyl-N-[[4-(trifluoromethyl)-phenyl]methyl]-1,3,4-thiadiazole-2-amine **100** exhibited most potent activity against *mycobacterium smegmatis* MC155 (MIC 26.46 µg/mL) compared to the Isoniazid control (12 µg/mL). The SAR analysis demonstrated that a smaller aromatic ring and electron-withdrawing groups favors activity [96].



100

5-substituted-1,3,4-thiadiazole-based fluoroquinolone derivatives were designed as potential antibacterial agents using a molecular hybridization approach. Compound **101a** was rewarded as the most active agent exhibiting antibacterial activity against *Escherichia coli* and *Staphylococcus aureus* with MIC values of 4µg/mL and 2µg/mL, respectively. Amongst the synthesized fluoroquinolone derivatives, compounds **101b** and **101a** were displayed significant antitubercular activity with 8 µg/mL MIC values for each. Most potent derivative, compound **101a** was docked against *Staphylococcus aureus* and *Mycobacterium tuberculosis* DNA gyrase enzymes and found capable for appropriate binding at target site [97].



101a, R₁ = 2,4-dichlorophenyl; R₂ = Ethyl;
101b, R₁ = 4-chlorophenyl; R₂ = Ethyl

101a-b

4. CONCLUSION

1,3,4-Thiadiazole is a unique pattern associated with various biological activities. The potency of the 4-thiazolidinone core is clearly evident from clinically used drugs such as acetazolamide, metazolamide and megestrol. Although antibacterial, anti-tubercular drugs, carbonic anhydrase inhibitors and antiulcer are the four main areas of clinical use, other potential targets remain to be explored. Most locations have been explored to improve the antibacterial and anti-tubercular profile of 1,3,4-thiadiazole, but none of the derivatives have shown promising anti-tubercular activity. The literature is extensively analyzed to provide a meaningful overview of the structural requirements for the business whenever possible.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Abas M, Bahadur A, Ashraf Z, Iqbal S, Rajoka MS, Rashid SG, Jabeen E, Iqbal Z, Abbas Q, Bais A, Hassan M. Designing novel anticancer sulfonamide based 2, 5-

- disubstituted-1, 3, 4-thiadiazole derivatives as potential carbonic anhydrase inhibitor. *J. Mol. Struct.* 2021;1246(1): 131145.
- Altıntop MD, Ciftci HI, Radwan MO, Sever B, Kaplancıklı ZA, Ali TF, Koga R, Fujita M, Otsuka M, Özdemir A. Design, synthesis, and biological evaluation of novel 1, 3, 4-thiadiazole derivatives as potential antitumor agents against chronic myelogenous leukemia: striking effect of nitrothiazole moiety. *Mol.* 2018;23(1):59.
- Shiri P. An overview on the copper-promoted synthesis of five-membered heterocyclic systems. *Appl. Organomet. Chem.* 2020;34(5):e5600.
- Shiri P. Novel Hybrid Molecules Based on triazole-β-lactam as Potential Biological Agents. *Mini Rev Med Chem* .2021;21(5):536-53.
- Mashayekh K, Shiri P. An overview of recent advances in the applications of click chemistry in the synthesis of bioconjugates with anticancer activities. *ChemistrySelect.* 2019;4(46):13459-78.
- Alizadeh A, Roosta A, Rezaiyehrad R, Halvagar M. Efficient one pot and chemoselective synthesis of functionalized 3-bromo-4, 5-dihydroisoxazole derivatives via 1, 3-dipolar cycloaddition reactions of nitrile oxides. *Tetrahedron.* 2017;73(48):6706-11.
- Alizadeh A, Roosta A, Halvagar M. Four-Component Regio-and Diastereoselective Synthesis of Pyrrolizidines Incorporating Spiro-Oxindole/Indanedione via 1, 3-Dipolar Cycloaddition Reaction of Azomethine Ylides. *ChemistrySelect.* 2019;4(1):71-4.

8. Alizadeh A, Roosta A, Halvagar MR. An efficient one-pot synthesis of highly substituted [1, 8] naphthyridin-1-phenyl-1-ethanone derivatives via a four-component reaction. *J. Iran. Chem. Soc.* 2017;14(10):2157-65.
9. Balaban AT, Oniciu DC, Katritzky AR. Aromaticity as a cornerstone of heterocyclic chemistry, *Chem. Rev.* 2004;104(5):2777-812.
10. Mitnik DG. A theoretical study on the aromaticity of thiadiazoles and related compounds, *J. Mol. Struct. THEOCHEM*, 2001;549(3):285-88.
11. Holla BS, Poorjary KN, Rao BS, Shivananda MK. New bis-aminomercaptotriazoles and bis-triazolothiadiazoles as possible anticancer agents. *Eur J Med Chem.* 2002;37(6):511–17.
12. Yousif E, Majeed A, Al-Sammarrae K, Salih N, Salimon J, Abdullah B. Metal complexes of Schiff base: preparation, characterization and antibacterial activity. *Arabian J Chem.* 2017;5(2):S1639-644.
13. Li Y, Geng J, Liu Y, Yu S, Zhao G. Thiadiazole – a promising structure in medicinal chemistry. *Chem. Med. Chem.* 2013;8(1):27–41.
14. Wermuth CG, Aldous D, Raboisson P, Rognan D. *The practice of medicinal chemistry.* 4th ed. London: Academic Press, Elsevier; 2015.
15. Brown N. *Bioisosteres in Medicinal Chemistry.* Weinheim, Germany: Wiley-VCH Verlag and Co., KGaA; 2012.
16. Tripathy R, Ghose A, Singh J. 1,2,3-Thiadiazole substituted pyrazolones as potent KDR/VEGFR-2 kinase inhibitors. *Bioorg Med Chem Lett.* 2007;17(6):1793-798.
17. Fosbinder RJ, Walter LA. Sulfanilamido derivatives of heterocyclic amines. *J. Am. Chem. Soc.* 1939;61(8):2032-33.
18. Beale JM, Block JH. *Wilson and Gisvold's textbook of organic medicinal and pharmaceutical chemistry.* 12th ed. Philadelphia: Lippincott Williams and Wilkins, Wolters Kluwer; 2011.
19. Kaur IP, Smitha R, Aggarwal D, Kapil M. Acetazolamide: future perspective in topical glaucoma therapeutics, *Int. J. Pharm.* 2002;248(1-2):1-4.
20. P. Wolf, Acute drug administration in epilepsy: A review, *CNS Neurosci. Ther.* 2011;17(5):442-48.
21. Russell MB, Ducros A. Sporadic and familial hemiplegic migraine: pathophysiological mechanisms, clinical characteristics, diagnosis, and management, *Lancet Neurol.* 2011;10(5):457-70.
22. Medindex [homepage on the Internet]. Available: <http://medindex.am/glossary/index.php/term/UMLS.+CSP-HL7-ICD9CM-NCI-NDFRT-RXNORM,METHAZOLAMIDE.xhtml>. Accessed March 7, 2017.
23. Sah P, Bidawat P, Seth M, Gharu CP. Synthesis of formazans from Mannich base of 5-(4-chlorophenyl amino)-2-mercapto-1,3,4-thiadiazole as antimicrobial agents, *Arab. J. Chem.* 2014;7(2):181-87.
24. Sunil D, Isloor AM, Shetty P, Satyamoorthy K, Prasad ASB. 6-[3-(4-Fluorophenyl)-1H-pyrazol-4-yl]-3-[(2-naphthyloxy)methyl][1,2,4]triazolo[3,4-b][1,3,4]thiadiazole as a potent antioxidant and an anticancer agent induces growth inhibition followed by apoptosis in HepG2 cells, *Arab. J. Chem.* 2010;3(4):211-17.
25. Zoumpoulakis P, Camoutsis C, Pairas G, Sokovic M, Glamoclija J, Potamitis C, Pitsas A. Synthesis of novel sulfonamide-1,2,4-triazoles, 1,3,4-thiadiazoles and 1,3,4-oxadiazoles, as potential antibacterial and antifungal agents. *Biological evaluation and conformational analysis studies*, *Bioorg. Med. Chem.* 2012 Feb 15;20(4):1569-83.
26. Kolavi G, Hegde V, Khazi I, Gadad P. Synthesis and evaluation of antitubercular activity of imidazo[2,1-b][1,3,4]thiadiazole derivatives, *Bioorg. Med. Chem.* 2006;14(9):3069-80.
27. Mazzone G, Pignatello R, Mazzone S. Synthesis and local anesthetic activity of alkylaminoacyl derivatives of 2-amino-1,3,4-thiadiazole, *Farmaco.* 1993;48(9):1207-24.
28. Kasimogullari R, Bulbul M, Arslan BS, Gokce B. Synthesis, characterization and antiglaucoma activity of some novel pyrazole derivatives of 5-amino-1,3,4-thiadiazole-2-sulfonamide, *Eur. J. Med. Chem.* 2010;45(11):4769-73.
29. Jatav V, Mishra P, Kashaw S, Stables JP. CNS depressant and anticonvulsant activities of some novel 3-[5-substituted 1,3,4-thiadiazole-2-yl]-2-styryl quinazoline-4(3H)-ones, *Eur. J. Med. Chem.* 2008;43(9):1945-54.
30. Kadi A, Al-Abdullah ES, Shehata IA, Habib EE, Ibrahim TM, El-Emam AA. Synthesis,

- antimicrobial and anti-inflammatory activities of novel 5-(1-adamantyl)-1, 3, 4-thiadiazole derivatives, *Eur. J. Med. Chem.* 2010;45(11):5006-11.
31. Clerici F, Pocar D, Guido M, Loche A, Perlini V, Brufani MJ. Synthesis of 2-amino-5-sulfanyl-1,3,4-thiadiazole derivatives and evaluation of their antidepressant and anxiolytic activity, *J. Med. Chem.* 2001;44(6):931-6.
 32. Hasui T, Matsunaga N, Ora T, Ohyabu N, Nishigaki N, Imura Y, Igata Y, Matsui H, Motoyaji T, Tanaka T, Habuka N, Sogabe S, Ono M, Siedem CS, Tang TP, Gauthier C, De Meese LA, Boyd SA, Fukumoto SJ. Identification of benzoxazin-3-one derivatives as novel, potent, and selective nonsteroidal mineralocorticoid receptor antagonists, *J. Med. Chem.* 2011;54(24):8616-31.
 33. Yu L, Gan X, D. Zhou, F. He, S. Zeng, D. Hu, Synthesis and antiviral activity of novel 1,4-pentadien-3-one derivatives containing a 1,3,4-thiadiazole moiety, *Molecules*, 2017;22(4):658.
 34. Khan I, Ali S, Hameed S, Rama NH, Hussain MT, Wadood A, Uddin R, Ul-Haq Z, Khan A, Ali S, Choudhary MI. Synthesis, antioxidant activities and urease inhibition of some new 1,2,4-triazole and 1,3,4-thiadiazole derivatives, *Eur. J. Med. Chem.* 2010;45(11):5200-7.
 35. Woolhouse MEJ, Sequeria SG. Host range and emerging and reemerging pathogens. *Emerg. Infect. Dis.* 2005;11(12):1842-47.
 36. Dye C. after 2015: Infectious diseases in a new era of health and development. *Philos. Trans. R. Soc. B Biol. Sci.* 2014;369(1645):20130426.
 37. Lozano R, M Naghavi K Foreman S, Lim K, Shibuya V, Aboyans J, Abraham T, Adair R, Aggarwal SY, Ahn, Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: A systematic analysis for the global burden of disease study 2010, *Lancet, The lancet.* 2012;380(9859):2095-128.
 38. Sharma PC, A. Sinhar, A. Sharma, H. Rajak, D.P. Pathak. Medicinal significance of benzothiazole scaffold: an insight view. *J. Enzyme. Inhib. Med. Chem.* 2013;28(2): 240-66.
 39. Johnson DS, Li JJ. The art of drug synthesis. Hoboken, NJ: Wiley-Interscience, John Wiley and Sons, Inc; 2007.
 40. Bhuva H, Sahu D, Shah BN, Modi DC, Patel MB. Biological profile of thiadiazole. *Pharmacology Online.* 2011;1(1):528-43.
 41. Othman AA, M. Kihel, S. Amara, 1,3,4-Oxadiazole, 1,3,4-thiadiazole and 1,2,4-triazole derivatives as potential antibacterial agents. *Arabian J. Chem.* 2019;12(7):1660-75..
 42. Sallam MS, El-Hashash M, Guirguis DB. Synthesis and antimicrobial activity of some novel substituted pyridazin-3(2H)-ones containing 1,3,4-thiadiazole moiety, *Med. Chem. Res.* 2016; 25(2):369-80.
 43. Rezki N, A.M. Al-Yahyawi, S.K. Bardaweel, F.F. Al-Blewi, M.R. Aouad, Synthesis of novel 2,5-disubstituted-1,3,4-thiadiazoles clubbed 1,2,4-triazole, 1,3,4-thiadiazole, 1,3,4-oxadiazole and/or schiff base as potential antimicrobial and antiproliferative agents, *Molecules* 2015 Sep;20(9):16048-67.
 44. Noolvi MN, Patel HM, Kamboj S, Cameotra SS. Synthesis and antimicrobial evaluation of novel 1,3,4-thiadiazole derivatives of 2-(4-formyl-2-methoxyphenoxy) acetic acid, *Arab. J. of Chem.* 2016;9(1):S1283-89.
 45. Bhardwaj V, Noolvi MN, Jalhan S, Patel HM. Synthesis, and antimicrobial evaluation of new pyridine imidazo [2,1b]-1,3,4-thiadiazole derivatives, *J. of Saudi Chem. Soci.* 2016;20(1):S406-10.
 46. Barot KP, Manna KS, Ghate MD. Design, synthesis and antimicrobial activities of some novel 1,3,4-thiadiazole, 1,2,4-triazole-5-thione and 1,3-thiazolan-4-one derivatives of benzimidazole, *J. of Saudi Chem. Soci* 2017;21(1):S35-43.
 47. El-Gazzar YI, Georgey HH, El-Messery SM, Ewida HA, Hassan GS, Raafat MM, M.A. Ewida, H.I. El-Subbagh, Synthesis, biological evaluation and molecular modeling study of new (1,2,4-triazole or 1,3,4-thiadiazole)-methylthio-derivatives of quinazolin-4(3H)-one as DHFR inhibitors, *Bioorg. Chem.* 2017;72(1):282-92.
 48. Muglua H, Yakanb H, Shouaib HA. New 1,3,4-thiadiazoles based on thiophene-2-carboxylic acid: Synthesis, characterization, and antimicrobial activities, *J. Mol. Struct.* 2020;1203(0): 127470..
 49. Gopi C, Sastry VG, Dhanaraju MD. Synthesis, spectroscopic characterization, X-ray crystallography, structural activity relationship and antimicrobial activity of some novel 4-(5-(10-(3-N, N-dimethylamino)propyl)-10H-phenothiazine-3-yl)-1, 3, 4-thiadiazole-2-yl)

- Azo dye/Schiff base derivatives, *Future J. Pharm. Sci.*, 2017;3(2):79-89.
50. Abo-Bakra AM, Hashemb HE. New 1,3,4-thiadiazole derivatives: Synthesis, characterization, and antimicrobial activity, *J. Heterocyclic. Chem.* 2019;56(3):1038-47.
 51. Vudhgiri S, Koude D, Veeragoni DK, Misra S, Prasad RBN, Jala RCR. Synthesis and Biological Evaluation of 5-Fatty-acylamido-1, 3, 4-Thiadiazole-2-Thioglycosides, *Bioorg. Med. Chem. Lett* 2017;27(15):3370-73.
 52. El-Naggar M, H.A. Sallam, S.S. Shaban, S.S. A. Wahab, A.E.E. Amr , M.E. Azab, E.S. Nossier, M.A. Al-Omar, Design, synthesis, and molecular docking study of novel heterocycles incorporating 1,3,4-thiadiazole moiety as potential antimicrobial and anticancer Agents, *Molecule* 2019;24(6):1066.
 53. Zhong X, X. Wang, L. Chen, X. Ruan, Q. Li, J. Zhang, Z. Chen, W. Xue, Synthesis and biological activity of myricetin derivatives containing 1,3,4-thiadiazole scaffold, *Chem. Cent. J.* 2017;11(1):1-9.
 54. Sahu JK, S. Ganguly, M. Yasir, Synthesis, SAR and molecular docking Studies of certain new derivatives of 1,2,4-triazolo [3,4- b][1,3,4] thiadiazole as potent antimicrobial agents, *Antiinfect Agents*, 2018;16(1):40-48.
 55. Muglu H, N. Sener, H.A.M. Emsaed, S. Ozkinal, O.E. Ozkan, M. Gur, *J. Mol. Struct.* 2018;1174(1):151-59.
 56. El-Badry YAM, M.S. Sallam, M.AA. El-Hashash, Efficient 1,3,4-thiadiazole-4,5-dihydropyridazin-3(2H)-ones as antimicrobial Agents, *Chem. Pharm. Bull.* 2018;66(4):427-33.
 57. Sekhar MM, Nagarjuna U, Padmavathi V, Padmaja A, Reddy NV, Vijaya T. Synthesis and antimicrobial activity of pyrimidinyl 1,3,4-oxadiazoles, 1,3,4-thiadiazoles and 1,2,4-triazoles, *Eur. J. Med. Chem*, 2018;145(1):1-10.
 58. Mutchu BR, Kotra V, Onteddu SR, Boddapati SNM, Bollikolla HB. Synthesis, cytotoxicity and antimicrobial evaluation of some new 2-aryl,5-substituted 1,3,4-oxadiazoles and 1,3,4-thiadiazoles, *Chemistry Africa* 2019;2(1):15-20.
 59. Dai H, G. Li, J. Chen, Y. Shi, S. Ge, C. Fan, H. He, Synthesis and biological activities of novel 1,3,4-thiadiazole-containing pyrazole oxime derivatives, *Bioorg. Med. Chem. Lett.* 2016;26(15):3818-21.
 60. Abu-Hashem AA, R.A.M. Faty, Synthesis, antimicrobial evaluation of some new 1, 3, 4-thiadiazoles and 1, 3, 4-thiadiazines, *Curr. Org. Synth.*, 2018;15(8):1161-70.
 61. Wu Q., H. Cai, T. Yuan, S. Li, X. Gan, B. Song, Novel vanillin derivatives containing a 1,3,4-thiadiazole moiety as potential antibacterial agents, *Bioorg. Med. Chem. Lett.* 2020;30(10):127113.
 62. Muglu H, N. Sener, H.A.M. Emsaed, S. Ozkinal, O.E. Ozkan, M. Gur, Synthesis and characterization of 1,3,4-thiadiazole compounds derived from 4-phenoxybutyric acid for antimicrobial activities, *J. Mol. Struct.* 2018;1174(1):151-59.
 63. Freitas HCN, Barbosab P, Bernardinob, V, Sueth- ntiagoc MSV, Wardelld JL, Wardelle DD. Ricardof TG, Melog EF, da Silvahi K. Salomaob CAM. Fraga, Synthesis and trypanocidal activity of novel pyridinyl-1,3,4-thiadiazole derivatives, *Biomed. Pharmacother.* 2020;127(1): 110162.
 64. Gur M, Sener N, Kastan CA, Ozkan OE, Mugu H, Elmaswaria MAM. Synthesis and characterization of some new heteroaromatic compounds having chirality adjacent to a 1,3,4-thiadiazole moiety and their antimicrobial activities, *J. Heterocycl. Chem.*, 2017;54(6):3578-90.
 65. Abdelhamid OA, Gomha SM, Shawali AS. Utility of N-aryl 2-aryloxyhydrazonopropanehydrazonoyl chlorides as precursors for synthesis of new functionalized 1,3,4-thiadiazoles with potential antimicrobial activity, *J. Adv. Res.* 2015;6(6):885-93.
 66. Lv X, Yang L, Fan Z, Bao X. Synthesis and antimicrobial activities of novel quinazolin-4(3H)-one derivatives containing a 1,2,4-triazolo[3,4-b][1,3,4]thiadiazole moiety, *J. of Saudi Chem. Soci.* 2018;22(1):101-9.
 67. Tahtaci H, Karacık H, Ece A, Er M, Seker MG. Design, synthesis, SAR and molecular modeling studies of novel imidazo[2,1-b][1,3,4] thiadiazole derivatives as highly potent antimicrobial agents, *Mol. Inf.* 2018;37(3):1700083.
 68. Nassar IF, S.R. Att-Allah, M.M. Hemdan, Utility of thiophene-2-carbonyl isothiocyanate as a precursor for the synthesis of 1,2,4-triazole, 1,3,4-oxadiazole and 1,3,4-thiadiazole derivatives with evaluation of their antitumor and antimicrobial activities,

- Phosphorus, Sulfur, Silicon Relat. Elem., 2018;193(10):630-6.
69. Karaburun AC, U.A. Cevik, D. Osmaniye, B.N. Saglık, B.K. Cavusoglu, S. Levent, Y.Ozkay, A.S. Koparal, M. Behcet, Z.A. Kaplancikli, Synthesis and evaluation of new 1,3,4-thiadiazole derivatives as potent antifungal agents, *Molecules* 2018;23(12):3266.
 70. Er M, A. Ozer, S. Direkel, T. Karakurt, H. Tahtaci, Novel substituted benzothiazole and Imidazo[2,1-b][1,3,4]Thiadiazole derivatives: Synthesis, characterization, molecular docking study, and investigation of their *in vitro* antileishmanial and antibacterial activities, *J. Mol. Struct.* 2019;1194(1):284-96.
 71. Tabatabaei Z.M., P. Foroumadi, M. Toolabic, F. Golib, S. Moghimi, S.K. Ardestani, A. Foroumadi, 2-(Bipiperidin-1-yl)-5-(nitroaryl)-1,3,4-thiadiazoles: synthesis, evaluation of *in vitro* leishmanicidal activity, and mechanism of action, *Bioorg. Med. Chem.* 2019;27(16):3682-91.
 72. Farhan ME, M.G. Assy, Heterocyclization of isoniazid: Synthesis and antimicrobial activity of some new pyrimidine, 1, 3-thiazole, 1, 2,4-thiadiazole, and 1, 2, 4-triazole derivatives derived from isoniazid, *Egypt. J. Chem.* 2019;62(2):171-80.
 73. Cascioferro T, B. Parrino, G.L. Petri, M.G. Cusimano, D. Schillaci, V.D. Sarno, S. Musella, E. Giovannetti, G. Cirrincione, P. Diana, 2,6-Disubstituted imidazo[2,1-b][1,3,4]thiadiazole derivatives as potent staphylococcal biofilm inhibitors, *Eur. J. Med. Chem.*, 2019;167:200-10.
 74. Kamboj VK, A. Kapoor, S. Jaina, Synthesis, antimicrobial, and antioxidant screening of aryl acetic acid incorporated 1,2,4-triazolo-1,3,4-thiadiazole derivatives, *J. Heterocycl. Chem.*, 2019;56(4):1376-82.
 75. Gur M. Synthesis, characterization, and antimicrobial properties of new 1,3,4-thiadiazoles derived from azo dyes, *J. Heterocyclic Chem.*, 2019;56(3):980-87.
 76. Karcz D, Matwijczuk A, Kaminski D, Creaven B, Ciszkowicz E, Szlachta KL, Starzak K. Structural features of 1,3,4-thiadiazole-derived ligands and their Zn(II) and Cu(II) complexes which demonstrate synergistic antibacterial effects with kanamycin, *Int. J. Mol. Sci.* 2020;21(16): 5735.
 77. Kerru N, L. Gummidi, S. Sandeep, V.H. Bhaskaruni, S.N. Maddila, S.B. Jonnalagadda, Ultrasound-assisted synthesis and antibacterial activity of novel 1,3,4-thiadiazole-1H-pyrazol-4-yl-thiazolidin-4-one derivatives, *Monatshefte fur Chemie* 2020;151(1):981-90.
 78. Mahmoud HK, A.A. Abbas, S.M. Gomha, Synthesis, antimicrobial evaluation and molecular docking of new functionalized bis(1,3,4-thiadiazole) and bis(thiazole) derivatives, *Polycycl. Aromat. Compd.* 2020;41(9):2029-41.
 79. Moustaf AH, D.H. Ahmed, M.T.M. El-Wassimy, M.F.A. Mohamed, Synthesis, antimicrobial studies, and molecular docking of some new dihydro-1,3,4-thiadiazole and pyrazole derivatives derived from dithiocarbazates, *Synth. Commun.*, 2021;51(4):570-84.
 80. Dagli M, M. Er, T. Karakurt, A. Onaran, H. Alici, H. Tahtaci, Synthesis, characterization, antimicrobial evaluation, and computational investigation of substituted imidazo[2,1-b][1,3,4]thiadiazole derivatives, *Chemistry Select*, 2020;38(5):11753-63.
 81. Fascio ML, CS Sepulveda, EB Damonte, N.B. D'Accorso, Synthesis and antiviral activity of some imidazo[1,2-b][1,3,4]thiadiazole carbohydrate derivatives, *Carbohydr Res.* 2019;480(1):61-66.
 82. Karabanovich G, Zemanova T, Smutny R, Szekeley M, Sarkan I, Centarova A, Vocat I, Pavkova P onka, J. Nmeek, J. Stolaikova, M.Vejsova, K.Vavrova, V. Klimesova, A. Hrabalek, P. Pavek, ST, Cole K, Mikusova J, Rohv, Development of 3,5-dinitrobenzylsulfanyl-1,3,4-oxadiazoles and thiadiazoles as selective antitubercular agents active against replicating and nonreplicating mycobacterium tuberculosis, *J. Med. Chem.*, 2016;59(6):2362-80.
 83. Ramprasad J, Nayak N, Dalimba UP, Yogeewari D, Sriram. One-pot synthesis of new triazole—Imidazo[2,1-b][1,3,4]thiadiazolehybrids via click chemistry and evaluation of their antitubercular activity, *Bioorg. Med. Chem. Lett.* 2015;25(19):4169-73.
 84. Babu KA, Singhvi I, Ravindra N, Shaik AB. Antimicrobial and antitubercular evaluation of some new 5-amino-1,3,4-thiadiazole-2-thiol derived schiff bases, *Rev. Roum. Chim.*, 2020;65(9):771-76
 85. Polkam N, Rayam P, Anireddy JS, Yennam S, Anantaraju HS, Dharmarajan S, Perumal Y, Kotapalli SS, Ummanni R, Balasubramanian S. Synthesis, *in vitro*

- anticancer and antimycobacterial evaluation of new 5-(2,5-dimethoxyphenyl)-1,3,4-thiadiazole-2-amino derivatives, *Bioorg. Med. Chem. Lett.* 2015;25(7):1398-402.
86. Syed MA, Reddy YRP, Chandrasekhar KB. Design, one-pot synthesis and biological evaluation of imidazo[2,1-b][1,3,4]thiadiazole derivatives for their anti-tubercular and anti-fungal activity, *J. App. Pharm. Sci.* 2018;8(07):021-27.
 87. Patel HM, Noolvi MN, Sethi NS, Gadad AK, Cameotra SS. Synthesis and antitubercular evaluation of imidazo[2,1-b][1,3,4]thiadiazole derivatives, *Arab. J. Chem.* 2017;10(1):S996-02.
 88. Ramprasad J, Nayak N, Dalimba U, Yogeewari P, Sriram D. One-pot synthesis of new triazole—Imidazo[2,1-b][1,3,4]thiadiazole hybrids via click chemistry and evaluation of their antitubercular activity, *Bioorg. Med. Chem. Lett.* 2015;25(19):4169-73.
 89. Patel H, Jadhav H, Ansari I, Pawara R, Surana S. Pyridine and nitro-phenyl linked 1,3,4-thiadiazoles as MDR-TB inhibitors, *Eur. J. Med. Chem.* 2019;167(1):1-9.
 90. Quintana C, Klahn AH, Artigas V, Fuentealba M, Biot C, Halloum I, Kremer L, Arancibia R. Cyrhetyrenyl and ferrocenyl 1,3,4-thiadiazole derivatives: Synthesis, characterization, crystal structures and in vitro antitubercular activity, *Inorg. Chem. Commun.* 2015;55(1):48–50.
 91. Joshi SD, More UA, Koli D, Kulkarni MS, Nadagouda MN, Aminabhavi TM. Synthesis, evaluation and in silico molecular modeling of pyrrolyl-1,3,4-thiadiazole inhibitors of InhA, 2015;59(1): 151-67.
 92. E. Taflana, H. Bayrakb, M. Erb, S.A. Karaoglu, A. Bozdeveci, Novel imidazo[2,1-b][1,3,4]thiadiazole (ITD) hybrid compounds: Design, synthesis, efficient antibacterial activity and antioxidant effects, *Bioorg. Chem.* 2019;89(1):102998.
 93. Tatar E, Kucukguzel SG, Sevgi Karakus ED, Clercq G, Andrei R, Snoeck, C, Pannecouque, S, Oktem Okullu N, Unubol, Tanil Kocagoz S, Kalayci F, Sahin I, Kucukguzel. Synthesis and biological evaluation of some new 1,3,4-thiadiazole and 1,2,4-triazole derivatives from L-methionine as antituberculosis and antiviral agents, *Marmara Pharm J.* 2015;19(2):88-102.
 94. Ramprasad J, Nayak N, Dalimba U, Yogeewari P, Sriram D. Ionic liquid-promoted one-pot synthesis of thiazole–imidazo[2,1-b][1,3,4]thiadiazole hybrids and their antitubercular activity, *Med. Chem. Commun.*, 2016;7(2):338-44.
 95. Jurupula Ramprasad, Nagabhushana Nayaka, Udayakumar Dalimba, Design of new phenothiazine-thiadiazole hybrids via molecular hybridization approach for the development of potent antitubercular agents, *Eur. J. Med. Chem.* 2015;106(1): 75-84.
 96. Sekhara DC, Rao DVV, Rao AT, Kumar UL, Jha A. Design and synthesis of 1,3,4-thiadiazole derivatives as novel anticancer and antitubercular agents, *Russ. J. Gen. Chem.*, 2020;63(11):4387-408.
 97. Demirci A, Karayel KG, Tatar E, Okullu SO, Unubol N, Tasli PN, Kocagoz ZT, Sahin F, Kucukguzel I. Synthesis and evaluation of novel 1,3,4-thiadiazole–fluoroquinolone hybrids as antibacterial, antituberculosis, and anticancer agents, *Turk. J. Chem.* 2018;42(3):839-58.