

## Phytochemical and biological activities of the genus *Terminalia*, emphasizing on

### *Terminalia paniculate* L.: a review

#### Abstract

*Terminalia* is the **second** major genus belonging of family Combretaceae. Species **from this genus** have been used in outmoded popular medicine globally. But then, a little study was stated on *Terminalia paniculate* L. (*TP*). This is a complete literature review of various *Terminalia* plants highlighting on *TP*, about its biological and isolated constituents. The objective of this study is to fascinate the attention of unfamiliar activities of *TP*, thus conducive to the growth of novel therapeutics which might benefit health of individuals suffering from illnesses. Existing data on *Terminalia* **was sourced from electronic databases** like Scopus, Goggle Scholar, ScienceDirect, PubMed, Chemical Abstract earth and Springer Link. Plant based studies have led to extraction of various categories of phytochemicals like, tannins, flavonoids, phenolic acids, triterpenes, triterpenoid glycosides, lignan and ligan derivatives in *Terminalia* plants. Extracts and their isolated constituents of various *Terminalia* plants reported a good spectrum of activities. *TP* consist of polyphenols such as ellagic acid, 3,3'-di-*O*-methyyl ellagic acid and 3,4,3'-*O*-trimethyl flavellagic acid. Studies on *Terminalia* genus has opened a diverse active chemical component. Many biochemical potentials have authenticated use of *Terminalia* in therapy of different ailments in plant-based medicine. Future studies desired to discover bioactive constituents' accountable **pharmacological mechanisms**. *Terminalia paniculata* is a tree native to southwest India (including the Western Ghats and Karnataka). Known in the timber trade as kindal. Though it has been used in the ancient system of medicine for **menstrual problems, cough, bronchitis, cardiac debility, hepatitis, diabetes and in obesity**, very little literature support is available on these biological activities, except for its few like acute and chronic anti-inflammatory activity. This review is an attempt to emphasize on the phytochemical and biological activities of terminalia genus and the species *Terminalia paniculate*.

**Keywords:** *TP*, ellagic acid, hepato-protective, nephron-protective.

## 1. INTRODUCTION:

*Terminalia paniculate* L. (Combretaceae) is a tropical tree with huge spreading in west and south India. In olden days, its extract was in the treatment of cholera, inflammation of parotid-glands, menstrual problems, cough, bronchitis, cardiac debility, hepatitis, diabetes and in obesity [1-6]. Juice of *TP* flowers were used as medicines for cholera, inflamed parotid glands and menstrual disorders in folk medicine [2,3]. The bark has 14 % tannins with a pyrogallol nucleus with gallic acid. Heartwood has ellagic acid, dimethyl ellagic acid, penta methyl flavellagic acid, trimethyl flavellagic acid and  $\beta$  sitosterol [7].

The *Terminalia* genus is one of the largest in family Combretaceae, with around 200 plant-species. These species are spread across tropical parts of the world, especially diverse in south-east Asia [1]. *Terminalia* gets its name after a latin word “*terminus*”, as leaves look at the tips of shoot. Mostly the trees are huge, in a height up to 75m<sup>3</sup>. Plants of this genus were extensively used in folk medicine in several countries for the therapy of various diseases like abdominal problems, bacterial attacks, common-cold, sore throats, conjunctivitis, diarrhea, dysentery, fever, gastric ulcers, headaches, heart diseases, hookworm, hypertension, jaundice, leprosy, nosebleed, edema, pneumonia and skin diseases.

In the previous reviews on *Terminalia* by Cock I. E. ,2015 and Fahmy NM *et al.*, 2015 phytochemical and biological activities of the genus were elaborated. This review also focuses on the proven medicinal properties of *Terminalia paniculate* along with the activities on the other species of the genus *Terminalia*. This review encourages the researchers to work on the biological activities of *Terminalia paniculate* which has been lacking scientific evidences for

all these years and thus directs future areas of research into the medicinal activities of this important species of the genus.

This review has 2 main sections, in first part, a phytochemical study of different components and their prevalence in *Terminalia* was reported, in the second part the biological activities led on various species was emphasized.

## 2. Part I: Phytochemicals

Phytochemical research performed on different *Terminalia* plant species have reported presence of tannins, pentacyclic triterpenes and their glycoside derivatives, flavonoids and other phenolic compounds [8-10].

Literature reveals rich source of tannins and pseudomamma, like gallic acid and its gallate esterase, chebulic and non-chebulic ellagitannins, ellagic acid derivatives and ellagic acid glycosides polyphenolics, which are potent natural antioxidants, have created great interest as budding therapeutics in various ailments. Many studies showed flavonoids and tannins with strong hepatoprotective potential in many animal models [5,6].

## 3. Part II: Biological activities

### 3.1 Hepatoprotective and nephroprotective studies

- *T. muelleri* constitutes polyphenolic-rich fraction having hepato-protective and nephron-protective potential in CCl<sub>4</sub>-generated hepato-toxicity and nephron-toxicities in rodents [11,29].
- Oral route of *T. arjuna* fruit juice prevented liver toxicity and oxidation of cadmium-caused liver toxicity in rodents [12,31].
- Manna reported protective actions of arjunolic acid, separated from bark of *T. arjuna*, against sodium arsenate-generated oxidation in mice hepatic cells [13,32].

- *In vitro* studies of hepatic cell lines with chebulic acid and neochebulic acid, extracted from *T. chebula* fruit ethanolic-extract, potentially reduced tert-butyl hydroperoxide-caused cytotoxicity, suppressed active oxygen radicals, and increased the liver GSH [14, 33].
- Corilagin, from *T. catappa* acted against galactosamine and lipopolysaccharide-induced hepato-toxicity in rodents at 1 mg/kg by suppressing oxidation and cell death [15, 34].
- Prior treatment of *T. bellerica* leaves extracts in CCl<sub>4</sub>-generated hepato-toxicity and nephron-toxicity, reported dose-dependent biochemical activity, while isolated gallic acid had strong hepatoprotective effects at 200 mg/kg [16, 35].

### 3.2 Anti-inflammatory studies

- Ethanolic stem extract of *T. phanerophlebia* and the isolated  $\beta$ -sitosterol from ethanolic stem extract prevented cyclooxygenase COX-II selectively [17, 36].
- *T. ferdinandiana* fruit extract reported a potent anti-inflammatory action in lipopolysaccharide-induced murine macrophages, by preventing COX-II and iNOS, also by preventing the synthesis of prostaglandin E<sub>2</sub> [18, 38].
- Chebulagic acid isolated from seeds of *T. chebula*, potentially delayed the inception and development of collagen-induced arthritis in rodents [19, 39].
- Anolignan B from the ethylacetate extract of *T. sericea* root had a non-selective preventive action on both the COX isoenzymes [20-25, 40].
- Punicalagin at 10 mg/kg and punicalin at 5 mg/kg from *T. catappa* leaves reported anti-inflammation against carragenan-induced paw edema in rodents [25,27, 41].
- Ursolic acid and 2 $\alpha$ ,3 $\beta$ ,2 $\epsilon$ 3-trihydroxyurs-12-en-28-oic acid from *T. catappa* ethanol extract of leaves reported anti-inflammatory potential, with a strong reduction (more than 50%) of edema on the mice ear pinna at 0.30 mg/ear [28, 29, 42].

### 3.3 Gastroprotective studies

- Chebulinic acid from fruit extract of *T. chebula* reported a gastroprotective action against duodenal ulcers caused by cold restraining (protection by 62.9%), aspirin- induced (protection by 55.3%), alcohol-induced (protection by 80.6%) and pyloric ligation-induced (protection by 66.6%) models. Chebulinic acid strongly decreases free acid levels by 49%, total acid levels by 38% and enhanced mucus production by 60%. Also, Chebulinic acid potentially suppressed H<sup>+</sup> K<sup>+</sup>-Pump invitro with an IC<sub>50</sub> of 65 µg/ml whereas, Omeprazole reported an IC<sub>50</sub> of 30.24 µg/ml, showing its potent antisecretory power [30, 43].
- Methanolic *T. arjuna* extract resulted in a potent decrease in lesion-index in diclofenac-induced ulceration, and a strong raise in pH, non-protein **sulfhydryl**, decreased glutathione, protein-bound carbohydrate complexes, mucus content with a good decrease in volume of gastric acid, free and total acid levels, pepsin levels, acid secretion, activity of lipid peroxidases and activity of myeloperoxidases [35, 44].
- Ethanolic *T. pallida* extract reported a potent anti-ulcer action against indomethacin, histamine and ethanol induced ulceration in rodents by increasing the antioxidant levels in mucosa, thus enhancing mucosal protection [36-45].

### 3.4 Antimicrobial and Antiviral studies

- *Terminalia* plant species were observed to possess a significant anti-microbial action on wide range of microbes. Water extract of *T. chebula* reported a strong antibacterial action on *H. pylori* with a MIC of 125 µg/ml and MBC of 150 µg/ml [46].
- Acetone *T. chebula* extract reported a significant antibacterial action on *Enterococcus faecalis*, *Bacillus subtilis* and *Klebsiella pneumoniae* [ 47].
- Casuarinin from *T. arjuna* bark extract, reported a strong antiviral action on *Herpes simplex* type II at 25 µM and prevented viral titers up to 1lakh-folds by preventing the attachment and

penetration of virus [48].

- Fyhrquist reported that methanolic extracts of roots and bark of *T. sambsiaca* possessed lesser MIC values than aqueous, butanolic and chloroform extracts against mycobacterium [49].
- The significant antibacterial action of ethyl acetate leaves extract of *T. muelleri* was credited to gallic acid [50].
- Antifungal action of 6 *Terminalia* leaf extracts (*T. proteinoids*, *T. brachystemma*, *T. sericea*, *T. gazensis*, *T. mollis* and *T. sambsiaca*) were evaluated on various fungi. It was reported that extract of acetone had highest antifungal action. Extract of *T. sericea* was most active against wide spectrum of microorganisms [51].
- Anolignan B from ethyl acetate extracts of *T. sericea* roots had a potent action against microbes with MIC 3.80 µg/ml with *Bacillus subtilis* and 31 µg/ml with *Escherichia coli* [40].
- Gallic acid from *T. nigrovenulosa* bark extracts reported a significant antifungal action against *Fusarium solani* [52].
- *T. macroptera* ethanolic extract of roots had promising action against microbes, with the lowest MIC against *Shigella dysenteriae*, *Staphylococcus aureus* and *Vibrio cholera* with a strong action against *Campylobacter* [53]. Additionally, *T. macroptera* root ethanolic extract reported an antimicrobial action against *Neisseria gonorrhoeae* with a MIC range of 100 µg/ml to 200 µg/ml, diethyl ether extract had most active MIC value ranging from 25 µg/ml to 50 µg/ml [54]. Also, it was presumed that punicalagin and terchebulin, were the constituent responsible for activity against *Helicobacter pylori* [55-58].

### 3.5 Cytotoxicity studies

- Methanolic extract of *T. chebula* fruits reported a reduction in cell-viability, anti-cell proliferatory effects, and resulting in apoptosis as dose increased on cancer lines. Also, it

resulted in cell death at reduced doses, and resulted in necrosis at increased doses. Chebulinic acid, tannic acid and ellagic acid, have IC<sub>50</sub> of 53 µg/ml, 59 µg/ml and 78 µg/ml respectively, posed the highest cytotoxicity amongst the fruits of *T. chebula* [59]. Moreover, chebulagic acid from the fruits of *T. chebula* extract reported an antiproliferative effect against HCT-15, COLO-205, MDA-MB-231, DU-145 and K562 cell-lines [60].

- Water extract of *T. catappa* leaves and its constituents *punicalagin* were active against bleomycin- caused genotoxicity in Chinese hamster ovary cells. In addition, *T. catappa* leaves extract resulted in a inhibitory effect on invasion and motility of metastatic A549 and Lewis lung carcinoma cells. The leaf ethanolic extract of *T. catappa* strongly prevented migration of squamous cellcarcinoma cells [61-63].
- Luteolin, gallic acid and gallic acid ethyl ester from methanolic extract stems and leaf of *T. arjuna* had a significant anti-neoplastic potential.
- Ivorenoside C from barks of *T. ivorensis* possessed an anti-proliferative action on MDA-MB-231 and HCT116 human cancer lines with IC<sub>50</sub> of 3.96 µM and 3.43 µM respectively.
- Leaves acetoneextract of *T. calamansanai* prevented viability of HL-60 cells.

### 3.6 Cardioprotective activities

- *T. arjuna* bark had been in use as a folk medicine as cardiotonic. The ethanolic bark extract of *T. arjuna* increased cardiac intracellular antioxidant status in CCl<sub>4</sub>-generated oxidation in rodents [57]. The antioxidant effect was similar to that of Vit C. Additionally, butanol fraction of *T. arjuna* bark extract possessed a protective effect on doxorubicin-generated cardiotoxicity by enhancing antioxidant enzymes, reducing serum creatine kinase levels and also by decreasing lipid peroxidation [61].
- Scientists reported that patients with refractory chronic congestive heart failure, on treatment

with *T. arjuna* bark extract as an additive medicine, resulted in a long-term enhancement in the symptoms of cardiac failure with an enhancement in left ventricular ejection [62]. A clinical trial was conducted to estimate role of *T. arjuna* in ischemic mitral regurgitation (IMR) after myocardial infarction (MI). Patients on treatment with *T. arjuna* as an adjuvant resulted strong prevention in IMR and decrease in frequency of angina.

- Pre-treatment with fruit extract of *T. pallida* upgraded myocardial damage in isoproterenol-induced MI in rodents and resulted in protective actions on cardiac muscle.
- Pre-treatment with *T. chebula* extract upgraded effects of isoproterenol on lipid peroxide formation [51,52].

### 3.7 Anti-hypertensive study

- Bark extract of *T. superba* resulted a strong antihypertensive action in spontaneously hypertensive rats, and also in glucose-induced hypertensive rats due to the withdrawal of sympathetic tone and enhancing antioxidant level [53,54].

### 3.8 Antiparasitic and molluscicidal studies

- *Invitro* nematicidal action of *T. nigrovenulosa* bark on *Meloidogyne incognita* was due to 3,4-dihydroxybenzoic acid [55].
- Ethyl acetate, acetone and methanol leaves and seeds extract of *T. chebula* resulted *invitro* ovicidal and larvicidal action on *Haemonchus contortus* [56]. Additionally, *T. chebula* fruit molluscicidal action was attributed for tannic acid which strongly prevented AChE, ACP and ALP action in the nervous system of *Lymnaea acuminata* [57].
- Ethanolic leaves extract of *T. catappa* had a molluscicidal action on snail intermediate hosts of schistosomiasis (*Biomphalaria pfeifferi* and *Bulinus globosus*) with *B. pfeifferi* being highly vulnerable.

### 3.9 Wound healing activity

- Topical application of *T. chebula* alcoholic leaves extract on rodents' wounds resulted a positive healing effect, by enhancing tensile strength of cells by around 41% and reducing epithelialization [59]. Additionally, tannin-rich fraction from *T. chebula* fruits were recognized for wound healing due to strong anti-bacterial and angiogenic activities [60].
- Topical administration of hydro-alcoholic extract of *T. arjuna* showed in a strong raise in tensile strength of wounds and epithelialization. This wound healing action was marked in tannin-rich portion compared to others.

### 3.10 Biological activities stated on *T. paniculata*

- Water extract of bark of *T. paniculata* decreased high blood glucose, HbA1c, creatinine, urea, ALT, AST levels and upturned the abnormal endogenic antioxidants and lipid levels to normal in Steptazocin-generated diabetic rats in contrast with untreated diabetic rats.
- Ethanolic extract of *T. paniculate* bark reported hepato-protective action and decreased abnormal serum parameters and lipid peroxides in paracetamol- generated liver toxicity in rodents [30].
- Water extract of *T. paniculata* bark strongly decreased carrageenan-caused paw edema volume [37].

## 4 Conclusion

A wide-ranging literature on *Terminalia* genus has discovered a diversity of constituents formed by this genus. **Tannins, flavonoids, phenolic acids, triterpenes, triterpenoid glycosides, lignan and lignan derivatives were found as the chief classes of secondary metabolites of this genus, responsible for the biological activities.** Additionally, present report showed that many biological studies conducted on various extracts and isolated contents from various species of this genus were engrossed on the evaluation of antimicrobial, antioxidant, hepatoprotective, anti-inflammatory, hypoglycemic, hypolipi

dimic, cytotoxic and wound healing effect. The different biological studies authenticated folk medicinal use of various *Terminalia* species.

Though various phytoconstituents and biological findings were reported from *Terminalia*, the researches have attended on a few species, with *T. chebula*, *T. bellerica*, *T. arjuna*, *T. catappa*, *T. horrida*, *T. superba*, *T. macroptera*, *T. pallida*, *T. ivorensis*, *T. sericea* and *T. alata* being greatest phytochemically and biologically evaluated species, parting a lush area for further research on other species like *TP* that have not been fully explored yet.

Acute toxicity study was carried out on *TP* with a dose of 1 000 mg/kg body weight. Under this, the considered parameters such as skin color, hair loss, eating and sleeping patterns and other behavioural observations were not altered showing that *PT* is safe orally at a dose of 1000 mg/kg body weight.

This review delivers a complete understanding of the phytochemicals and biological activities of various *TP* which might benefit in the development of new alternate medicines for the therapy of various illnesses.

## References

1. De Morais Lima GR, de Sales IR, Caldas Filho MR, de Jesus NZ, de Sousa Falcão H, et al. (2012) Bioactivities of the genus Combretum (Combretaceae): a review. *Molecules* 17: 9142-9206.
2. Saxena VG, Mishra KK, Vishwakarma, Saxena A (2013) A comparative study on quantitative estimation of tannins in *Terminalia chebula*, *Terminalia bellerica*, *Terminalia arjuna* and *Saraca indica* using spectrophotometer. *Asian J Pharm Clin Res* 6: 148-149.
3. Stace CA (2007) *Combretaceae*: Springer 9: 67-82.
4. Eloff JN, Katerere DR, McGaw LJ (2008) The biological activity and chemistry of the southern African Combretaceae. *J Ethnopharmacol* 119: 686-699.
5. Yang MH, Vasquez Y, Ali Z, Khan IA, Khan SI (2013) Constituents from *Terminalia* species increase PPAR $\alpha$  and PPAR $\beta$  levels and stimulate glucose uptake without enhancing adipocyte differentiation. *J Ethnopharmacol* 149: 490-498.
6. Bag A, Bhattacharyya SK, Chattopadhyay RR (2013) The development of *Terminalia*

- chebula Retz. (Combretaceae) in clinical research. *Asian Pac J Trop Biomed* 3: 244-252.
7. Pellati F, Bruni R, Righi D, Grandini A, Tognolini M, et al. (2013) Metabolite profiling of polyphenols in a *Terminalia chebula* Retzius ayurvedic decoction and evaluation of its chemopreventive activity. *J Ethnopharmacol* 147: 277-285.
  8. Dixit D, Dixit AK, Lad H, Gupta D, Bhatnagar D (2013) Radioprotective effect of *Terminalia Chebula* Retzius extract against irradiation-induced oxidative stress. *Biomed Aging Pathol* 3: 83-88.
  9. Liu M, Katerere DR, Gray AI, Seidel V (2009) Phytochemical and antifungal studies on *Terminalia mollis* and *Terminalia brachystemma*. *Fitoterapia* 80: 369- 373.
  10. Garcez FR, Garcez WS, Miguel DL, Serea AA, Prado FC (2003) Chemical constituents from *Terminalia glabrescens*. *J Braz Chem Soc* 14: 461-465.
  11. Li DQ, Zhao J, Xie J, Li SP (2014) A novel sample preparation and on-line HPLC–DAD–MS/MS–BCD analysis for rapid screening and characterization of specific enzyme inhibitors in herbal extracts: Case study of  $\alpha$ -glucosidase. *J Pharm Biomed Anal* 88: 130-135.
  12. Gao H, Huang YN, Xu, Kawabata J (2007) Inhibitory effect on  $\alpha$ -glucosidase by the fruits of *Terminalia chebula* Retz. *Food Chem* 105: 628-634.
  13. Senthilkumar GP (2008) Biochemical studies on the effect of *Terminalia chebula* on the levels of glycoproteins in streptozotocin-induced experimental diabetes in rats. *J Agric Food Chem* 6: 105-115.
  14. Ramachandran S, Rajasekaran A, Manisenthilkumar KT (2012) Investigation of hypoglycemic, hypolipidemic and antioxidant activities of aqueous extract of TP bark in diabetic rats. *Asian Pac J Trop Biomed* 2: 262-268.
  15. Nampoothiri SV, Prathapan A, Cherian OL, Raghu KG, Venugopalan VV, et al. (2011) In vitro antioxidant and inhibitory potential of *Terminalia bellerica* and *Emblica officinalis* fruits against LDL oxidation and key enzymes linked to type 2 diabetes. *Food Chem Toxicol* 49: 125-131.
  16. Anam K, Widharna RM, Kusrini D (2009)  $\alpha$ -Glucosidase Inhibitor Activity of *Terminalia* Species. *Int J Pharm* 5: 277-280.
  17. Wansi JD, Lallemand MC, Chiozem DD, Toze FA, Mbaze LM, et al. (2007)  $\alpha$ -Glucosidase inhibitory constituents from stem bark of *Terminalia superba* (Combretaceae). *Phytochemistry* 68: 2096-2100.
  18. Nagappa AN, Thakurdesai PA, Venkat Rao N, Singh J (2003) Antidiabetic activity of *Terminalia catappa* Linn fruits. *J Ethnopharmacol* 88: 45-50.
  19. Kameswara Rao B, Renuka Sudarshan P, Rajasekhar MD, Nagaraju N, Appa Rao Ch (2003) Antidiabetic activity of *Terminalia pallida* fruit in alloxan induced diabetic rats. *J Ethnopharmacol* 85: 169-172.
  20. Latha R, Daisy P (2011) Insulin-secretagogue, antihyperlipidemic and other protective effects of gallic acid isolated from *Terminalia bellerica* Roxb. in streptozotocin-induced diabetic rats. *Chemico-Biological Interactions* 189: 112-118.
  21. Israni DA, Patel KV, TR G (2010) Anti-hyperlipidemic activity of aqueous extract of *Terminalia chebula* and Gaumutra in high cholesterol diet fed rats. *Int J Pharm Pharm Sci* 1: 48-59.

21. Ram A, Lauria P, Gupta R, Kumar P, Sharma VN (1997) Hypocholesterolaemic effects of *Terminalia arjuna* tree bark. *J Ethnopharmacol* 55: 165-169.
22. Shaila HP, Udupa SL, Udupa AL (1998) Hypolipidemic activity of three indigenous drugs in experimentally induced atherosclerosis. *Int J Cardiol* 67: 119-124.
23. Viswanatha GL, Vaidya SK, Krishnadas N, Rangappa S (2010) Antioxidant and antimutagenic activities of bark extract of *Terminalia arjuna*. *Asian Pac J Trop Biomed* 3: 965-970.
24. Dixit D, Dixit AK, Lad H, Gupta D, Bhatnagar D (2013) Radioprotective effect of *Terminalia Chebula Retzius* extract against  $\gamma$ -irradiation-induced oxidative stress. *Biomedicine and Aging Pathology* 3: 83-88.
25. Pfundstein B, El Desouky SK, Hull WE, Haubner R, Erben G, et al. (2010) Polyphenolic compounds in the fruits of Egyptian medicinal plants (*Terminalia bellerica*, *Terminalia chebula* and *Terminalia horrida*): Characterization, quantitation and determination of antioxidant capacities. *Phytochemistry* 71: 1132-1148.
26. Bajpai M, Pande A, Tewari SK, Prakash D (2005) Phenolic contents and antioxidant activity of some food and medicinal plants. *Int J Food Sci Nutr* 56: 287-291.
27. Fahmy NM, Al-Sayed E, Abdel-Daim MM, Karonen M, Singab A (2015) Protective effect of *Terminalia muelleri* against carbon tetrachloride-induced hepato- nephro toxicity in mice and characterization of its bioactive constituents. *Pharm Biol* 1-11.
28. Eesha BR, Mohanbabu AV, Meena KK, Babu S, Vijay M, et al. (2011) Hepatoprotective activity of TP against paracetamol induced hepatocellular damage in Wistar albino rats. *Asian Pac J Trop Biomed* 4: 466-469.
29. Ghosh J, Das J, Manna P, Sil PC (2010) Protective effect of the fruits of *Terminalia arjuna* against cadmium-induced oxidant stress and hepatic cell injury via MAPK activation and mitochondria dependent pathway. *Food Chem* 123: 1062-1075.
30. Manna P, Sinha M, Pal P, Sil PC (2007) Arjunolic acid, a triterpenoid saponin, ameliorates arsenic-induced cyto-toxicity in hepatocytes. *Chem Biol Interact* 170: 187-200.
31. Lee HS, Jung SH, Yun BS, Lee KW (2007) Isolation of chebulic acid from *Terminalia chebula* Retz. and its antioxidant effect in isolated rat hepatocytes. *Arch Toxicol* 81: 211-218.
32. Kinoshita S, Inoue Y, Nakama S, Ichiba T, Aniya Y (2007) Antioxidant and hepatoprotective actions of medicinal herb, *Terminalia catappa* L. from Okinawa Island and its tannin corilagin. *Phytomedicine* 14: 755-762.
33. Jadon A, Bhadauria M, Shukla S (2007) Protective effect of *Terminalia belerica* Roxb. and gallic acid against carbon tetrachloride induced damage in albino rats. *J Ethnopharmacol* 109: 214-218.
34. Nair JJ, Aremu AO, Van Staden J (2012) Anti-inflammatory effects of *Terminalia phanerophlebia* (Combretaceae) and identification of the active constituent principles. *S Afr J Bot* 81: 79-80.
35. Talwar S, Nandakumar K, Nayak PG, Bansal P, Mudgal J, et al. (2011) Anti-inflammatory activity of TP bark extract against acute and chronic inflammation in rats. *J Ethnopharmacol* 134: 323-328.

36. Tan AC, Hou DX, Konczak I, Tanigawa S, Ramzan I, et al. (2011) Native Australian fruit polyphenols inhibit COX-2 and iNOS expression in LPS- activated murine macrophages. *Food Res Int* 44: 2362-2367.
37. Nair V, Singh S, Gupta YK (2010) Anti-arthritic and disease modifying activity of *Terminalia chebula* Retz. in experimental models. *J Pharm Pharmacol* 62: 1801-1806.
38. Eldeen IM, Elgorashi EE, Mulholland DA, van Staden J (2006) Anolignan B: a bioactive compound from the roots of *Terminalia sericea*. *J Ethnopharmacol* 103: 135-138.
39. Lin CC, Hsu YF, Lin TC (1999) Effects of punicalagin and punicalin on carrageenan-induced inflammation in rats. *Am J Chin Med* 27: 371-376.
40. Fan YM, Xu LZ, Gao J, Wang Y, Tang XH, et al. (2004) Phytochemical and antiinflammatory studies on *Terminalia catappa*. *Fitoterapia* 75: 253-260.
41. Mishra V, Agrawal M, Onasanwo SA, Madhur G, Rastogi P, et al. (2013) Anti- secretory and cyto-protective effects of chebulinic acid isolated from the fruits of *Terminalia chebula* on gastric ulcers. *Phytomedicine* 20: 506-511.
42. Devi RS, Narayan S, Vani G, Shyamala Devi CS (2007) Gastroprotective effect of *Terminalia arjuna* bark on diclofenac sodium induced gastric ulcer. *Chem Biol Interact* 167: 71-83.
43. Gupta M, Mazumder UK, Manikandan L, Bhattacharya S, Senthilkumar GP, et al. (2005) Anti-ulcer activity of ethanol extract of *Terminaliapallida* Brandis. in Swiss albino rats. *J Ethnopharmacol* 97: 405-408.
44. Malekzadeh F, Ehsanifar H, Shahamat M, Levin M, Colwell RR (2001) Antibacterial activity of black myrobalan (*Terminalia chebula* Retz) against *Helicobacter pylori*. *Int J Antimicrob Agents* 18: 85-88.
45. Kathirvel A, Sujatha V (2012) In vitro assessment of antioxidant and antibacterial properties of *Terminalia chebula* Retz. leaves. *Asian Pac J Trop Biomed* 2: S788-S795.
46. Cheng HY, Lin CC, Lin TC (2002) Antiherpes simplex virus type 2 activity of casuarinin from the bark of *Terminalia arjuna* Linn. *Antiviral Res* 55: 447-455.
47. Fyhrquist P, Laakso I, Garcia Marco S, Julkunen-Tiitto R, Hiltunen R (2014) Antimycobacterial activity of ellagitannin and ellagic acid derivate rich crude extracts and fractions of five selected species of *Terminalia* used for treatment of infectious diseases in African traditional medicine. *S Afr J Bot* 90: 1-16.
48. Anam K, Suganda A, Sukandar E, Kardono LBS (2010) Antibacterial effect of component of *Terminalia muelleri* Benth. against *Staphylococcus aureus*. *Int J Pharm* 6: 407-412.
49. Masoko P, Picard J, Eloff JN (2005) Antifungal activities of six South African *Terminalia* species (Combretaceae). *J Ethnopharmacol* 99: 301-308.
50. Nguyen DM, Seo DJ, Lee HB, Kim IS, Kim KY, et al. (2013) Antifungal activity of gallic acid purified from *Terminalia nigrovenulosa* bark against *Fusarium solani*. *Microb Pathog* 56: 8-15.
51. Silva O, Duarte A, Pimentel M, Viegas S, Barroso H, et al. (1997) Antimicrobial activity of *Terminalia macroptera* root. *J Ethnopharmacol* 57: 203-207.
52. Silva O, Ferreira E, Vaz Pato M, Caniça M, Gomes ET (2002) In vitro anti- *Neisseria*

- gonorrhoeae activity of *Terminalia macroptera* leaves. FEMS Microbiol Lett 217: 271-274.
53. Silva O, Viegas S, de Mello-Sampayo C, Costa MJ, Serrano R, et al. (2012) Anti-*Helicobacter pylori* activity of *Terminalia macroptera* root. Fitoterapia 83: 872-876.
  54. Kuete V, Tabopda TK, Ngameni B, Nana F, Tshikalange TE, et al. (2010) Antimycobacterial, antibacterial and antifungal activities of *Terminalia superba* (Combretaceae). S Afr J Bot 76: 125-131.
  55. Carpano SM, Spegazzini ED, Rossi JS, Castro MT, Debenedetti SL (2003) Antifungal activity of *Terminalia australis*. Fitoterapia 74: 294-297.
  56. Srivastava SK, Srivastava SD, Chouksey BK (2001) New antifungal constituents from *Terminalia alata*. Fitoterapia 72: 106-112.
  57. Saleem A, Husheem M, Härkönen P, Pihlaja K (2002) Inhibition of cancer cell growth by crude extract and the phenolics of *Terminalia chebula* retz. fruit. J Ethnopharmacol 81: 327-336.
  58. Row LR, Raju RR (1967) Chemistry of *Terminalia* species-XI: Isolation of 3,4,3'-O-trimethyl flavellagic acid from TP Roth. Tetrahedron 23: 879-884.
  59. Nadkarni AK: TP Roxb. In Dr KM Nadkarni's Indian Materia Medica. Volume 1. 3rd edition. Mumbai, India: Popular Prakashan;1996:931.
  60. Talwar S, Nandakumar K, Nayak PG, Bansal P, Mudgal J, Mor V, Rao CM, Lobo R: Anti-inflammatory activity of TP bark extract against acute and chronic inflammation in rats. J Ethnopharmacol 2011, 134:323-328.