

# Review Article

## Recent Approaches of Drugs and Vaccines Used in Covid 19 : A Review

### ABSTRACT:

A novel SARS-CoV-2 coronavirus developed in December 2019, triggering a life-threatening pneumonia outbreak in China, and has since expanded worldwide, producing a pandemic. Because of the urgent need to control the disease and the dearth of specific and effective treatment options, FDA-approved medicines that have proven efficacy against comparable viruses are being used. In vitro, chloroquine, remdesivir, lopinavir/ritonavir or ribavirin are effective at inhibiting SARS-CoV-2. The preliminary findings of a variety of clinical trials utilising various chloroquine or hydroxychloroquine delivery procedures all indicate a positive outcome. They may not be effective in cases of consistently high viremia, and data on ivermectin (another antiparasitic medication) is currently unavailable. Intriguingly, azithromycin, a macrolide antibiotic, in conjunction with hydroxychloroquine may provide therapeutic benefits. Favipiravir, tocilizumab, and azithromycin types are among the other treatment options being investigated.

*Keywords: SARS-CoV 2, MOA, Drugs, Vaccines*

### 1. INTRODUCTION:

Coronavirus sickness, a highly communicable virus because of the SARS, has made a huge impact on global demography; killing over 3.8 People around the world who have died because of the pandemic, causing it the worst worldwide health disaster since the 1918 disaster. After the first case was it was found in Dec 2019 in Wuhan, Hubei Province, China. The disease increases quickly, prompting the WHO to declare it a global pandemic on March 11, 2020. Since it was COVID disease has ravaged the world and has been declared a global pandemic in several countries throughout the world, and it has overloaded several medical care systems. The covid disease continues too many nations are experiencing a second or third wave of infections, which can be linked in large part to the rise of variant versions of this virus. The fact that significant Clinical research has contributed to a better understanding of covid disease and Has turned into a problem growing concern, as Covid disease continues to cause problems around the world, with most nations are in the early stages of a second or third wave of breakouts[1]. Since the outbreak began, four SARS-CoV-2 VOCs have been detected, according to a current WHO epidemiological statement as of June 22, 2021: Alpha was the UK reported the first variation of concern in late Dec 2020; Beta was first shown in South Africa in Dec 2020; Gamma was first identified in early Jan 2021 in Brazil. In Dec 2020, Delta for the first time was discovered in India. This review research aims to give a comprehensive summary of the most recent pharmacological treatment options for Covid disease. The efficacy of different existing COVID-19 and variant vaccines is also briefly discussed in this review (1).

### 2. COVID19 SYMPTOMS:

Different persons are affected by the coronavirus in different ways. The majority of infected patients will experience mild to moderate symptoms and will recover without the need for hospitalization. The following are the most common signs and symptoms:

- Fever/cough
- Tiredness
- A change in flavour or odour

#### 2.1 Fewer common symptoms include:

- Sore throat pains and aches cause a headache.
- Diarrhoea,

- 
- The appearance of a rash on the skin or irritated eyes, or colouration of the fingers or toes
  - Breathing difficulties, shortness of breath, speech or mobility loss, and confusion are all serious signs.
  - Chest discomfort

Mild symptomatic people who manage their symptoms at home. When a person is infected with the virus, symptoms appear after 5–6 days, but it can take up to 14 days [2].

## 2.2 Risk factors:

- Chronic kidney disease is a condition that affects the kidney so Possessive
- Obesity
- Heart failure or coronary artery disease are serious heart disorders.
- Sickle cell disease is a type of sickle cell illness.
- Diabetes type 2

The following are some of the conditions that can cause severe COVID-19 illness:

- Asthma ranges from mild to severe.
- Diseases that influence your brain's blood arteries and blood flow
- Cystic fibrosis is a disease that affects the lungs.
- Blood pressure that is too high
- A compromised immune system as a result of a blood or bone marrow transplant or corticosteroid prescription
- Dementia
- Hepatitis is a disease of the liver.
- Pregnancy
- Lung tissue that has been damaged or scarred (pulmonary fibrosis)
- Smoking
- Thalassemia
- Type 1 diabetes
- Depression
- Anxiety
- Schizophrenia

It has symptoms that are comparable to toxic shock and illness, which causes inflammation in the blood vessels of children [3].

According to experts, the following symptoms were the most common in COVID-19 patients :

- 99.9% fever
- 70% exhaustion
- 59 per cent of coughing
- 40% suffer from a lack of appetite
- 35% of body aches
- 31% of people experience shortness of breath.
- 27% mucus/phlegm

## 3. DRUGS:

### 3.1 Remdesivir:

Remdesivir is a medication that is a nucleoside analogue. It is justified to employ a broad-spectrum antiviral with invitro coronavirus action. The action mechanism of remdesivir is the Remdesivir-triphosphate (RDV-TP) is a mono phosphoramidite prodrug of remdesivir that inhibits RNA-dependent RNA polymerases (RdRps) [4]. Adenosine triphosphate is used by Remdesivir-TP to incorporate into viral RNA strands. Once fused to viral RNA at a location I, RDV-TP terminates the RNA combination at position i+3. Because RDV-TP does not produce rapid chain termination, it

---

appears to resist viral exoribonuclease editing. In Table No.1. Remdesivir's method of action has been shown in vitro to work against the Arenaviridae, Flaviviridae, Filoviridae, Paramyxoviridae, Pneumoviridae, and Coronaviridae virus families, indicating that it has broad antiviral activity [4]. Remdesivir's activity against the Coronaviridae family was initially demonstrated in 2017, sparking interest in the drug as a potential covid disease treatment [8]. Remdesivir has been confirmed as a non-obligate chain terminator of RdRp from coronavirus and similar coronaviruses, as well as MERS-CoV, in several clinical trials for covid disease [12, 13].

### 3.2 Chloroquine and hydroxychloroquine:

Its anti-SARS-CoV-2 efficacy in vitro and immunomodulatory features justify its use. Through their method of action, chloroquine and hydroxychloroquine block viral enzymes [14]. Inhibition of the ACE2 cell receptor, acidification of the cell membrane's outer surface to impede viral combination, and immunomodulation of cytokine discharge are all options [15]. Based on hydroxychloroquine's in vitro antiviral activity, the optimal dose design as the therapy of Coronavirus 2 SARSCoV-2 was predicted. However, preliminary results from randomised controlled studies in hospitalised patients comparing hydroxychloroquine with or without azithromycin to placebo indicated no difference in clinical status or overall mortality [16, 17]. According to the findings of randomised controlled studies, using hydroxychloroquine as a postexposure preventative infection with the covid virus or symptoms of covid sickness was not prevented [18].

### 3.3 Lopinavir/Ritonavir:

Classification of Lopinavir/Ritonavir is an Inhibitor of HIV protease. Additional coronaviruses may be active, according to in vitro and animal model research. In Table No.1 it works by inhibiting covid virus reproduction and activity [20]. Lopinavir/ritonavir is an FDA-approved HIV medication combination, that was recommended as an antiviral treatment against covid disease in the early phases of the pandemic. A randomised control pilot research found no advantage from the drug therapy compared to a minimum standard of care in individuals hospitalized has the severe covid disease. In both hospitalized and non-hospitalized patients, it is currently not advised for covid disease treatment [21].

### 3.4 Ivermectin:

Classification of Ivermectin is Antiparasitic. The rationale for use of Ivermectin is it prohibits covid virus replication in cell cultures.

In any case, pharmacokinetic and pharmacodynamic studies suggest that dosages up to 100 times higher than those approved in people would be required to achieve the plasma focuses required for the antiviral effect seen in vitro [23].

In Table No.1 Ivermectin is transported within the cell protein that the virus employs to increase infection by inhibiting the antiviral response of the host. Ivermectin is an anti-parasitic medicine that has been licensed by the FDA and is used to take care of Covid disease all over the world. It was created after an in vitro investigation showed that it could prevent coronavirus proliferation [24]. A randomised controlled experiment in which 476 adult Covid disease patients were randomly assigned Ivermectin 300mcg/kg body weight or placebo for five days did not result in substantial improvement or the desired symptom relief. Both hospitalized and non-hospitalized individuals should avoid ivermectin treatment for Covid disease [24].

### 3.5 Favipiravir:

Favipiravir is an inhibitor of RNA polymerase that is dependent on RNA currently being researched. The following rationale for its use favipiravir is a broad-spectrum antiviral that has been discovered effective in the laboratory against RNA viruses.

Favipiravir acts by blocking the RNA polymerase that is an RNA-dependent enzyme, which prevents the synthesis of viral RNA [25]. China's NMPA has included this drug in a list of potential covid virus treatments. Favipiravir group participants' lab tests for the covid virus were negative after the treatment was administered [26]. In addition to interferon is 5 million U twice daily via aerosol inhalation, 35 patients with a median age of 43 years were given 1600mg twice a day as a loading dosage and 600mg twice daily. In a study conducted, 45 patients were given lopinavir 400mg or ritonavir 100mg (RTV) every day twice by aerosol inhalation, as well as 5 million U IFN-a twice a day

---

(middle age was 49 a long time). They noticed a short time for the virus to be cleared and considerable imaging of the chest has improved in the FPV group with low ADRs. In a study conducted, 45 patients were given lopinavir 400mg or ritonavir 100mg (RTV) twice a day by aerosol inhalation, as well as IFN-a 5 million U twice daily. They noticed a short In the FPV group; there was a significant improvement in chest imaging and a minimal rate of adverse effects. The results of a recent trial with 120 coronavirus patients in Wuhan, China [27].

### 3.6 REGN-COV2 (Casirivimab and Imdevimab)

Classification of Casirivimab is a monoclonal antibody. Casirivimab is a recombinant human IgG1 monoclonal antibody that binds to the coronavirus spike protein's receptor binding domain, which is involved in the viral connection, combination, and cell entry. The coronavirus spike protein is destroyed by casirivimab in conjunction with imdevimab [28]. In Table No.1 Casirivimab is a cocktail of two non-competing IgG1 antibodies that have been shown to reduce viral load in vivo by targeting the RBD on the coronavirus spike protein, preventing virus infection (14). In a break analysis of 275 patients from a continuous double-blinded test including non-hospitalized patients with corona disease who were given a placebo, 2.4g of casirivimab, or Imdevimab at random. The safety profile of this cocktail antibody was also evaluated in this investigation. According to early data from a Phase 3 trial of REGN-COV, corona disease resulted in a 70% decrease in hospitalization or a 70% reduction in non-hospitalized patients. There is in vitro data available on the impact of REGN-COV2 on the two novel coronavirus variants of concern that demonstrate retained activity [29].

### 3.7 Glucocorticoids:

Dexamethasone is a glucocorticoid hormone. Corticosteroids decrease Leukocyte migration to inflammatory regions, as well as vasodilation and capillary permeability. Corticosteroids bind to the glucocorticoid receptor, causing gene expression alterations with a wide range of downstream effects that might take hours to days to manifest. In Table No.1 Glucocorticoids inhibit phospholipase A2, which prevents the formation of arachidonic acid derivatives, as well as Death and demargination of neutrophils they inhibit the transcription factors NF-Kappa B and other inflammatory transcription factors while boosting the production of anti-inflammatory genes like interleukin-10. Corticosteroids at lower amounts are anti-inflammatory, however, higher doses suppress the immune system. Sodium levels rise and potassium levels fall when high amounts of glucocorticoids bind to the mineralocorticoid receptor over time [30]. Corona disease has been associated with cytokine-induced inflammatory lung damage, which is characterised by an increase in inflammatory markers. Glucocorticoid effectiveness in covid disease patients was not thoroughly reported in the early phases of the epidemic. The recovery preliminary study found that giving dexamethasone to hospitalized people was given to patients with clinically suspected or lab-confirmed coronavirus. Dexamethasone or frequent consideration for patients who received intrusive mechanical breathing or oxygen support for clinically suspected or lab-confirmed coronavirus had a reduced 28-day death rate than those who did not. Dexamethasone, alone or in combination with remdesivir, is used in hospitalized patients who require supplementary oxygen or non-obtrusive or invasive mechanical respiration is currently regarded as the standard of therapy, depending on the severity of the condition [31].

### 3.8 Azithromycin

Classification of Azithromycin is Macrolide Antibacterials and the Rationale for use is macrolides may have immunomodulatory characteristics, and azithromycin may prevent bacterial superinfection, making them suitable for adjuvant therapy.

To grow, Bacteria require a unique protein synthesis pathway that is facilitated by ribosomal proteins [32, 33]. It obstructs bacterial protein synthesis by preventing the movement phase the binding of the 50S ribosomal subunit as well as protein synthesis Label. As a result, various bacterial illnesses have been brought under control. Macrolides In pulmonary inflammatory illnesses, they exhibit immunomodulatory effects [34]. They may aid in reducing cytokine overproduction associated with respiratory virus infections. Preventing cytokines from prompting neutrophils (PMNs) to chemotaxis to the lungs is one example of immunomodulatory approaches. Mucus hypersecretion is inhibited (20). In Table No.1 Azithromycin is a macrolide antibiotic is a type of antibiotic that is used to treat bacterial infections [35]. This macrolide possesses antibacterial, antiviral, and immunomodulatory properties, making it potentially useful in viral infections such as the corona virus. The high tissue accumulation compensates for the poor absorption, resulting in a 37 per cent oral bioavailability. Azithromycin is

400 to 1,000 times more concentrated in epithelial cells, fibroblasts, lymphocytes, and alveolar macrophages than in blood [36, 37]. Chemotactic drug delivery raises local drug concentrations by releasing stored azithromycin through blood phagocytes and other cells that move to inflamed and dirty tissues. As a result of this, azithromycin has a half-life of 68–79 hours. This medication has an amazing ability to permeate lung tissue and keep medication concentrations stable [38].

### 3.9 Tocilizumab:

Tocilizumab is a monoclonal antibody that has been shown to target the interleukin-6 receptors alpha receptor demonstrated to be useful in the treatment of rheumatoid arthritis. The information on how this agent was utilized is distorted. When compared to placebo, Tocilizumab did not result in a significant change in clinical status or a decrease in 28-day mortality in a randomised controlled trial comprising 438 hospitalised patients with severe coronavirus pneumonia, 294 of them were given tocilizumab and 144 were given placebo [39]. Another randomized, two-fold visually impaired fake treatment controlled trial involving persons with confirmed extreme corona disease that comprised A study of 243 patients who were randomly assigned to receive tocilizumab or a placebo discovered that it was ineffective in preventing death [40]. The REMAP-CAP and RECOVERY preliminary (not yet released) showed a mortality advantage in individuals with fast respiratory decompensation [41]. Interleukin 6 is a pro-inflammatory cytokine produced by T cells, B cells, lymphocytes, monocytes, and fibroblasts. Antibody synthesis, cytotoxic T-cell isolation, and development are all steps in the cytotoxic T-cell process are all boosted by IL-6. By binding to IL-6 receptors in the solvent and on the membrane layer, tocilizumab suppresses IL-6-mediated inflammation. C-reactive protein, serum amyloid A, fibrinogen, haptoglobin, and -1-antichymotrypsin are all blood markers for inflammation and all produced in response to IL-6, but not fibronectin, egg whites, or transferrin [42].

**Table.1 Covid19 Drugs**

Drugs	Classification	Rationale for use	Brand name	Generic name	Mechanism of action	Route of Administration
Remdesivir	Antiviral - Nucleoside Analogue	Broad spectrum	Veklury	Remdesivir	Mono phosphoramidate prodrug of (RDV - TP)	Intravenous
Chloroquine and Hydroxychloroquine	Antimalarial	Invitro against corona virus	Aralen	Chloroquine	Inhibition of viral enzymes. Acidification of the surface of the cell membrane.	Intramuscular or Subcutaneous.
Lopinavir or Ritonavir	Inhibitor of HIV protease	Other coronaviruses may have activity based on in vitro and animal model investigations	Kaletra	Lopinavir/Ritonavir	Suppress corona virus replication	Oral
Ivermectin	Antiparasitic	Inhibits the replication of coronavirus	Sklice, Soolantara, Stromectol	Ivermectin	Suppress the virus infections.	Oral, Intravenous (IV), Topical route

Favipiravir	Antiviral	Antiviral <b>has</b> broad spectrum efficacy in vitro against RNA viruses.	Avigan	Favipiravir	RNA polymerase (RdRp) inhibitor that inhibits viral RNA synthesis.	Oral, IV
Azithromycin	Macrolide antibacterial	Macrolides may have immunomodulatory characteristics and can be used as an adjuvant therapy to prevent bacterial infection.	Azasite	Azithromycin	Immunomodulatory properties in pulmonary inflammatory disorders, Inhibit cytokine production, accelerating neutrophil apoptosis.	Iv, Oral
Casirivimab and Imdevimab	Monoclonal antibody	Cytokine release syndrome	REGEN-COV, Ronaprave	Casirivimab and Imdevimab	Inactivation of <b>the</b> spike protein	Iv, Subcutaneous
Tocilizumab	Monoclonal antibody	Cytokine release syndrome	Actemra, RoActemra	Tocilizumab	T-cell activation	Iv, Subcutaneous

#### 4. VACCINES;

##### 4.1 Covishield: (ChAdOx1 nCoV-19 antibody):

In Table No.2 After two doses, a preliminary evaluation of a 70.4% clinical efficacy and a favourable safety profile **has** been proven in a multinational randomised control trial to 64% protect to avoid corona disease. In various places throughout the world that have not yet gotten formal approval, the Covishield has been approved to prevent corona disease. The FDA has granted emergency use authorisation or approval in the US [43].

##### 4.2 Moderna (mRNA-1273):

People receiving 2 doses of Moderna antibody 28 days apart in a trial 3 randomly selected, observer-blinded, placebo-controlled **trials** were found to be 94.1% effective in preventing corona disease, with no safety concerns other than short-term local and systemic reactions(26). Based on the results of this vaccine effectiveness preliminary trial, the FDA issued a EUA for the use of the **modern** antibody to prevent corona disease on December 18, 2020. Up to seven distinct Vaccines were developed natively by Covaxin (India), Sputnik V (Russia), and CoronaVac (China), incorporating protein-based and inactivated antibodies certified for crisis use in various countries throughout the world to prevent corona disease [43].

**Table.2 Covid19 Vaccines**

Vaccines	Scientific name	Firm	Dosing schedule	Route Administration of	Storage
----------	-----------------	------	-----------------	-------------------------	---------

<b>COVISHIELD (WHO Approved)</b>	ChAdOx1nCoV-19	M/s Serum Institute of India Pvt. Ltd.	Two doses Day 01 & Day 84	Intramuscular	2-8°C
<b>COVAXIN (WHO Approved)</b>	Whole-Virion Inactivated SARSCoV-2 Vaccine	M/s Bharat Biotech	Two doses, Day 0 & 28	Intramuscular	2-8°C
<b>MODERNA</b>	mRNA-1273 COVID-19 vaccine	M/s Cipla Ltd. (Importer)	Two doses, Day 0 & 28	Intramuscular	-25°C to 15°C
<b>JOHNSON AND JOHNSON (WHO Approved)</b>	Ad26.COV2-S (recombinant)	M/s Johnson & Johnson Pvt. Ltd. (Importer) & Biological Limited	Single-dose	Intramuscular	-25°C to 15°C & 2-8°C
<b>SPUTNIK-V</b>	Recombinant Adenovirus vector Based SARS-CoV-2 liquid vaccine Gam COVID Vac	M/s Dr Reddy's Lab. Ltd. (Importer)	Two doses, Day 0 & Day 21	Intramuscular	-18°C
<b>ZYCOV-D</b>	Novel CoronaVirus-2019-nCov Vaccine (recombinant DNA)	Cadila Healthcare Limited	Three doses (Day 0, 28 And 56)	Intradermal	2-8°C
<b>NOVAVAX NUVAXOVID (WHO Approved)</b>	Protein subunit Vaccine	Serum Institute of India	Two Dose	Intramuscular	2-8°C
<b>COMIRNATY (WHO Approved)</b>	RNA Based vaccine also referred to as Tozinameran, BNT162b2	Pfizer/BioNTech	Two Dose (Three week interval)	Intramuscular	2-8°C
<b>CORONAVAC (WHO Approved)</b>	Inactivated SARSCoV-2 Vaccine	Sinovac Biotech	Two Dose (Four week interval)	Intramuscular	2-8°C

## 5. CONCLUSION:

COVID-19 has been tested using a variety of antiviral medications, both separately and in combination, that have previously been used against SARS and MERS, as well as FDA-approved and laboratory tests. Despite extensive research, no viable COVID-19 antiviral medication or vaccine is currently available. While many innovative therapeutic techniques are being developed more clinical trials are essential during these key periods to ensure the safety and efficacy of innovative medications. While many medications are successful against a variety of respiratory viral infections, including influenza and some coronavirus strains, building resistance to existing antiviral medications remains a significant concern.

## 6. CONFLICT OF INTEREST:

---

Authors are declared that there is no conflict of interest with each other. The products used for this research are commonly and predominantly used products in our area of research and country. There is no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by the personal efforts of the authors.

## 7. REFERENCES

1. Chan J F, To K K, Tse H, Jin D Y, & Yuen K Y. (2013). Interspecies transmission and emergence of novel viruses: lessons from bats and birds. *Trends in microbiology*, 21(10), 544–555.
2. Lei J, Kusov Y, & Hilgenfeld R. (2018). Nsp3 of coronaviruses: Structures and functions of a large multi-domain protein. *Antiviral research*, 149, 58–74.
3. Agostini M L, Andres E L, Sims A C, Graham R L, Sheahan T P, Lu X, et al. (2018). Coronavirus Susceptibility to the Antiviral Remdesivir (GS-5734) Is Mediated by the Viral Polymerase and the Proofreading Exoribonuclease. *mBio*, 9(2), e00221-18.
4. Wang M, Cao R, Zhang L, Yang X, Liu J, Xu M, et al. (2020). Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell research*, 30(3), 269–271.
5. Brown AJ, Won JJ, Graham RL, Dinnon KH, Sims AC, Feng JY, et al. (2019). Broad spectrum antiviral remdesivir inhibits human endemic and zoonotic deltacoronaviruses with a highly divergent RNA dependent RNA polymerase. *Antiviral research*, 169, 104541.
6. De Wit, Feldmann F, Cronin J. Prophylactic and therapeutic remdesivir (GS-5734) treatment in the rhesus macaque model of MERS-CoV infection. *Proc Natl Acad Sci U S A*. 2020 Feb 13. [Epub ahead of print] PMID: 32054787
7. Ko W, Rolain J, Lee N. Arguments in favor of remdesivir for treating SARS-CoV-2 infections. *Int J Antimicrob Agents* 2020 Mar 6. [Epub ahead of print] PMID: 32147516
8. Gordon CJ, Tchesnokov EP, Feng JY. The antiviral compound remdesivir potently inhibits RNA-dependent RNA polymerase from Middle East respiratory syndrome coronavirus. *J Biol Chem* 2020 Feb 24. [Epub ahead of print] PMID: 32094225
9. Warren TK, Jordan R, Lo MK, Ray AS, Mackman RL, Soloveva V, et al. (2016). Therapeutic efficacy of the small molecule GS-5734 against Ebola virus in rhesus monkeys. *Nature*, 531(7594), 381–385.
10. Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, Kalil AC, et al. Study Group Members (2020). Remdesivir for the Treatment of Covid-19 - Final Report. *The New England journal of medicine*, 383(19), 1813–1826.
11. Goldman JD, Lye D, Hui DS, Marks KM, Bruno R, Montejano R, et al. GS-US-540-5773 Investigators (2020). Remdesivir for 5 or 10 Days in Patients with Severe Covid-19. *The New England journal of medicine*, 383(19), 1827–1837.
12. Spinner CD, Gottlieb RL, Criner GJ, Arribas López JR, Cattelan AM, Soriano Viladomiu A, et al. GS-US-540-5774 Investigators (2020). Effect of Remdesivir vs Standard Care on Clinical Status at 11 Days in Patients With Moderate COVID-19: A Randomized Clinical Trial. *JAMA*, 324(11), 1048–1057.

- 
13. Zhang R, & Mylonakis E. (2021). In inpatients with COVID-19, none of remdesivir, hydroxychloroquine, lopinavir, or interferon  $\beta$ -1a differed from standard care for in-hospital mortality. *Annals of internal medicine*, 174(2), JC17.
  14. Gao J, Tian Z, & Yang X. (2020). Breakthrough: Chloroquine phosphate has shown apparent efficacy in treatment of COVID-19 associated pneumonia in clinical studies. *Bioscience trends*, 14(1), 72–73.
  15. Wang M, Cao R, Zhang L, Yang X, Liu J, Xu M, et al.(2020). Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell research*, 30(3), 269–271.
  16. RECOVERY Collaborative Group, Horby P, Mafham M, Linsell L, Bell J L, Staplin N, Emberson JR, et al.(2020). Effect of Hydroxychloroquine in Hospitalized Patients with Covid-19. *The New England journal of medicine*, 383(21), 2030–2040.
  17. Zhang R, & Mylonakis E. (2021). In inpatients with COVID-19, none of remdesivir, hydroxychloroquine, lopinavir, or interferon  $\beta$ -1a differed from standard care for in-hospital mortality. *Annals of internal medicine*, 174(2), JC17.
  18. Mitjà O, Corbacho-Monné M, Ubals M, Alemany A, Suñer C, Tebé C, et al. BCN-PEP-CoV2 Research Group (2021). A Cluster-Randomized Trial of Hydroxychloroquine for Prevention of Covid-19. *The New England journal of medicine*, 384(5), 417–427.
  19. Boulware DR, Pullen M F, Bangdiwala AS, Pastick KA, Lofgren SM, Okafor EC, et al. (2020). A Randomized Trial of Hydroxychloroquine as Postexposure Prophylaxis for Covid-19. *The New England journal of medicine*, 383(6), 517–525.
  20. Chu CM, Cheng VC, Hung IF, Wong MM, Chan KH, Chan KS, et al. HKU/UCH SARS Study Group (2004). Role of lopinavir/ritonavir in the treatment of SARS: initial virological and clinical findings. *Thorax*, 59(3), 252–256.
  21. Liu X, Wang XJ. Potential inhibitors for 2019-nCoV coronavirus M protease from clinically proven medicines. *J Genet Genomics* 2020 Feb 13. [Epub ahead of print] PMID: 32173287
  22. Cao B, Wang Y, Wen D, Liu W, Wang J, Fan G, et al.(2020). A Trial of Lopinavir-Ritonavir in Adults Hospitalized with Severe Covid-19. *The New England journal of medicine*, 382(19), 1787–1799.
  23. Grieco DL, Menga LS, Cesarano M, Rosà T, Spadaro S, Bitondo MM, et al. COVID-ICU Gemelli Study Group (2021). Effect of Helmet Noninvasive Ventilation vs High-Flow Nasal Oxygen on Days Free of Respiratory Support in Patients With COVID-19 and Moderate to Severe Hypoxemic Respiratory Failure: The HENIVOT Randomized Clinical Trial. *JAMA*, 325(17), 1731–1743.
  24. Caly L, Druce JD, Catton MG, Jans DA, & Wagstaff KM. (2020). The FDA-approved drug ivermectin inhibits the replication of SARS-CoV-2 in vitro. *Antiviral research*, 178, 104787.
  25. López-Medina E, López P, Hurtado IC, Dávalos DM, Ramirez O, Martínez E Dí, et al. (2021). Effect of Ivermectin on Time to Resolution of Symptoms Among Adults With Mild COVID-19: A Randomized Clinical Trial. *Shiraki K and Daikoku T. Favipiravir, an anti-influenza drug against life-threatening RNA virus infections. Pharmacol Ther.* 2020 Feb 22. [Epub ahead of print] PMID:32097670 *MA*, 325(14), 1426–1435.
  26. Tegally H, Wilkinson E, Giovanetti M, Iranzadeh A, Fonseca V, Giandhari J, et al. (2021). Detection of a SARS-CoV-2 variant of concern in South Africa. *Nature*, 592(7854), 438–443.
  27. Cai Q, Yang M, Liu D, Chen J, Shu D, Xia J, et al.(2020). Experimental Treatment with Favipiravir for COVID-19: An Open-Label Control Study. *Engineering (Beijing, China)*, 6(10), 1192–1198.

- 
28. Baum A, Ajithdoss D, Copin R, Zhou A, Lanza K, Negron N, et al. (2020). REGN-COV2 antibodies prevent and treat SARS-CoV-2 infection in rhesus macaques and hamsters. *Science (New York, N.Y.)*, 370(6520), 1110–1115.
  29. Weinreich DM, Sivapalasingam S, Norton T, Ali S, Gao H, Bhore R, et al. Trial Investigators (2021). REGN-COV2, a Neutralizing Antibody Cocktail, in Outpatients with Covid-19. *The New England journal of medicine*, 384(3), 238–251.
  30. Yasir M, Goyal A, Bansal P & Sonthalia S. (2021). *Corticosteroid Adverse Effects*. In StatPearls. StatPearls Publishing.
  31. RECOVERY Collaborative Group, Horby P, Lim W S, Emberson JR, Mafham M, Bell JL, Linsell L, et al. (2021). Dexamethasone in Hospitalized Patients with Covid-19. *The New England journal of medicine*, 384(8), 693–704.
  32. Gautret P, Lagier JC, Parola P, Hoang VT, Meddeb L, Mailhe M, et al. (2020). Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *International journal of antimicrobial agents*, 56(1), 105949.
  33. Amsden G. W. (2005). Anti-inflammatory effects of macrolides--an underappreciated benefit in the treatment of community-acquired respiratory tract infections and chronic inflammatory pulmonary conditions?. *The Journal of antimicrobial chemotherapy*, 55(1), 10–21.
  34. Kanoh, S., & Rubin, B. K. (2010). Mechanisms of action and clinical application of macrolides as immunomodulatory medications. *Clinical microbiology reviews*, 23(3), 590–615.
  35. Beigel JH, Tomashek KM, Dodd LE, et al. Remdesivir for the treatment of Covid-19 — preliminary report. *N Engl J Med*. 2020 [cited 2020May 25]; NEJMoa2007764. Available from: <http://www.nejm.org/doi/10.1056/NEJMoa2007764>
  36. Tyteca D, Van Der Smissen P, Mettlen M, Van Bambeke F, Tulkens PM, Mingeot-Leclercq MP, & Courtoy, P. J. (2002). Azithromycin, a lysosomotropic antibiotic, has distinct effects on fluid-phase and receptor-mediated endocytosis but does not impair phagocytosis in J774 macrophages. *Experimental cell research*, 281(1), 86–100.
  37. Zheng S, Matzneller P, Zeitlinger M, & Schmidt S. (2014). Development of a population pharmacokinetic model characterizing the tissue distribution of azithromycin in healthy subjects. *Antimicrobial agents and chemotherapy*, 58(11), 6675–6684.
  38. Rosas IO, Bräu N, Waters M, Go RC, Hunter BD, Bhagani S, et al. (2021). Tocilizumab in Hospitalized Patients with Severe Covid-19 Pneumonia. *The New England journal of medicine*, 384(16), 1503–1516.
  39. Stone JH, Frigault MJ, Serling-Boyd NJ, Fernandes AD, Harvey L, Foulkes AS, et al. BACC Bay Tocilizumab Trial Investigators (2020). Efficacy of Tocilizumab in Patients Hospitalized with Covid-19. *The New England journal of medicine*, 383(24), 2333–2344.
  40. REMAP-CAP Investigators, Gordon AC, Mouncey PR, Al-Beidh F, Rowan KM, Nichol AD, Arabi YM, et al.(2021). Interleukin-6 Receptor Antagonists in Critically Ill Patients with Covid-19. *The New England journal of medicine*, 384(16), 1491–1502.
  41. Tanaka T, Narazaki M, & Kishimoto T. (2014). IL-6 in inflammation, immunity, and disease. *Cold Spring Harbor perspectives in biology*, 6(10), a016295.
  42. Voysey M, Clemens SAC, Madhi SA, Weckx LY, Folegatti PM, Aley PK, et al. Oxford COVID Vaccine Trial Group. Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa, and the UK. *Lancet*. 2021 Jan 09;397(10269):99-111.

- 
43. Baden LR, El Sahly HM, Essink B, Kotloff K, Frey S, Novak R, et al. COVE Study Group (2021). Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *The New England journal of medicine*, 384(5), 403–416.