

Original Research Article

ASSESSMENT OF RELATION OF SERUM URIC ACID & MICRO ALBUMINURIA IN THE PATIENTS OF DIABETES MELLITUS TYPE-2 FOR EARLY DETECTION OF DIABETIC NEPHROPATHY

ABSTRACT

Background: Diabetic Nephropathy is one of the serious and life perilous complications of diabetes mellitus. Prolong duration of diabetes, no proper care and management, poor glycemic control in diabetic nephropathy can lead to End Stage Renal Disease (ESRD).

Objective: To assess the relation of serum uric acid & micro albuminuria in the patients of diabetes mellitus for early detection of diabetic nephropathy.

Methodology: This cross-sectional study was conducted in the Department of Medicine, LUMHS Jamshoro, with a total of 80 diabetic patients as case study subjects and 80 non-diabetic healthy participants as controls. Random glucose was measured using the gluco oxidase technique, and serum uric acid was measured using the uricase enzyme method in a calorimeter. Immunoturbidimetry kit technique was used to calculate microalbuminuria.

Results: The mean value of random blood sugar (RBS) in the control group was 135 ± 10.11 mg/dl, but RBS in the case study group was 224 ± 13.52 mg/dl, indicating that RBS in the case study group was extremely significant ($P < 0.001$). The mean serum uric acid level in the control group was 4.7 ± 0.6 mg/dl, while it was 6.6 ± 1.4 mg/dl in the case study group, which is

highly significant ($P < 0.001$) with correlation ($r = 0.34$). Albumin in urine in the control group was 10.2 ± 1.78 mg/gm, while it was 54 ± 10.34 mg/gm in the case study group, which is a significant ($P < 0.05$) increase in the case study group with correlation ($r = 0.28$). In diabetics, serum uric acid and microalbuminuria show a positive connection.

Conclusion: This study concluded that there is strong positive relation of serum uric acid and micro albuminuria in diabetic patients for the early detection of diabetic nephropathy.

Key Words: Serum Uric Acid, Micro albuminuria, Diabetes Mellitus, Diabetic nephropathy

INTRODUCTION

Hyperglycemia, as well as disturbances in carbohydrate, protein, and lipid metabolism, are characteristics of the metabolic syndrome, which is associated with diabetes mellitus (DM).¹Diabetes mellitus is the leading cause of morbidity worldwide.²Diabetes today affects more than 4% of the world's population.³Diabetic mellitus is more common in underdeveloped nations. Nearly 10% of Pakistan's population is affected with diabetes mellitus.⁴Sedentary behavior, high BMI, obesity, and poor eating habits are all risk factors for the development of diabetes and related consequences such as nephropathy, retinopathy, and atherosclerosis.^{5,6,7}Diabetic nephropathy is one of the most significant consequences of glomerulosclerosis, and if left untreated, it can lead to end-stage renal disease, which is fatal to humans.^{8,9}Diabetic patients with nephropathy have a higher risk of death, with a 37 to 40 times increase in mortality.⁹Micro albumin is one of the predictive factors for nephropathy in people with type 2 diabetes.^{10,11}When diabetic nephropathy progresses, albumin may be present in the urine sample at the time of diagnosis, but levels between 30 and 300 mg/day indicate micro albuminuria. When diabetic nephropathy progresses, macro albumin urea may develop.^{12, 13}

The purine metabolism produces uric acid as a byproduct molecule.¹⁴ Nearly 30% of it is used in the small intestine, with the remaining 50-60% excreted through the renal system.¹⁵ Due to a lack of the uricase enzyme in humans and a significant amount of uric acid in serum, humans have a higher rank of serum uric acid levels than other animals. Uric acid in serum is mostly excreted by urine.¹⁶ When renal activities are interrupted owing to a disease in the kidney, it can lead to an increase in blood uric acid levels due to a decrease in uric acid outflow through the urine.¹⁷ Serum uric acid levels differ by gender, with a typical range of up to 7mg/dl in men and up to 6mg/dl in women.¹⁸ Different investigations have found increased blood uric acid levels in many renal conditions, including early kidney diseases, diabetes, metabolic syndrome, hypertension, and so on.¹⁹

The goal of this study was to see if there was a link between microalbuminuria and serum uric acid in type 2 diabetes patients.

METHODOLOGY

Between July and December 2019, a cross-sectional research was undertaken at LUMHS Jamshoro. A total of 80 diagnosed cases of type-2 diabetes (59 men and 21 females) with an age range of 40 to 60 years were recruited from medical OPDs and the LUMHS diabetic clinic as case study participants, while 80 healthy people (60 males and 20 females) in the same age range served as control subjects. This study excluded individuals with serum creatinine level more than 1.5mg/dl, macroalbuminuria, diabetic retinopathy, type 1 diabetes, hypertension, heart failure, hypothyroidism, gout, history of current fever and urinary tract infection and those on diuretic or ACE inhibitor medicines. On the spot, a blood sample of 5 cc was taken for examination of random blood glucose levels, serum uric acid, serum creatinine, and HbA1c percent, while a urine sample was taken for albumin detection. The hexokinase technique was used to assess the random blood glucose level. Serum creatinine was determined using a kit, and serum uric acid was determined using a calorimeter and the uricase enzyme technique. Albumin in urine samples is measured using an immunoturbidimetry kit. SPSS version 22 was used for the statistical analysis.

RESULTS

The relationship of several factors between the control and s=case study groups is given in Table 1. In the case study group, random blood glucose levels were significantly higher, and HbA1c percent levels were also significantly higher. In the case study group, blood creatinine and serum uric acid levels increased considerably, following the same pattern as in microalbuminuria. Micro albuminuria was found in only 11 cases of control people, whereas it was found in 49 cases of type-2 diabetes mellitus.

These findings also revealed a substantial relationship between serum uric acid, microalbuminuria, and random blood sugar levels.

Table: No: 01

Parameters under study with their significant & correlation values

Parameter	Control Group	Case Study Group	P.value	r. value
RBS (mg/dl)	135 ± 10.11	224 ± 13.52 **	< 0.001	0.30
HbA1c%	5.3 ± 0.5	9.7 ± 2.1*	< 0.05	0.26
S.Creatinine (mg/dl)	0.71 ± 0.34	1.51 ± 0.52*	<0.05	0.11
S. Uric Acid (mg/dl)	4.7 ± 0.6	6.6 ± 1.4**	<0.001	0.34
Albumin in urine (mg/gm)	10.2 ± 1.78	54 ± 10.34 *	<0.05	0.28




Figure No: 01 Serum Uric Acid Level




Figure No: 02 Albumin Level in Urine




Figure No:03 Serum Uric Acid with Association of Glycemic Control in Control & Case Study Group




Figure No:04 MicroAlbumin urea with Association of Glycemic Control in Control & Case Study Group

DISCUSSION

Diabetes and its consequences have a significant negative influence on human health. Diabetic nephropathy is a diabetic condition that damages the kidney over time, eventually leading to end-stage renal failure (ESRD). Nephropathy caused by diabetes increases the risk of kidney injury by a factor of ten.¹⁰ In diabetic nephropathy, serum creatinine levels can be used as a predictor for assessing kidney function.²⁰ The presence of albumin in the urine is also a warning sign that kidney damage is beginning.²¹ Kidney damage in diabetics is linked to hyperglycemia for a variety of reasons, including hexosamine activation,²² an increase in intracellular reactive oxygen species, and non-enzymatic glycation of proteins, which raises renal capillary pressure and, eventually, the appearance of protein in urine.^{23,24} As a result, DM combined with micro albuminuria implies renal impairment.²⁵ This study reveals that there is strong relation with DM, hyper uricemia and micro albuminuria.

Uric acid is one of the components associated with glucose metabolism.²⁶ Uric acid levels may rise as a result of oxidative stress,²⁷ which may contribute to the development of diabetes. Another mechanism linking uric acid to the development of diabetes is that it inhibits the formation of endothelial nitric oxide, resulting in endothelial dysfunction and insulin resistance, as evidenced by the higher levels of uric acid found in DM by several researchers in their investigations.^{28, 29}

Serum uric acid levels were higher in newly diagnosed cases of diabetes, according to Anju Gill et al (2013).³⁰ According to Abbas Dehghan et al (2008)³¹, those with high blood uric acid levels are more likely to acquire type 2 diabetes. After doing study, C.K. Kramer et al. (2009)³² proposed that hyperuricemia can induce glucose intolerance. Our findings backed with the findings of RanparvarAlamdarm et al.(2006)³³, who found that blood glucose levels are directly

related to micro albuminuria. Our findings are backed up by ShokoofehBonakdaran et al.(2011)³⁴, who found a substantial link between hyperuricemia and microalbuminuria. The development of nephropathy in diabetes mellitus persistently declines the renal functions which can be irreversible in nature. Treatments available not for the restorations of renal damage which has been done but for the back up support for further renal damage which can lead ESRD.

UNDER PEER REVIEW

CONCLUSION

This study revealed that elevated level of serum uric acid along microalbuminuria in the patients of type-2 diabetes mellitus may be the one of the predicting and early indicator for the development of diabetic nephropathy in the diabetic population.

UNDER PEER REVIEW

REFERENCES

1. Vieira R, Souto SB, Sánchez-López E, López Machado A, Severino P, Jose S, Santini A, Fortuna A, García ML, Silva AM, Souto EB. Sugar-lowering drugs for type 2 diabetes mellitus and metabolic syndrome—review of classical and new compounds: part-I. *Pharmaceuticals*. 2019 Dec;12(4):152.
2. Zheng Y, Ley SH, Hu FB. Global aetiology and epidemiology of type 2 diabetes mellitus and its complications. *Nature Reviews Endocrinology*. 2018 Feb;14(2):88-98.
3. Halajur U, Setiawan I. The Relationship of Knowledge With The Self-Management of Diabetes Clients Diabetes Mellitus. *International Journal of Science, Technology & Management*. 2021 Jan 27;2(1):358-62.
4. Nawaz MS, Nawaz MS, Shah KU, ul Mustafa Z, Ahmed A, Ahmed HS, Khan TM, Bukhsh A, Ali S. Prevalence and determinants of restless leg syndrome in type 2 diabetes mellitus (T2DM) in Pakistan. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2021 Mar 1;15(2): 525-8.
5. Omorogiwa A, Oaikhena GA, Okioya P, Akubueze D, Owobu E, Enahoro I, Okoduwa O, Otoghile T. Diabetes mellitus: prevalence amongst university staff in southern Nigeria and attitude towards routine glycemic/glucosuric checkup. *International Journal of Biomedical and Health Sciences*. 2021 Jun 3;6(1).
6. Balwan WK, Saba N, Rasool N. A SYSTEMATIC REVIEW OF OBESITY—AN INVITED DISEASE. *Journal of Natural Remedies*. 2021;21(1):2.
7. Fareed M, Salam N, Khoja AT, Mahmoud AM, Ahamed M. Life style related risk factors of type 2 diabetes mellitus and its increased prevalence in Saudi Arabia: A brief review. *International Journal of Medical Research & Health Sciences*. 2017;6(3):125-32.
8. Wei TT, Yang LT, Guo F, Tao SB, Cheng L, Huang RS, Ma L, Fu P. Activation of GPR120 in podocytes ameliorates kidney fibrosis and inflammation in diabetic nephropathy. *ActaPharmacologicaSinica*. 2021 Feb;42(2):252-63.
9. Kebede SA, Tusa BS, Weldesenbet AB, Tessema ZT, Ayele TA. Incidence of Diabetic Nephropathy and Its Predictors among Type 2 Diabetes Mellitus Patients at University of Gondar Comprehensive Specialized Hospital, Northwest Ethiopia. *Journal of Nutrition and Metabolism*. 2021 Aug 27;2021.

10. El Helaly RM, Elzehery RR, El-Emam OA, El Domiaty HA, Elbohy WR, Aboelenin HM, Salem NA. Genetic association between interleukin-10 gene rs1518111 and rs3021094 polymorphisms and risk of type 1 diabetes and diabetic nephropathy in Egyptian children and adolescents. *Pediatric Diabetes*. 2021 Jun;22(4):567-76.
11. Kocak MZ, Aktas G, Duman TT, Atak BM, Savli H. Is Uric Acid elevation a random finding or a causative agent of diabetic nephropathy?. *Revista da Associação Médica Brasileira*. 2019 Oct 10;65:1155-60.
12. Kovačević M, Kovačević M, Prodanović V, Čančar O, Mališ S, Čančar V, Đukanović L. Analysis of risk factors for progression of diabetic nephropathy in patients with type 2 diabetes. *Srpski arhiv za celokupno lekarstvo*. 2021(00):13-.
13. Aglua I. Association of Microalbuminuria with High HbA1C levels in Melanesian Adults with Diabetes of at-least 1 Year Duration. *medRxiv*. 2021 Jan 1.
14. Ahola AJ, Harjutsalo V, Forsblom C, Pouwer F, Groop PH. Depression Is Associated With Progression of Diabetic Nephropathy in Type 1 Diabetes. *Diabetes care*. 2021 Jan 1;44(1):174-80.
15. Hu J, Xu W, Yang H, Mu L. Uric acid participating in female reproductive disorders: a review. *Reproductive Biology and Endocrinology*. 2021 Dec;19(1):1-1.
16. Leander J, Sunnåker M, Rekić D, Aksenov S, Eriksson UG, Johansson S, Parkinson J. A semi-mechanistic exposure–response model to assess the effects of verinurad, a potent URAT1 inhibitor, on serum and urine uric acid in patients with hyperuricemia-associated diseases. *Journal of Pharmacokinetics and Pharmacodynamics*. 2021 Mar 17:1-7.
17. Bhattarai B, Joshi AR, Lama PY, Bataju M, Singh S. Correlation of serum uric acid with intelligence in medical and dental students of a Medical College in Kathmandu. *Journal of KIST Medical College*. 2021 Jan 31;3(1):14-7.
18. Leander J, Sunnåker M, Rekić D, Aksenov S, Eriksson UG, Johansson S, Parkinson J. A semi-mechanistic exposure–response model to assess the effects of verinurad, a potent URAT1 inhibitor, on serum and urine uric acid in patients with hyperuricemia-associated diseases. *Journal of Pharmacokinetics and Pharmacodynamics*. 2021 Mar 17:1-7.
19. Mori K, Furuhashi M, Tanaka M, Numata K, Hisasue T, Hanawa N, Koyama M, Osanami A, Higashiura Y, Inyaku M, Matsumoto M. U-shaped relationship between

serum uric acid level and decline in renal function during a 10-year period in female subjects: BOREAS-CKD2. *Hypertension Research*. 2021 Jan;44(1):107-16.

20. Liu N, Xu H, Sun Q, Yu X, Chen W, Wei H, Jiang J, Xu Y, Lu W. The Role of Oxidative Stress in Hyperuricemia and Xanthine Oxidoreductase (XOR) Inhibitors. *Oxidative Medicine and Cellular Longevity*. 2021 Mar 26;2021.
21. Jaaban M, Zetoune AB, Hesenow S, Hessenow R. Neutrophil-lymphocyte ratio and platelet-lymphocyte ratio as novel risk markers for diabetic nephropathy in patients with type 2 diabetes. *Heliyon*. 2021 Jul 1;7(7):e07564.
22. Iacobini C, Vitale M, Pesce C, Pugliese G, Menini S. Diabetic complications and oxidative stress: A 20-year voyage back in time and back to the future. *Antioxidants*. 2021 May;10(5):727.
23. Khater SI, Mohamed AA, Arisha AH, Ebraheim LL, El-Mandrawy SA, Nassan MA, Mohammed AT, Abdo SA. Stabilized-chitosan selenium nanoparticles efficiently reduce renal tissue injury and regulate the expression pattern of aldose reductase in the diabetic-nephropathy rat model. *Life Sciences*. 2021 May 31:119674.
24. Babel RA, Dandekar MP. A review on cellular and molecular mechanisms linked to the development of diabetes complications. *Current Diabetes Reviews*. 2021 May 1;17(4):457-73.
25. Feng C, Wu M, Chen Z, Yu X, Nie Z, Zhao Y, Bao B. Effect of SGLT2 inhibitor on renal function in patients with type 2 diabetes mellitus: a systematic review and meta-analysis of randomized controlled trials. *International Urology and Nephrology*. 2019 Apr;51(4):655-69.
26. Ya BL, Liu Q, Li HF, Cheng HJ, Yu T, Chen L, Wang Y, Yuan LL, Li WJ, Liu WY, Bai B. Uric acid protects against focal cerebral ischemia/reperfusion-induced oxidative stress via activating Nrf2 and regulating neurotrophic factor expression. *Oxidative Medicine and Cellular Longevity*. 2018 Oct;2018.
27. DeFronzo RA, Reeves WB, Awad AS. Pathophysiology of diabetic kidney disease: impact of SGLT2 inhibitors. *Nature Reviews Nephrology*. 2021 May;17(5):319-34.
28. Di Bonito P, Valerio G, Licenziati MR, Campana G, Del Giudice EM, et al. "Uric acid, impaired fasting glucose and impaired glucose tolerance in youth with overweight and obesity." *Nutrition, Metabolism and Cardiovascular Diseases*. 2021 Feb 8;31(2):675-80.

29. Ganesh G, Krishnamurthy N, Ashakiran S. A study of serum uric acid and its correlation with blood glucose and Egfr in type 2 diabetes mellitus patients at a rural tertiary care centre in Kolar. *J Clin Biomed Sci* 2012;2(2):79-89.
30. Gill A, Kukreja S, Malhotra N, Chhabra N. Correlation of the Serum Insulin and the Serum Uric Acid Levels with the GlycatedHaemoglobin Levels in the Patients of Type 2 Diabetes Mellitus. *Journal of Clinical and Diagnostic Research*, 2013.
31. Dehghan A., van Hoek M., Sijbrands E.J., Hofman A., Witteman J.C. High serum uric acid as a novel risk factor for type 2 diabetes. *Diabetes Care* 2008; 31(2):361-362.
32. Kramer CK, von Muhlen D, Jassal SK, et al. Serum uric acid levels improve prediction of incident type 2 diabetes in individuals with impaired fasting glucose: the Rancho Bernardo Study. *Diabetes Care*.2009 Jul;32(7):1272–1273.
33. Iranparvar A M, Aminisani N, Bashardoost B, Shamshirgaran SM, Khodamoradzadeh M, Shokrabadi M, Olomi B. Prevalence and risk factors of microalbuminuria in type 2 diabetic patients in a diabetic clinic of Ardabil-Iran. *Int J EndocrinolMetab* 2006; 4: 8-12.
34. Shokoofeh Bonakdaran,1 Maryam Hami, Mohammad TaghiShakeri, Hyperuricemia and Albuminuria in Patients With Type 2 Diabetes MellitusIranian Journal of Kidney Diseases, 5(1)January 2011IJKD 2011;5:21-4.