

ANTIFUNGAL EFFECT OF HENNA (*LAWSONIA INERMIS*) EXTRACT ON PATHOGENIC FUNGI

Abstract

The study aimed at undertaking preliminary phytochemical studies and antifungal activities of *Lawsonia inermis* leaf extracts against clinical *Candida* isolates from female patients attending Dalhatu Araf Specialist Hospital (DASH) Lafia, Nasarawa State. HVS (High Vaginal Swab) samples were collected from 185 subjects and transported to the laboratory for analysis. Microbial culture and isolations were done on Sabouraud dextrose agar (SDA), Blood agar, Potato dextrose agar (PDA) and Sabouraud dextrose broth. Identification of clinical isolates was done following standard guideline for *Candida* identification including microscopic, cultural and biochemical characteristics (sugar utilization and fermentation). Antifungal susceptibility tests of the plant extracts at different concentrations were carried out against *Candida* isolates. Distilled water and ketoconazole drug served as negative and positive control respectively. Zones of inhibitions, the minimum inhibitory concentration (MIC) and the minimum fungicidal concentration (MFC) of the extracts were determined. Data were analysed on the Minitab 16.0 software for descriptive (mean with standard error) and inferential statistics and Chi Square at 95% confidence limit. In conclusion, *L.inermis* leaf has been shown to have antifungal properties since it contained quality phytochemicals in sufficient quantity that may be explored in the synthesis of drugs against some species of *Candida*. This finding is crucial in the management and control of candidiasis in the study.

INTRODUCTION

Plants are the primary source of medicine and they provide a potent channel in the search for new drugs (Shah *et al.*, 2006). Plants and their organs house many active ingredients which are precursors in drug synthesis Therefore, medicinal plants are important in pharmacological

research and drug development (Adebolu and Oladimeji, 2005; Aguoru *et al.*, 2015a; Shah *et al.*, 2006).

L. inermis is a perennial shrub native to tropical and sub-tropical regions of Africa, Southern Asia, Northern Australia and semi-arid zones. The plant has been introduced to other parts of the world as exotic plants with high invasiveness and tolerance to extreme conditions (Aguoru *et al.*, 2015b; Rao *et al.*, 2005). These properties make the plant suitable for research and further exploitation in both cosmetic and pharmaceutical industries. One of the challenges in public health today most especially in Nigeria and other Tropical countries is the high prevalence of infections of bacterial or fungal or protozoan or helminthic origin (Adebolu and Oladimeji, 2005; Elham and Elbasheir, 2014; Hassan *et al.*, 2004; Umeh *et al.*, 2005). One of the recalcitrant fungal infections is caused by many species of *Candida* although *C. albicans* is considered the most clinically important pathogens among *Candida* species (Manolakaki *et al.*, 2010). Many *candida* species are harmless as normal commensals in humans especially on mucosal surfaces of the gastrointestinal and genital tracts but they are also opportunistic pathogens that are infectious when opportunity present itself (Kourkoumpetis *et al.*, 2011).

An antimicrobial is an agent or a substance sourced from plants, animal and even microorganisms that have the ability to kill microorganisms or pathogens by inhibit their growth (Ajayi *et al.*, 2008; Prescott *et al.*, 2002). Antimicrobial medicines can be grouped according to the microorganisms they act primarily against. For example, antibacterial are used against bacteria and antifungal are used against fungi. They can also be classified according to their function. Agents that kill microbes are called microbiocidal, while those that merely inhibit their growth are called microbiostatic (Ajayi *et al.*, 2008; Prescott *et al.*, 2002). In

1928, Alexander Fleming became the first to discover a natural antimicrobial fungus known as *Penicillium rubens*. He named the substance extracted from the fungus penicillin and in 1942 it was successfully used to treat a *Streptococcus* infection. Penicillin also proved successful in the treatment of gonorrhoea and pneumonia, which were potentially fatal to patients. Infectious diseases have continued to be a major concern for many nations all over the world, with the current increasing trends of multi-drugs resistance among emerging and re-emerging fungal pathogens to the available modern drugs or antibiotics (Prescott *et al.*, 2002).

The potential medicinal benefits of *Lawsonia inermis* have received attention in pharmaceutical biotechnology (Abdelgadir *et al.*, 2010; Aguwa, 2008; Badoni *et al.*, 2014; Erzincan, 2017). The plant is commonly known as henna, mignonette and Egyptian privet in English. In Nigeria, it is called Laali in Yoruba and Leeli in Hausa (Aguwa, 2008). The plant was reported to be used in the treatment of ulcer, anaemia, diabetes, jaundices and other ailments in India (Chauhan *et al.*, 2007). Ancient history of India describes diverse uses and appreciable role of henna in natural health medicines (Chauhan *et al.*, 2007). Henna is also well recognized as a medicinally valuable plant in other parts of the world (Borade *et al.*, 2011; Badoni *et al.*, 2014; Ali *et al.*, 2015). Varghese *et al.* (2010) described the plant as a natural dye plant of various therapeutic roles in medicine. Currently, pharmacological claims exist that *L. inermis* possesses antiviral, antibacterial, antifungal, anti-inflammatory, tuberculostatic and anticarcinogenic activities (Varghese *et al.*, 2010). In Nigeria, henna plant is commonly used among women for body beautification as part of their culture. The dye obtained from the plant may be fixed on finger nails, faces, palms, hands, legs, stomach and any other part of the body (Aguoru *et al.*, 2015b). The dye is also used in other parts of the world in terms of cosmetic purposes as traditional symbols of identity (Rao *et al.*, 2005).

Vaginal candidiasis with high prevalence rate among women has been implicated in cases of infertility, early miscarriages and damages to the female reproductive system (Kourkoumpetis *et al.*, 2011). Drugs that are effective against pathogenic *Candida* species are either expensive or being resisted by the pathogen (Cazarolli *et al.*, 2008; Kourkoumpetis *et al.*, 2011).

A major task in public health today is to search for cheap, readily available plants as sources of highly effective antifungal drugs. It is important to continuously undertake studies on the antimicrobial activities of plants in the quest for potent antimicrobial plants that can effectively combat many human pathogenic microbes including *Candida* species without any form of resistance. Therefore, this research attempted to study the antifungal efficacy of *L. inermis* leaf extracts against clinical *Candida* species isolates from female patients with cases of infertility currently undergoing treatments at the gynecology unit of DASH Hospital in Lafia, Nasarawa State, Nigeria.

METHODOLOGY

Study Area

Nasarawa State lies within the Guinea savannah region of Central Nigeria with a total land area of 27,116.8 km². This study was carried out in Lafia, Nasarawa state at Dalhatu Araf Specialist Hospital Lafia.

Study Design and Sampling

A cross sectional study design was carried out among female patients attending the Gynecology unit. This study was carried out in Lafia, Nasarawa state it of the Dalhatu Araf Specialist Hospital Lafia (DASH). The hospital was specifically chosen due to its high level of patronage

among women, referral cases from other hospitals on gynecological cases, quality service delivery and state of the earth facilities

Clinical Isolates

The stock isolates for this study were obtained from female patients attending Dalhatu Araf Specialist Hospital Lafia, Nasarawa State. Samples of vaginal discharge in HVS were collected using sterile disposable speculum. All collected samples were accompanied by data involving patients' age, sex, sample and code in addition to date of sample collection.

Leaf extraction procedures

Leaf extraction (both fresh and dried leaves) was obtained using cold maceration method as described in Umeh *et al.* (2005). Fifty (50g) of the powdered and fresh milled plant material (leaves) was weighed into clean sterile bottles separately. The weighed-out leaves plant was extracted using 250ml of water and ethanol separately in a tightly covered bottle and left for 48hours at room temperature and were filtered into sterile beakers, and filtrates collected were re-filtered using Whatmans No. 1 filter paper into sterile sample bottles. They were labeled appropriately and stored in plastic bags at -20°C for further analyses (Umeh *et al.*, 2005).

Preparation of stock drug solution (ketoconazole)

Stock drug solution (1280 mg/liter) was prepared by adding 50 ml of dimethyl sulfoxide to 64 mg of ketoconazole and was allowed to stand for 30 minutes. It was then dispensed in small amount and stored at -20°C (Warnock, 2007).

Determination of minimum inhibitory concentration (MIC) of extract

Theoretically, MIC is the lowest concentration of antimicrobial agent that inhibits growth which are determined visually (show turbidity) after a standard incubation period of 18-24h at 35°C

(Warnock, 2007). MIC was taken as measurement parameter to quantify the effects of antifungal agents. Based on the preliminary screening, the extract that revealed potent antifungal activity was further tested to determine the minimum inhibitory concentration (MIC) for each fungal sample. The MICs of these extracts was determined by Sabouraud broth micro dilution method (Warnock, 2007). Zones of inhibitions of the extracts at different concentrations were measured using the metre rule in milimetre. Inhibition (susceptibility) and resistance level of the treatment levels were noted.

Statistical Analysis Data Analysis

Data were analysed by the Minitab 16.0 software. Quantitative data were subjected to descriptive and inferential statistics. One Way ANOVA, T-tests and Chi-Square were applied appropriately at (P > 0.05) was considered significance.

Sample size

Sample size was determined using a conventional sample size calculator adopted in Emeribe *et al.* (2015) as given below:

$$n = \frac{Z^2 pq}{e^2}$$

n= sample size,

Z= the value of the z-table at 95% with confidence interval of 1.96

p=prevalence of *albicans* infection among women at 14% (0.14) (Emeribe *et al.*, 2015)

q= 1-p

d= sampling error or degree of precision at 5%

$$n = \frac{1.96^2 \times 0.14 (0.86)}{0.05^2} = \frac{3.8416 \times 0.1204}{0.0025} = \frac{0.4625}{0.0025} \quad n = 185$$

RESULT

Prevalence of pathogenic *Candida* (*C. albicans*) and the age distribution of female patients with vaginal candidiasis. A total of 72 females were infected resulting in prevalence of 38.9% out of the total sample size of 185 female patients. Among them, 28.1% were less than 20 years of age while 23.6% were between 31 and 40 years. 20 and years had high Infection (52.8%). *C. albicans* infection was associated with age group ($\chi^2 = 61.1944$, $P < 0.05$) as shown in Table 1.

Table 2 gives the results of quantitative phytochemical analysis of ethanolic leaf extracts of *L. inermis*. Tannin concentration recorded the highest value among the phytochemicals quantified with an average value of 5.21 ± 0.01 while glycoside had the lowest amount recorded with an average of 0.21 ± 0.00 . Alkaloid ranged from 2.40% to 2.46% with a mean of 2.44 ± 0.02 . Terpenes ranged from 2.15% to 2.26% with a mean of 2.21 ± 0.03 . Flavonoid ranged from 0.31% to 0.36% with a mean of 0.34 ± 0.01 . Reducing sugar had an average of 1.13 ± 0.01 ranging from 1.12% to 1.14%. Steroid and anthraquinone had average values of 0.33 ± 0.00 and 0.45 ± 0.01 respectively. Saponin ranged from 1.30% to 1.34% with a mean of 1.32 ± 0.01 . Differences recorded in the quantity of phytochemicals in the *L. inermis* leaf were significant ($F = 12121.96$, $P < 0.05$). Tannin, alkaloid and terpenes were the first three highest active ingredients while flavonoid, steroid and glycoside were the least three active ingredients present (Figure 3). The hierarchical relationship is given below:

Table 3 showed the zones of inhibition and the minimum inhibitory concentration (MIC) of fresh and dried ethanolic extracts of *L. inermis* at different concentrations against *Candida* isolates. FLEE (Fresh leaf ethanolic extract) at 500mg/ml gave the highest zone of inhibition of 30mm with a mean value of 28.7 ± 0.88 while DLEE (Dry leaf ethanolic extract) of the same concentration gave its highest zone of inhibition of 28mm with an average of 23.7 ± 2.33 inhibition. At 250mg/ml, FLEE and DLEE yielded average zones of inhibition of 20.0 ± 0.58 and 18.0 ± 0.56 respectively. Similar result was observed in the positive control (ketoconazole) which gave 15mm zone of inhibition at 200mg/ml. At 125mg/ml of leaf extract, zone of inhibition was 16.7 ± 1.20 in FLEE and 15.0 ± 1.15 in DLEE. From the results FLEE gave higher values than DLEE on the average of inhibition capacity as shown in Figure 4. Also zones of inhibition reduced as the extract concentration reduced. The minimum concentration of extract where inhibition was observed was 62.5mg/ml. Here, FLEE and DLEE gave average zone of inhibition of 7.0 ± 0.58 and 5.0 ± 1.00 respectively. No inhibition was observed at extract concentration below 62.5mg/ml in both FLEE and DLEE. Also, no inhibition was recorded under distilled water employed as negative control. No significant difference was observed in the inhibitory capacity of FLEE and DLEE as both extract types gave similar results ($T= 0.24$, $P= 0.813$, $P>0.05$). However, significant positive relationship was established between extract concentration and inhibition in FLEE ($r =+0.945$, $P<0.05$) and DLEE ($r =+0.932$, $P<0.05$). In the two types of extracts, coefficients of determination R^2 were 89.3% and 86.9% respectively.

Table 4 revealed the susceptibility and resistance of five species of *Candida* to different concentrations of the ethanolic extracts of *L. inermis*. *Candida albicans* tested and identified were 72 isolates. Extract concentration of 250mg/ml gave the highest percentage of inhibition at 27.8% and 15.6mg/ml followed by 25% inhibited. Three (3) isolates showed resistance at

7.8mg/ml with 4.2%. The species of *C. glabrata* identified and tested were 45 isolates out of which 8 isolates (17.8%) showed resistance at the lowest extract concentration. 15 (33.3%) and 10 (22.2%) of the isolates were inhibited by 250mg/ ml of the extract while 62.5mg/ml of the extracts were also inhibited respectively. A total of 30 isolates were identified as *C. Krusei*. Out of these, 19 (63.3%) showed resistance at the lowest concentrations while 250mg/ml inhibited 11 (36.7%) of the isolates. There were 24 *C. parapsilosis* identified and tested. Out of these 29.2% were inhibited by 62.5mg/ml of the extract while 20.8% were inhibited at 250mg/ml. only 8.3% of the isolates showed resistance to the extract at the lowest concentration used (7.8mg/ml). *C. tropicalis* isolates were 14 in number out of which 28.6% showed resistance at the lowest concentration used but 35.7% were inhibited by 125mg/ml of the extract.

From the 185 *Candida* species isolated, 52(29.2%) were inhibited by 250mg/ml as the highest inhibition recorded followed by 32 (17.3%) isolates inhibited by 62.5mg/ml of the extracts. The lowest concentration of 7.8mg/m resulted in 36 (19.5%) isolated which showed resistance. Using Chi square analysis, resistance and susceptibility to extracts were significantly associated with specific *Candida* species ($\chi^2 = 9.87065$, $P= 0.043$, $P<0.05$) and the extraction concentration used ($\chi^2 = 241.497$, $P= 0.000$, $P<0.05$).

Table 1: Prevalence of *Candida albicans* caused Vaginal Candidiasis and Age Distribution of Patients

Age (years)	Number	Percentage from <i>C. albicans</i> infected subjects (n=72) (%)	Percentage from total sample size (n=185) (%)
<20	13	18.1	7.03
20-30	38	52.8	20.54
31-40	17	23.6	9.19
41-50	4	5.6	2.16

>50	0	0	0
Total	72		38.92

χ^2 (age and Infection) = 61.1944, P=0.000, **P<0.05**

Table 2: Quantitative Phytochemical Analysis of *Lawsonia inermis* Ethanolic Leaf

	Alk	Flav	Reds	Glyc	Phen	Sap	Tan	Ste	Terp	Anth
Rep	2.45 ^b	0.31 ^h	1.14 ⁱ	0.21 ⁱ	1.89 ^d	1.30 ^e	5.21 ^a	0.32 ^h	2.15 ^c	0.47 ^g
1										
Rep	2.40 ^b	0.36 ^h	1.13 ^f	0.2 ⁱ	1.89 ^d	1.32 ^e	5.20 ^a	0.33 ^h	2.26 ^c	0.44 ^g
2										
Rep	2.46 ^b	0.34 ^h	1.12 ^f	0.21 ⁱ	1.93 ^d	1.34 ^e	5.23 ^a	0.33 ^h	2.22 ^c	0.44 ^g
3										
	2.44 ^b	0.34 ^h	1.13 ^f	0.21 ⁱ	1.90 ^d	1.32 ^e	5.21 ^a	0.33 ^h	2.21 ^c	0.45 ^g
Mean										
S.E	±0.02	±0.01	±0.01	±0.00	±0.01	±0.01	±0.01	±0.00	±0.03	±0.01

Means that do not share a letter are significantly different

F= 12121.96, P= 0.000 (**P<0.05**)

Legend:

Alk= Alkaloids

Flav= Flavonoids

Reds= Reducing sugar

Glyc= Glycoside

Phen= Phenol

Sap= Saponins

Tan= Tannins

Ste= Steroid

Terp= Terpenes

Anth= Anthraquinones

3.9	NI	NI	NI	NI	NI	NI	NI	NI
1.95	NI	NI	NI	NI	NI	NI	NI	NI
0.98	NI	NI	NI	NI	NI	NI	NI	NI
Distilled water	0	0	0	0.0±0.00	0	0	0	0.0±0.00

Ketoconazole

(200mg/ml)

15

15

Key:

NI = No Inhibition

FLEE= Fresh leaf ethanolic extract

DLEE= Dried leaf ethanolic extract

FLEE and DLEE: T= 0.24, P= 0.813 (P>0.05) DF = 17

FLEE: Pearson correlation (R) of Extract Concentration and Inhibition = 0.945, P = 0.000 (R²= 89.3%)

DLEE: Pearson correlation (R) of Extract Concentration and Inhibition = 0.932, P = 0.000 (R²= 86.9%)

Table 4: Measurement of Susceptibility and Resistance of *Candida* Species to Different Concentrations of *Lawsonia inermis* Ethanolic Extracts

Inhibitory diameter zone	S	S	S	S	S	S	R	
	5mm	10mm	15mm	20mm	25mm	30mm	00mm	
Concentration of extract	500	250	125	62.5	31.3	15.6	7.8	
	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	mg/ml	
Isolates	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)	
<i>C. albicans</i>	72	5 (6.9)	20 (27.8)	8 (11.1)	13 (18.1)	5 (6.9)	18 (25)	3 (4.2)
<i>C. glabrata</i>	45	0 (0.0)	15	12 (26.7)	10 (22.2)	0 (0.0)	0 (0.0)	8 (17.8)

			(33.3)					
<i>C. krusei</i>	30	0 (0.0)	11	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	19
			(36.7)					(63.3)
<i>C. parapsilosis</i>	24	3(12.5)	5 (20.8)	4 (16.7)	7 (29.2)	0 (0.0)	3(2.5)	2 (8.3)
<i>C. tropicalis</i>	14	0 (0.0)	3 (21.4)	5 (35.7)	2 (14.3)	0 (0.0)	0 (0.0)	4 (28.6)
Total	185	8(4.3)	54(29.2)	29(15.6)	32(17.3))	5(2.7)	21(11.4)	36(19.5)

)

χ^2 (Inhibition and Species) = 9.87065, P= 0.043 (P<0.05)

χ^2 (Inhibition and Extract concentration) = 241.497, P= 0.000 (P<0.05)




Figure 1: Percentage Resistance and Susceptibility of *Lawsonia inermis* Extracts against *Candida species*

DISCUSSION AND CONCLUSION

This work, *L. inermis* leaf was composed of high in tannin (>5%). This is a powerful active principle found in plants that possess amazing healing properties and those that are highly poisonous to insects as a defensive mechanism (Talib, 2011). Tannin exerts biocidal effects through diverse mechanisms possibly through microbial enzyme inactivation. In the work of Cowan (1999), condensed tannins glued to the cell walls of bacteria; prevent growth and protease activity. These are the likely mechanisms of tannin against pathogenic fungi. The antifungal activity of *Lawsonia inermis* leaf reported in this study may be attributed to the high quantity of tannin and the moderate level of alkaloid, terpene, phenol and saponin.

It is also possible that other phytochemicals that were present in low quantity in *L.inermis* leaf might possess antifungal properties regardless of their level. This finding aligns with some phytochemical studies where active ingredients such as reducing sugar, anthraquinone, flavonoid, steroid, glycoside were found in low quantity (Khan *et al.*, 2015; Mudi *et al.*, 2011). In other studies, some of these active ingredients were found in large amount in all parts of the plant (Kelmanson *et al.*, 2002). Regardless of the sources, these are active principles with known antimicrobial properties. For instance, flavonoid protects plants against predation due to their antimicrobial nature (Dixon and Steale, 1999). The marked antimicrobial property of flavonoids against a variety of bacterial and fungal pathogens is mediated by their action on the microbial cell membranes (Cazarolli *et al.*, 2008).

Results have pointed out the degree of relatedness among the active ingredients. This is important in the screening of varieties and cultivars of *L.inermis* for their phytochemicals. This approach is popular in plant studies most especially in plant breeding programme (Olasan *et al.*, 2018). In this case, the presence of one phytochemical may indicate the presence of the other within the leaf as in the case of flavonoid and terpene (+0.999), alkaloid and glycoside (+0.988), anthraquinone and terpene (+0.933), saponin and phenol (+0.866) and alkaloid and tannin (+0.849).

The present report aligns with previous investigation on the antifungal activity of *L.inermis*. In previous work for instance, ethanolic extract of the whole plant showed antifungal activity against *Candida albicans*, *Cryptococcus neoformans*, *Trichophyton mentagrophytes*, *Microsporium canis* and *Aspergillus niger* (Muhammad and Muhammad, 2005). Bark decoction of henna inhibited the activity of protopectinase and polygalactouronase enzymes present in fungi (Bhakuni *et al.*, 2000).

Both fresh and dried leaf ethanolic extracts at 500mg/ml and 250mg/ml were able to inhibit fungal isolates as high as 18-30mm zones of inhibition. The extracts were as effective as ketoconazole drug at these concentrations. Thus inhibition of fungal isolated was tied to the concentration of the extracts in either fresh or dried leaf. In this work, extract concentration explained fungal inhibition by 89%. Other unforeseen factors such as nutrient availability and other environmental factors accounted for the remaining 11% of the observed inhibition not attributed to the *L. inermis* extracts. Similar findings were reported in the antimicrobial activities of other plants where effective inhibition of pathogens was observed only at high concentrations of the extracts used (Kurek *et al.*, 2011). In the present study, the minimum fungicidal concentration (MFC) of leaf extract (prepared in either fresh or dried form) ranged from 22.7mg/ml to 47.0mg/ml. The observed significant differences in the MFC of the extracts could be attributed to the differences in the inherent resistant level of the isolates of different species with different genetic make-up.

From the results, five species of *Candida* were isolated and identified among the female patients attending gynaecology unit of Dalhatu Araf Specialist Hospital Lafia. This shows that candidiasis is still rampant and it is caused by diverse species including *Candida albicans*, *Candida tropicalis*, *Candida glabrata*, *Candida krusei* and *Candida parapsilosis*. Also, each of the species may be associated with different pathological effects in the female genital tract. Although only *C.albicans* is considered pathogenic, all the *Candida* species identified are opportunistic infectious yeasts (Erzincan, 2017). The most common species isolated was *C.albicans* (38.9%) followed by *C. glabrata* (24.3%) and *C. krusei* (16.2%). *C. parapsilosis* and *C. tropicalis* accounted for 12.9% and 7.5% respectively. Result was consistent with existing reports in literature that *C. albicans* is the most commonly isolated species in human genital tract

causing vaginal candidiasis (Manolakaki *et al.*, 2010). The prevalence of candidiasis caused in Lafia was higher than the 6.5% *C. albicans* and 7.5% Non albicans *Candida spp.* prevalence rates reported by Emeribe *et al.* (2015) among non-pregnant women in Abuja, Nigeria. In this work, results have shown that those within 20-30 years are higher risk. Similar finding was reported by Emeribe *et al.* (2015) when candidiasis was more pronounced among non-pregnant women of 20-30 years of age. In this age group, women are sexually active group and they are within the optimal state of conception and child bearing. These are those with more cases of infertility at DASH Lafia. From this work, high prevalence of approximately 39% of pathogenic vaginal candidiasis might be responsible for the inability of the affected patients to conceive. Moreover, other non pathogenic opportunistic *Candida* species with high prevalence (such as *C.glabrata* and *C. krusei*) might become pathogenic in immune-compromised female patients. Results are consistent with previous claims that there is an increasing global prevalence of infections caused by various *Candida* species (Spivak *et al.*, 2017).

The five species of *Candida* isolated among 185 subjects may pose serious epidemiological consequence because cases of synergistic interaction among pathogenic and non-pathogenic *Candida* species are possible in female genital tract. This condition may present serious opportunistic infections. Synergism may also make candidiasis difficult to treat because the organisms are likely to escape drug therapy through enhanced resistance. Differences in the genetic make-up of the species may impact on drug resistance. This view is widely upheld in epidemiological studies of many infectious diseases caused by multiple species of a genus (Mishra *et al.*, 2011). Medically important species of fungi such as *Candida* species have been reported to have multiple drug resistance and this has led to the search for plants that can effectively combat fungal infections including candidiasis (Mishra *et al.*, 2011). In this work, *C.*

albicans showed the highest susceptibility to *L. inermis* extract followed by *C. parapsilosis* and *C. glabrata* meanwhile many isolates of *C. krusei* and *C. tropicalis* displayed high resistance to the extract regardless of the solvent type (ethanol or petroleum ether). This shows the extent of applicability of drug synthesis from *L.inermis* leaf against the five tested species of *Candida*, although biostatic potency may be improved during drug synthesis. These findings aligned with the work of Elham and Elbasheir (2014) in the *in vitro* activity of *L. inermis* against some pathogenic fungi where inhibition was found to depend on species of fungi.

Ethical approval and Consent:

Ethical approval was obtained from the Hospital ethical committee. Informed consent was also obtained from each participant prior to commencement of the study.

REFERENCE

Abdelgadir, E.H., Ahmed, R.H., Adam, S.I.Y. and Hussein, A.M. (2010). Evaluation of toxicological activity of the aqueous extract of *Lawsonia inermis* seeds on wistar rats. *Journal of Pharmacology and Toxicology*, 5:324-333.

Adebolu, T.T. and Oladimeji, S.A. (2005). Antimicrobial activity of leaf of *Ocimum gratissimum* on selected diarrhea causing bacteria in South Western Nigeria, *African Journal of Biotechnology*, 4(7):682-690.

- Aguoru, C.U., Akombo, K and Olasan, J.O. (2015a). Qualitative and Quantitative Phytochemical Analysis of the leaf, stem bark and root of *Bombax ceiba* (Red Silk Cotton Tree) in North Central Nigeria. *International Journal of Sciences*, 4(5): 1-5.
- Aguoru, C.U., Okpe, F. O and Olasan, J.O. (2015b). Comparative Staining Efficacy of Safranin and *Lawsonia inermis* L. Aqueous Ethanolic Leaf Extract on Epidermal Cells of *Allium cepa* L. *International Journal of Current Research in Biosciences and Plant Biology*, 2(4): 108-111.
- Aguwa, C. N. (2008). Toxic effects of the methanolic extract of *Lawsonia inermis* roots. *International Journal of Crude Drug Research*, 10:15-10
- Ajayi, I. A. Jonathan, S. G. Adewuyi, A. and Oderinde, R. A. (2008). Antimicrobial screening of the essential oil of some herbal plants from Western Nigeria. *World Applied Sciences Journal*. 3(1): 79-81.
- Akinpelu, D. A. and Onakaya, T. M. (2006). Antimicrobial activities of medicinal plants used in folkore remedies in South-Western Nigeria. *African Journal of Biotechnology*. 5 (11): 1078-1081.
- Al-Ani, F. K. Younes, F.A, and Al-Rawashdeh, F. O (2002). Ringworm Infection in Cattle and Horses in Jordan. *Acta of Veterinary Science*, 71:55-60.
- Ali, B.H., Bashir, A.K. and Tanira, M.O.M. (1995). Anti-inflammatory, antipyretic and analgesic effects of *Lawsonia inermis* L (Henna) in rats. *Pharmacology*, 51: 356-363.

Alper, J. (1998). Effort to combat microbial resistance lags. *American Society of Microbiology News* 64:440–441.

Al-Rubiay, K. K., Jaber, N. N., Al-Mhaawe, B. H. and Alrubaiy, L. K. (2008). Antimicrobial efficacy of Henna extracts. *Oman Medical Journal*, 23(4):1-4.

Alviano, D.S. and Alviano, C. S. (2009). Plant extracts: Search for new alternatives to treat microbial diseases. *Current Pharmaceutical Biotechnology Journal*, 10: 106-121.

Ashnagar, A. and Shiri, A. (2011). Isolation and characterization of 2-hydroxy 1- 1,4 naphthoquinone (lawsone) from the powdered leaves of henna plant marketed in Ahwaz city of Iran. *International Journal of Chemical Research*, 3: 1941-1944.

Asi, M. N. and Hossein, H. (2008). Review of pharmacological effects of *Glycyrrhiza* sp. and its bioactive compounds. *Phytotherapy Research*.22 (6): 709–24.

Badoni, S.R., Semwal, D.K., Combrink, S., Cartwright-Jones, C. and Viljoen, A. (2014). *Lawsonia inermis* L.: ethnobotanical, phytochemical and pharmacological aspects. *Journal of Ethnopharmacology*, 155 (1): 80-103.

Bandow, J.E., Brotz, H., Leichert, L. I. O., Labischinski, H. and Hecker, M. (2003). Proteomic approach to understanding antibiotic action. *Antimicrobial Agents Chemotherapy*, 47: 948-955.

- Bandyo, U. Biswas, K. Chattopadhyay, I. and Banerjee, R.K. (2002). Biological Activities and medicinal properties of neem (*Azadirachta indica*). *Current Science*, 82(11):1336-1345.
- Barker, J. (2006). Antibacterial drug discovery and structure-based design. *Drug Discovery Today*. 11: 391-404.
- Bartroli, J., Turmo, E., Alguero, M., Boncompte, E. and Vericat, M. L. (1998). New azole antifungals. 3. Synthesis and antifungal activity of 3-substituted-4 (3H)-quinazolinones. *Journal of Medicinal Chemistry*. 41: 1869-1882.
- Bhakuni, D.S, Dhar, M.I, Dhar, M.M., Dhawan, B.N., Gupta, B. and Srimal, R.C. (2000), Screening of Indian plants for biological activity, *Indian Journal of Experimental Biology*, 15: 91-97.
- Bussmann, R.W., Gilbreath, G.G., Solio, J. Lutura, M., Lutuluo, R., Kunguru, K., Wood, N. and Mathenge S. G. (2006). Plant use of the Maasai of Sekenani Valley, Maasai Mara Kenya. *Journal of Ethnobiology and Ethnomedicine*, 2: 22.
- Cazarolli, L. H., Zanatta, L. and Alberton, E. H. (2008). Flavonoids: prospective drug candidates. *Mini Reviews in Medicinal Chemistry*, 8 (13): 1429–1440.
- Chauhan, M. G. and Pillai, A. P. G. (2007). *Lawsonia inermis*: Microscopic profile of powdered drugs used in Indian system of medicine. *Jamnagar Gujrat*, 2: 84-85.
- Cheesebrough, M. (2002). District Laboratory Practice in Tropical Countries. Tropical Health Technology Publishers, Great Britain. 40-56pp.

Chopra, R. N., Nayer, S. L., and Chopra, I. C. (1956). Glossary of India Medicinal Plants, CSIR Publications, New Delhi, pp 151.

Cowan, M. M. (1999). Plant products as antimicrobial agents. *Clinical Microbiology and Review*. 12 (4): 564-582.

Cuong, N. X., Nguyen, N. X., Thao, N. P., Nam, N. H. and Kim, Y. H. (2010). Inhibitors of osteoclastogenesis from *Lawsonia inermis* leaves. *Bioorganic and Medicinal Chemistry Letters*, 20(6):4782-4784.

Dasgupta, T., A.R. Rao and P.K. Yadava, (2003). Modulatory effect of henna leaf (*Lawsonia inermis*) on drug metabolising phase I and phase II enzymes, antioxidant enzymes, lipid peroxidation and chemically induced skin and stomach papillomagenesis in mice. *Molecular Cell Biochemistry*, 245: 11-22.

Davidson, P. M. and Naidu, A. S. (2000). Phytophenols in Natural Food. Antimicrobial Systems, pp. 265–293, CRC Press.

Deborah, D.E, Bhavani, R.S., Kumar, R. Y and Bharath, K A (2011), Formulation and Evaluation of *Lawsonia inermis* paste. *International Journal of Research in Pharmaceutical and Biomedical science*, 2 (2): 687-690.

Di-Domenico, B., (1999). Novel antifungal drugs. *Current Opinion in Microbiology*, 2(5): 509-515.

Dixon, R. A. and Steele, C. L. (1999). Flavonoids and isoflavonoids, a gold mine for metabolic engineering. *Trends in Plant Science*, 4 (10): 394–400.

- Endrini, S., Rahmat, A., Ismail, P., Taufiq-Yap, Y. H. and Othamn, F. (2012). Effects of Henna (*Lawsonia inermis*) on the Apoptotic pathway of human liver carcinoma cell lines. *Journal of Biological Chemistry*, 7(3):321
- Elham A and Elbasheir A.M (2014). Invitro activity of *Lawsonia Inermis* (Henna) against some pathogenic fungi. *Journal of Mycolog*, 13: 18-23.
- Emeribe, A., Abdullahi, N.I., Onyia, J. and Ifunaya, , A.L. (2015). Prevalence of vulvovaginal candidiasis among non-pregnant women attending a tertiary health care facility in Abuja, Nigeria. *Research and Report in Tropical Medicine*, 6: 37-42
- Erzincan, U. (2017). Antifungal Activity of *Lawsonia inermis* (Henna) against Clinical *Candida* isolates. *Journal of Science and Technology*, 10(2): 196-202.
- Faizi, S. and Ali, M. (1999). Shamimin: A new Flavonol C-glycoside from leaves of *Bombax ceiba*. *Planta Medicine*, 65: 383–385
- Ferrazzano, G.F., Amato, I., Ingenito, A., Zarrelli, A., Pinto, G. and Pollio, A. (2011). Plant polyphenols and their anti-cariogenic properties: A review *Molecules*, 16 (2): 1486-1507.
- Ficker, C. E., Arnason, J. T., Vindas, P. S., Alvarez, L. P. and Akpagana, K. (2003). Inhibition of human pathogenic fungi by ethnobotanically selected plant extracts. *Mycoses*, 46 (1-2): 29-37.

- Gorji, A., (2003). Pharmacological treatment of headache using traditional Persian medicine. *Trends Pharmacological Science*, 24: 331-334.
- Gull, I., Sohail, M., Shahbaz, M.A. and Muhammed, A.A. (2013), Phytochemical, toxicological and antimicrobial evaluation of *Lawsonia inermis* extracts against clinical isolates of pathogenic bacteria and fungi. *Annals of Clinical Microbiology*, 12:36-42.
- Gupta, S., Ali, M., Pillai, K. K. and Alam, M. S. (1993). Evaluation of anti-inflammatory activity of some constituents of *Lawsonia inermis*. *Fitoterapia*, 64(4):365-366.
- Habbal, O. A., Al-Jabri, A. A., El-Hag, A. H., Al-Mahrooqi, Z.H. and Al-Hashmi, N. A. (2005). In vitro antimicrobial activity of *Lawsonia inermis* Linn (Henna): A pilot study on the Omani Henna. *Saudia Medical Journal*, 26:69-72.
- Hanna, R., Maciej, J.N., Lapinsky, I and Adamowicz, L. (1998). Molecular structure and infra-red spectre of 2-hydroxy-1-4-naphthquinone: Experimental matrix isolation and theoretical Hatree-Fock and post Hatree-Fock study. *Spectrochimica Acta*, 54: 1091: 1103.
- Hassan, M.M., Oyewole, A. O., Amupitan, T. O., Abdullahi, M. S. And Okonkwo, E. M. (2004). Preliminary phytochemical and antibacterial investigation of crude extracts of the root bark of *Detarium microcopum*. *Journal of the Chemical Society of Nigeria*, 29(1): 26-29.
- Hiroshi, S. (2007). Shimizu's Textbook of Dermatology, Hokkaido University, 355pp.

Ibrahim U. J. (2014). A Progress Report on the Development of Spatial Data Infrastructure in Nasarawa State - Nigeria. Office of the Governor, Lafia, Nasarawa State, NAGIS Service Center Office.

Jagdish, C. (2010) Dermatophytoses. Medical Mycology, 1st Edition, 106-107.

Jamuna, S., Senguttuvan, K. and Krishnamoorthy, K. (2014). Phytochemical analysis and evaluation of leaf and root parts of the medicinal herb *Hypochoeris radiata* L. for in vitro antioxidant activities. *Asian Pacific Journal of Tropical Medicine*. 4: 359-367

Jiny, V.K., Silvipinya, K.S, Resmi, S and Jolly, C.I. (2010), *Lawsonia inermis*: A natural dye of various therapeutic uses a review. *Cosmeceuticals*, 5:10-16.

Kamal, M. and Jawaid, T (2010). Pharmacological activities of *Lawsonia inermis* Linn: A review. *International Journal of Biomedical Research*, 2:62-68.

Kelmanson, J. F., Jager, A. K. and Staden, J. V. (2002). Zulu Medicinal plants with antibacterial activity. *Journal of Ethnopharmacology*, 69:241-246.

Khan, M. M., Ali, A., Jain, D.C., Bhakuni, R. S., Zaim, M., Thakur, R. S. (2015), Occurrence of some antiviral sterols in *Artemisia annua*. *Plant Science Journal*, 75(2):161-165.

Kourkoumpetis, T.K., Velmahos, G.C., Ziakas, P.D. and Malolakaki, D. (2011). The effect of cumulative length of hospital stay on the antifungal resistance of *Candida* strains isolated from critically ill surgical patients. *Mycopathologia*, 171 (2): 85-91

Kurek, A., Grudniak, A. M., Kraczkiewicz-Dowjat, A. and Wolska, K. I. (2011). New antibacterial therapeutics and strategies. *Polish Journal of Microbiology*, 60 (1): 3–12.

- Lanternier, F. Pathan, S. Vincent, Q. B and Liu, L. (2013). Deep dermatophytosis and inherited CARD9 deficiency. *The New England Journal of Medicine*; 369(18): 1704-1714.
- Larone, D.H. (2000) medically important fungi: A guide to identification,, American Society for Microbiology. 3rd Edition. 402pp
- Lavhate, M. S. and Mishra, S.H. (2007). A review: nutritional and therapeutic potential of *Ailanthus* species. *Pharmacognosy Review*, 1(1):105-113.
- Lekouch, N., A. Sedki, A. Nejmeddine and S. Gamon, (2001). Lead and traditional Moroccan pharmacopoeia. *Science Total Environment.*, 280: 39-43.
- Manolakaki, D., Velmahos, G. and Kourkoumpetis, T. (2010). Candida infection and colonization among trauma patients. *Virulence*, 1(5): 367-375.
- Mikhaeil, B. R., Badria, F. A., Maatooq, G. T. and Mohammed, M. A. (2004). Antioxidant and immunomodulatory constituents of Henna leaves. *Zeitschrift fur Naturforsch*, 59C:468-476.
- Mishra, A., Kumar, S., Bhargava, A., Sharma, B. and Pandey, A.K. (2011). Studies on *in vitro* antioxidant and anti staphylococcal activities of some important medicinal plants. *Cellular and Molecular Biology*, 57: 16-25.
- Mishra, A.K., Mishra, A., Kehri, H. K., Sharma, B. and Pandey, A. K. (2009). Inhibitory activity of Indian spice plant *Cinnamomumzeylanicum* extracts against *Alternaria solani* and *Curvularia lunata*, the pathogenic dematiaceous moulds. *Annals of Clinical Microbiology Antimicrobials*, 8: 9 doi: 10.1186/1476-0711-8-9.

- Mudi, S. Y., Ibrahim, H. and Bala, M. S. (2011). Acute toxicity studies of the aqueous root extract of *Lawsonia inermis* Linn. in rats. *Journal of Medicinal Plants Research*, 35: 5123-5126
- Muhammad, H. S. and Muhammad, S. (2005). The use of *Lawsonia inermis* in the management of burn wound infections. *African Journal of Biotechnology*, 4:934-937
- Muhsin, T.M. Aubaid, A.H and Al-Duboon AH (1997). Extracellular enzyme activities of dermatophytes and yeast isolates on solid media. *Mycoses*, 40 (11-12):465-469.
- Nadjib, M. R., Zahia, B. A., Mohammed, B., Kebir, B. and Nouredine, C. B. (2013), Antimicrobial activities of Henna extract and some synthetic naphthoquinones derivatives. *American Journal of Medical and Biological Research*, 1(1): 16-22
- Norman, R. F. (1996). Biological and Phytochemical screening of Plants; *Journal of Pharmaceutical sciences*. 55(3): 243-251.
- Odebiyi, O. O. and Sofowora, E. A. (1978). Phytochemical screening of Nigeria medicinal plants II. *Lloydia*,41: 2234-246.
- Ogbonna, C. I., Enweani, I. B. and Ogueri, S. C. (2008). The distribution of ringworm infections among Nigerian nomadics fulani herdsmen. *Mycopathologia*, 96 (1): 45-51.
- Okpanachi, O., Aguru, C.U., Azua, E.T., and Olasan, J.O. (2019). Ethnobotanical uses and people's perception of indigenous fruits and vegetable crops in Nasarawa State, Nigeria, *International Journal of Current Research in Food Science and Agriculture*, 1(1): 1-7

- Okerulu, I. O. and Ani, C. J. (2001). The phytochemical analysis and antibacterial screening of extracts of *Terracarpidium conophorum*. *Journal of Chemical Society of Nigeria*, 26(1): 53-56.
- Olasan, J.O., Aguoru, C.U., Omoigui, L.O. and Ekefan, E.J. (2018). Character Association Studies in Groundnut (*Arachis hypogaea* L.) *American Journal of Plant Sciences*, 9(7):1531-1543
- Oumeish, O.Y., 2001. The cultural and philosophical concepts of cosmetics in beauty and art through the medical history. *Clinical Dermatology.*, 19: 375-386.
- Philpot, C. M. (1999). The use of nutritional tests for the differentiation of dermatophytes. *Sabouraudia*, 15 (2):141-50.
- Pink, J., Planchon, J.S.M., Tagliarino, C., Varnes, M.E, Siegel, D. and Boothman, D.A. (2000), “NAD(P)H, Quinone oxidoreduotae activity is the principal determinant if beat-lapachone cytotoxicity”. *Journal of Biological chemistry*, 275:5410-5424.
- Ponnu, S., Santhi, D. K., Jacob, N. and Suresh, B. (2003). Safety measures with herbs. *Indian Pharmacist*, 2, pp: 9-12.
- Prescott, L., Harley, J. and Klein, D. A. (2002). *Microbiology*. 5thed. 950.
- Rippon, J. W. (2007). The pathogenic fungi and the pathogenic actinomycetes. *Medical mycology*, 2nd ed., W. B. Saunders Co., Philadelphia. p. 154-248.
- Savoia, D. (2012). Plant-derived antimicrobial compounds: alternatives to antibiotics. *Future Microbiology*, 7 (8): 979–990.

- Shah, J. S., Shah, M. B., Goswami, S. S., and Santani, D. D. (2006). Mechanism of action of antiulcer activity of bark extracts of *Manikarahexandra* against experimentally induced gastric ulcer in rats. *Pharmacognosy Magazine*, 2:40-45
- Sharma, V.K. (1990). Tuberculostatic activity of *Lawsonia inermis* Linn). *Tubercle*, 71: 293-295.
- Sidigia, I., Nyaigotti-Chacha, C. and Kanunah, M. P. (1990). Traditional medicine in Africa. East African Educational Publishers, Nairobi, 500pp.
- Spivak, E.S., Hanson, K. and Emily, S. (2017). *Candida auris*: An emerging fungal pathogen. *Journal of Clinical Microbiology*, 56(2): 35-43
- Stary, F. (1996). The Natural guide to medicinal herbs and plants. Barnes and Noble Inc., New York. 622pp.
- Talib, W.H., (2011). Anticancer and Antimicrobial Potential of Plant-Derived Natural Products. In: Phytochemicals-Bioactivities and Impact on Health. Rasooli, I., (Eds.). Intech publications, Croatia, pp: 141-158
- Thomas, PA (2003). Current Perspectives on Ophthalmic Mycoses. *Clinical Microbial and Review*,16(4):730-797.
- Umeh, E. U., Oluma, H. O. A. and Igoli, J. O. (2005). Antibacterial screening of four local plants using an indicator-based microdilution technique. *Africa Journal of Traditional, Complementary and Alternative Medicines*, 2(3): 238-243.
- Varghese, J. K., Silvipriya, K. S., Resmi, S., Jolly, C. I. (2010). *Lawsonia inermis*: A natural dye

of various therapeutic uses: A Review. *Cosmoceuticals*, 1(1): 1-5.

Warnock, D. W. Evans E. G. V and Richardson, M. D. (2007)“Methods with antifungal drugs,” in *Medical Mycology a Practical Approach*, IRL Press, Oxford, UK, Eds., chapter 11, pp. 235–247,

Warycha, M. A. Leger, M. Tzu, J. Kamino, H and Stein J. (2011) Deep dermatophytosis caused by *Trichophyton rubrum*. *Dermatology Online Journal*; 17 (10): 21.

Wawrzkievicz, K. Wolski, T and Lobarzewski, J. (2003). Screening the keratinolytic activity of dermatophytes in vitro. *Mycopathologia*. 114 (1):1-8.

Xu, R., Zhao, W., Xu, J., Shao, B. and Qin, G. (1996). Studies on bioactive saponins from Chinese medicinal plants. *Advances in Experimental Medicine and Biology*. 404: 371–382.

Zafriri, D., Ofek, I., Adar, R., Pocino, M. and Sharon, N. (1989). Inhibitory activity of cranberry juice on adherence of type 1 and type P fimbriated *Escherichia coli* to eucaryotic cells. *Antimicrobial Agents Chemotherapy*, 33: 92-98

